

Topological representations of motion groups and mapping class groups – a unified functorial construction

Martin Palmer and Arthur Soulié

22nd December 2020

Abstract

Braid groups, mapping class groups and similar groups of geometric origin typically have “wild” representation theory. One would therefore like to construct representations of these groups by topological or geometric means, in order to be able to understand them with topological or geometric tools. As one very important example, Lawrence and Bigelow have constructed families of linear representations of the classical braid groups starting from actions on the twisted homology of configuration spaces, which were then used by Bigelow and Krammer to prove the linearity of the braid groups.

We give a unified construction of such topological representations: in each dimension d , we construct a large family of representations of a category \mathcal{UD}_d whose automorphism groups contain all mapping class groups and *motion groups* in dimension d . There are three parameters that one may vary in the construction: a submanifold $Z \subset \mathbb{R}^d$ and two integers $\ell \geq 2$ and $i \geq 0$. In particular, varying the parameter ℓ leads to a “pro-nilpotent” tower of representations, of which we give several non-trivial examples in dimensions 2 and 3. The richer structure of the category \mathcal{UD}_d (beyond its automorphisms) may moreover be used to organise the representation theory of the family of groups in question.

This recovers and unifies many previously-known constructions, including those of Lawrence-Bigelow, as well as the *Long-Moody construction*, using an iterative variant of our construction. We also discuss some of the new families of representations that we obtain in dimensions 2 and 3, for the surface braid groups, loop braid groups and mapping class groups.

1. Introduction

The representation theory of *mapping class groups* and *motion groups* is very rich, and the subject of much active research – see Birman and Brendle’s survey [BB05, §4] or Margalit’s expository paper [Mar19] for instance. Their representation theory is in particular known to be *wild*, meaning (roughly) that there can be no classification system for their irreducible representations with finitely many parameters. For the braid groups on $n \geq 6$ strands, this follows from the work of Erdmann and Nakano [EN02] on the representation theory of Hecke algebras; for $n = 3$ (and thus $n = 4$ due to the surjection $\mathbf{B}_4 \twoheadrightarrow \mathbf{B}_3$) it follows from work of Krugljak and Samoilenko [KS80].

In order to understand the representation theory of these groups, we must therefore view them not just as abstract groups, but use their associated geometry and topology. Combining this philosophy (of studying these groups via their *topology*) with the philosophy of representation theory itself (of studying groups via *linear algebra*), one is led naturally to homology — more precisely, the idea of studying these groups via their actions on the homology of topological spaces naturally associated to them.

Another philosophical idea adopted in this paper is to treat simultaneously *families of groups* that belong together geometrically. Here a family of groups means a collection of groups G_i indexed by a partially-ordered set (typically the natural numbers \mathbb{N}) equipped with morphisms $G_i \rightarrow G_j$

2020 *Mathematics Subject Classification*: 18B30, 18B40, 20C07, 20J05, 55R91, 57M07, 57M10, 57R19.

Key words and phrases: Homological representations, mapping class groups, surface braid groups, loop braid groups, motion groups, Lawrence-Bigelow representations, Long-Moody construction.

whenever $i \leq j$. The idea is to require representations to respect the natural coherences between the different groups in the family: this is more meaningful, since the groups naturally arise with certain coherences between them. It should also make the representation theory a little less wild, by using more of the natural structure of the families of groups and the ensuing imposed constraints.

For the family of braid groups \mathbf{B}_n (where the inclusions are induced by adding a strand on the left), Lawrence [Law90] and Bigelow [Big01] constructed well-known families of linear representations, called the *Lawrence-Bigelow representations*, following different methods. They may be defined via actions on twisted homology groups of configuration spaces of unordered points in a marked 2-disc. The most famous of these are the family of (reduced) *Burau representations* originally introduced by Burau in [Bur35] and the family of *Lawrence-Krammer-Bigelow representations*, which Bigelow [Big01] and Krammer [Kra02] independently proved to be faithful.

Our goal is to develop a much more general construction, which:

- applies simultaneously to all mapping class groups and motion groups in a given dimension,
- respects the coherences between these groups, encoded by defining the representations on a category containing all of these groups as (normal subgroups of) its automorphism groups,
- produces a much wider family of representations,
- produces interesting representations also over non-commutative rings.

In dimension d , we consider a certain topological category \mathcal{UD}_d , whose automorphism groups are (roughly) diffeomorphism groups of d -manifolds fixing a given configuration of submanifolds in their interior. Its associated discrete category $\pi_0(\mathcal{UD}_d)$ therefore contains all mapping class groups of d -manifolds, as well as all motion groups in d -manifolds (as normal subgroups; cf. §4.3.3). Our construction produces “twisted representations” of this category, namely functors

$$\pi_0(\mathcal{UD}_d) \longrightarrow \text{Mod}_\bullet$$

to the category Mod_\bullet of right-modules over rings. This contains the category Mod_R for every ring R , and it is an important question when our representations take values in such a subcategory; we give a general criterion for this in §5.1.5. The construction depends on three parameters:

- a submanifold $Z \subseteq \mathbb{R}^d$ and a subgroup G of its mapping class group $\pi_0(\text{Diff}(Z))$,
- an integer $\ell \geq 2$,
- an integer $i \geq 0$.

Each of these parameters may be varied to obtain interesting representations. For example, the family of Lawrence-Bigelow representations depends on an integer parameter $k \geq 1$; our construction recovers this in the case when Z is a 0-dimensional manifold of size k (and $d = 2$, $\ell = 2$, $i = k$). In dimensions $d \geq 3$, moreover, it becomes interesting to take higher-dimensional submanifolds of \mathbb{R}^d for Z , in particular in the case of the *loop-braid groups* (the group of motions of an n -component unlink in \mathbb{R}^3).

The parameter $\ell \geq 2$ controls the ground ring over which the representation is defined: it is the group-ring of a group of nilpotency class at most $\ell - 1$. There are cases where the output of our construction is independent of ℓ (hence the ground ring is commutative), but there are also many interesting cases where we obtain an infinite tower of representations as $\ell \rightarrow \infty$. In particular, the family of Lawrence-Krammer-Bigelow representations is the $\ell = 2$ term of such a tower.

The parameter $i \geq 0$ controls the degree in which we take homology. In the case of the Lawrence-Bigelow representations, there is only one interesting degree in which we can take homology (the homology in other degrees being trivial). However, more generally there can be many interesting degrees in which to take homology. For example, if we consider the family of loop-braid groups and the “naive” analogue of the Lawrence-Bigelow representations (taking Z to be a 0-dimensional manifold of size k), then there are non-trivial homology groups in all degrees $k \leq i \leq 2k$. These particular representations will be studied in more detail in the sequel paper [PS21].

There are also several natural variations of our construction. First, one may change the “flavour” of (twisted) homology that we use: notably *Borel-Moore* homology rather than ordinary homology (also cohomology, compactly-supported cohomology, etc.). Second, there is also a “solvable” (as opposed to “nilpotent”) variant of our construction, where the parameter ℓ controls the degree of solvability of the group(-ring) over which the representations are defined; see §5.1.6. We focus in

this paper on the nilpotent version since the lower central series of the relevant groups is generally better-understood than their derived series. In particular, the stopping or non-stopping of the lower central series of certain “mixed” motion groups determines whether or not we obtain an infinite tower of representations as $\ell \rightarrow \infty$ (see above); this question is answered for mixed versions of classical, loop- and surface braid groups in the companion paper [DPS21], joint with Jacques Darné.

Finally, there is also a variant of our construction that may be iterated, in the sense that, instead of a single (twisted) representation, it gives an *endofunctor of the category of twisted representations*. It depends on the same parameters as above, plus one additional parameter, which we now explain. The parameters Z, G, ℓ determine a functor $a: \mathcal{C} \rightarrow \text{Grp}$ from any subcategory $\mathcal{C} \subseteq \pi_0(\mathcal{UD}_d)$ to the category of groups. The additional parameter is then a functor $\chi: a \rtimes \mathcal{C} \rightarrow \mathcal{C}$, where \rtimes is the Grothendieck construction and this functor should be thought of morally as a “stabilisation map”. The parameters Z, G, ℓ, i, χ then determine an endofunctor of the category of twisted representations of \mathcal{C} with values in Mod_\bullet . This is inspired by and recovers the *Long-Moody construction*; see §6.

In the remainder of this introduction, we give more precise details of our constructions, results and particular examples of homological representations that we construct.

Coherent families of representations. In order to deal with representations with some *compatibility conditions*, we are interested in collections of representations $\{\varrho_n: G_n \rightarrow GL_R(M_n)\}_{n \in \mathbb{N}}$ together with preferred maps $m_n: M_n \rightarrow M_{n+1}$ satisfying the property that the restriction of ϱ_{n+1} to G_n with respect to m_n is ϱ_n . Then we say that the representations $\{M_n\}_{n \in \mathbb{N}}$ form a *family of linear representations* of the groups $\{G_n\}_{n \in \mathbb{N}}$. Here we are assuming for simplicity that the family of groups is indexed by \mathbb{N} and the morphisms $G_n \rightarrow G_{n+1}$ are injective.

This notion can be encoded in a functorial way. Let \mathcal{G} be the groupoid with objects indexed by non-negative integers, with the groups $\{G_n\}_{n \in \mathbb{N}}$ as automorphism groups and with no morphisms between distinct objects. For instance, we consider the braid groupoid β to deal with braid groups and the decorated surfaces groupoid \mathcal{M}_2 for the mapping class groups of surfaces; see §4.4 and §4.5. Suppose that we have chosen a category $\mathcal{C}_{\mathcal{G}}$ containing \mathcal{G} as its underlying groupoid and with a preferred morphism $\iota_n: n \rightarrow n+1$ for each object n , satisfying $\iota_n \circ g = g \circ \iota_n$ for each $g \in G_n$. In all of the examples addressed in this paper, such a category $\mathcal{C}_{\mathcal{G}}$ is constructed through *Quillen’s bracket construction* using a monoidal structure on \mathcal{G} (or \mathcal{G} will be a module over another monoidal groupoid); see §3. We denote by Mod_R the category of right R -modules for some ring R . A functor $\mathcal{C}_{\mathcal{G}} \rightarrow \text{Mod}_R$ then gives us a family of representations of the family of groups $\{G_n\}_{n \in \mathbb{N}}$. Moreover, if $\mathcal{C}_{\mathcal{G}}$ has been constructed from \mathcal{G} by “freely adjoining” the new morphisms ι_n (as will be the case in our examples), then a functor $\mathcal{C}_{\mathcal{G}} \rightarrow \text{Mod}_R$ is *equivalent* to a family of representations of $\{G_n\}_{n \in \mathbb{N}}$, in which case it will be referred to as a *global representation*.

Functorial homological constructions. Our overall procedure for defining *global homological representations* is summarised in the diagram (1.1) below. A more elaborate version of this is described in diagram (2.10) and Definition 2.21, but the essential ideas are the same. The desired output is the diagonal functor $\mathcal{C}_{\mathcal{G}} \rightarrow \text{Mod}_R$, a (coherent) family of representations of $\{G_n\}_{n \in \mathbb{N}}$. This is constructed in three steps:

- We first construct a topologically-enriched category $\mathcal{C}_{\mathcal{G}}^t$ whose π_0 recovers $\mathcal{C}_{\mathcal{G}}$. We will always construct the category $\mathcal{C}_{\mathcal{G}}$ using Quillen’s bracket construction, so we explain in §3.1 how this construction may be lifted to topologically-enriched categories to produce an appropriate $\mathcal{C}_{\mathcal{G}}^t$. This requires a technical result about commuting Quillen’s bracket construction with π_0 (*cf.* Lemma 3.8), which depends on a fibration condition which we verify in all of our examples; see §3.3. In our examples, its morphism spaces may be identified (*cf.* Proposition 4.8) with certain embedding spaces between manifolds.

In particular, we construct (see §3.2) topologically-enriched categories \mathcal{UD}_d that are designed to contain all diffeomorphisms of d -manifolds (equipped with configurations), together with all embeddings between such manifolds that correspond to splittings into boundary connected

summands. Its associated discrete category $\pi_0(\mathcal{UD}_d)$ thus contains all mapping class groups and motion groups in dimension d .

- The key geometric step is to construct a (continuous) functor F from \mathcal{C}_G^t to Cov_Q , the category of topological spaces equipped with regular coverings with fixed deck transformation group Q . This geometric construction is the subject of §5.
- The remaining steps simply encode the idea of taking twisted homology of covering spaces. The functor Lift takes a regular covering with deck transformation group Q to the corresponding bundle of $\mathbb{Z}[Q]$ -modules, the functor $-\otimes V$ takes the fibrewise tensor product with a $(\mathbb{Z}[Q], R)$ -bimodule V (“specialising the coefficients”), producing a bundle of R -modules, and finally H_i is the twisted homology functor in degree i . See §2.3–§2.5 for more details.

$$\begin{array}{ccccccc}
\mathcal{C}_G^t & \xrightarrow{F} & \text{Cov}_Q & \xrightarrow{\text{Lift}} & \text{Top}_{\mathbb{Z}[Q]} & \xrightarrow{-\otimes V} & \text{Top}_R & \xrightarrow{H_i} & \text{Mod}_R \\
\pi_0 \downarrow & & & & & & & & \\
\mathcal{C}_G & \dashrightarrow & & & & & & &
\end{array}
\quad (1.1)$$

$L_i(F; V)$

We also consider the slightly more general setting of *twisted* representations, where groups act on the base ring R as well as on modules: we then denote the appropriate target category by Mod_R^{tw} . In this setting, the target of the functor F is the analogous larger category Cov_Q^{tw} . Even more generally, we allow the ring R and the deck transformation group Q to vary. The corresponding larger categories in this case are denoted by Mod_\bullet and Cov_\bullet ; they contain Mod_R^{tw} and Cov_Q^{tw} respectively for each R and Q . These are described in more detail in §2.2.

There are also variants of this construction for *relative homology* where we work with categories of pairs of spaces, and for *Borel-Moore homology* where we restrict to categories of locally compact spaces and proper maps. In particular, using Borel-Moore homology is especially interesting when the image of the functor F consists of configuration spaces of points in a surface; see §5.2.

Some of these representations have been defined and studied before — at least at the level of individual groups, i.e. when restricted to the individual automorphism groups of \mathcal{C}_G — and indeed one purpose of describing this general procedure for constructing homological representations is to give a *unified* description for various different representations appearing in the literature, as well as discovering new constructions by comparing representations coming from different settings in this unified context.

Constructions of representations for motion groups. The first type of groups to which we apply our construction are *motion groups*: given a closed submanifold A of the interior of a manifold M , the motion group $\text{Mot}_A(M)$ is the fundamental group of $\text{Emb}(A, \mathring{M})/\text{Diff}(A)$, the space of embeddings of A into the interior of M modulo diffeomorphisms of A . If A is orientable, we may also consider $\text{Mot}_A^+(M)$, which is the fundamental group of $\text{Emb}(A, \mathring{M})/\text{Diff}^+(A)$. Important examples of (M, A) are:

- $(\mathbb{D}^2, n \text{ points})$ — corresponding to the classical braid groups $\mathbf{B}_n \cong \text{Mot}_n(\mathbb{D}^2)$;
- $(S, n \text{ points})$ — corresponding to the braid groups $\mathbf{B}_n(S) \cong \text{Mot}_n(S)$ on a surface S ;
- (\mathbb{D}^3, U_n) , where U_n is an n -component unlink in the interior of \mathbb{D}^3 — corresponding to the *loop-braid groups* $\mathbf{LB}_n \cong \text{Mot}_{U_n}^+(\mathbb{D}^3)$ and the *extended loop-braid groups* $\mathbf{LB}'_n \cong \text{Mot}_{U_n}(\mathbb{D}^3)$.

In §5.1, we construct a functor $\mathcal{UD}_d \rightarrow \text{Cov}_\bullet$ depending on two parameters: a closed submanifold $Z \subseteq \mathbb{R}^d$ and an integer $\ell \geq 1$. (There is also a third parameter, which we are ignoring here for simplicity.) The motion group $\text{Mot}_A(M)$ is a normal subgroup of the automorphism group of (M, A) in $\pi_0(\mathcal{UD}_d)$, so diagram (1.1) applied to this functor gives a (possibly twisted) representation of $\text{Mot}_A(M)$.

To give an idea of how this functor $F: \mathcal{UD}_d \rightarrow \text{Cov}_\bullet$ is constructed, we explain how to construct $\pi_0(F): \pi_0(\mathcal{UD}_d) \rightarrow \pi_0(\text{Cov}_\bullet)$ restricted to $\text{Mot}_A(M)$. To do this, we have to give a regular covering together with an action up to homotopy of $\text{Mot}_A(M)$. The key idea is to use the fact that $\text{Mot}_A(M)$ acts up to homotopy on the space of embeddings (modulo diffeomorphisms) of Z into $\mathring{M} \setminus A$. The fundamental group of this space is the motion group $\text{Mot}_Z(M \setminus A)$, and one then has to define appropriate quotients of this group so that the $\text{Mot}_A(M)$ -action lifts to the corresponding regular

covering. To do this, we use the following diagram, which we explain below:

$$\begin{array}{ccccccc}
1 & \longrightarrow & \text{Mot}_Z(M \setminus A) & \longrightarrow & \text{Mot}_{Z \sqcup A}(M) & \longrightarrow & \text{Mot}_A(M) \longrightarrow 1 \\
& & \downarrow \phi & & \downarrow & & \downarrow \\
1 & \longrightarrow & Q & \longrightarrow & \frac{\text{Mot}_{Z \sqcup A}(M)}{\Gamma_\ell(\text{Mot}_{Z \sqcup A}(M))} & \longrightarrow & \frac{\text{Mot}_A(M)}{\Gamma_\ell(\text{Mot}_A(M))} \longrightarrow 1.
\end{array}$$

The top row consists of the split-surjection $\text{Mot}_{A \sqcup Z}(M) \rightarrow \text{Mot}_A(M)$ given by forgetting the manifold Z (we always require M to have non-empty boundary; the splitting uses this), whose kernel is $\text{Mot}_Z(M \setminus A)$. Write $\Gamma_\ell(G)$ for the ℓ -th term of the lower central series of a group G ; this explains two of the quotients, and the diagram is completed by defining Q to be the image of $\text{Mot}_Z(M \setminus A)$ in the bottom-middle group. An easy diagram chase shows that the action of $\text{Mot}_A(M)$ on $\text{Mot}_Z(M \setminus A)$ preserves $\ker(\phi)$, which is equivalent to saying that the action lifts to the corresponding regular covering. (We also have to identify the geometric action of $\text{Mot}_A(M)$ on $\text{Mot}_Z(M \setminus A)$ with the action coming from the splitting on the top row; cf. Lemma 4.25.)

The quotient group Q is often the same for a whole family of motion groups (such as the three mentioned above); we call this phenomenon *Q-stability*, and in this case the functor F restricted to this family takes values in $\text{Cov}_Q^{\text{tw}} \subset \text{Cov}_\bullet$. In the case $\ell = 2$, moreover, we show in §5.1.5 that the functor takes values in $\text{Cov}_Q \subset \text{Cov}_\bullet$, so the resulting representations are always untwisted.

If we consider all $\ell \geq 1$ simultaneously, these representations may be packaged together into a “pro-nilpotent” tower of representations of the family of motion groups; see §5.1.7. Whether or not this tower stops at some finite stage depends on the lower central series of the “mixed” motion group $\text{Mot}_{Z \sqcup A}(M)$. This is investigated for the classical, surface and loop braid groups in [DPS21].

Many well-known representations arise as particular instances of this general homological construction. First of all, for the classical braids groups, we take Z to be a set of $k \geq 1$ unordered points in the interior of \mathbb{D}^2 . Quillen’s bracket construction defines a category $\mathfrak{U}\beta$ having the braid groupoid β as its underlying groupoid.

Theorem 1.1 (see Proposition 5.28 and Corollary 5.34) *Choosing the quotients by the Γ_2 -term and taking homology in degree k , the above procedure defines functors*

$$\mathfrak{L}\mathfrak{B}_1: \mathfrak{U}\beta \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}]} \quad \text{and} \quad \mathfrak{L}\mathfrak{B}_k: \mathfrak{U}\beta \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}^2]} \quad (\text{for } k \geq 2),$$

which encodes the Lawrence-Bigelow representations [Law90; Big04]. In particular, $\mathfrak{L}\mathfrak{B}_1$ and $\mathfrak{L}\mathfrak{B}_2$ encode the reduced Burau representations Lawrence-Krammer-Bigelow representations respectively.

If $k = 2$, applying the procedure for the quotient by the Γ_ℓ -term for $\ell \geq 3$ provides twisted representations $\mathfrak{L}\mathfrak{B}_{2,\ell}$ which are not equivalent to $\mathfrak{L}\mathfrak{B}_2$. If we use Borel-Moore homology, the twisted representations of braid groups encoded by $\mathfrak{L}\mathfrak{B}_{2,\ell}^{\text{BM}}$ are faithful.

More generally, we consider the braid groups on a connected compact surface S with one boundary-component and take for Z a set of $k \geq 1$ unordered points in the interior of S . Quillen’s bracket construction provides a category $\langle \beta, \beta^S \rangle$ having the braid groups as automorphism groups. If S is orientable of genus $g \geq 1$ we denote it by $\Sigma_{g,1}$, if it is non-orientable of genus $h \geq 1$ we denote it by $\mathcal{N}_{h,1}$.

Theorem 1.2 (see Propositions 5.35 and 5.37) *Choosing the quotients by the Γ_2 -term and taking homology in degree k , the above procedure defines functors*

$$\begin{aligned}
\mathfrak{L}_k(\Sigma_{g,1}, \Gamma_2): \langle \beta, \beta^{\Sigma_{g,1}} \rangle &\longrightarrow \text{Mod}_{\mathbb{Z}[Q_{(k,2)}(\Sigma_{g,1})]} \\
\mathfrak{L}_k(\mathcal{N}_{h,1}, \Gamma_2): \langle \beta, \beta^{\mathcal{N}_{h,1}} \rangle &\longrightarrow \text{Mod}_{\mathbb{Z}[Q_{(k,2)}(\mathcal{N}_{h,1})]},
\end{aligned}$$

where $Q_{(k,2)}(\Sigma_{g,1}) \cong \mathbb{Z}^{2g} \oplus (\mathbb{Z}/2\mathbb{Z})^{d_k}$ and $Q_{(k,2)}(\mathcal{N}_{h,1}) \cong \mathbb{Z}^h \oplus (\mathbb{Z}/2\mathbb{Z})^{d_k}$, with $d_k = 1$ if $k = 1$ and $d_k = 2$ if $k \geq 2$. In the orientable case, if we take Γ_3 quotients instead, we obtain

$$\mathfrak{L}_k(\Sigma_{g,1}, \Gamma_3): \langle \beta, \beta^{\Sigma_{g,1}} \rangle \longrightarrow \text{Mod}_{\mathbb{Z}[Q_{(k,3)}(\Sigma_{g,1})]}^{\text{tw}},$$

where $Q_{(k,3)}(\Sigma_{g,1}) = \ker(\mathbf{B}_{k,n}(\Sigma_{g,1})/\Gamma_3 \rightarrow \mathbf{B}_n(\Sigma_{g,1})/\Gamma_3)$, which is independent of n for $n \geq 3$.

For orientable surfaces, the option of taking the quotients by Γ_3 is actually a reinterpretation of the work of An and Ko in [AK10] — see [BGG17] — to extend some homological representations from the classical braid groups to the surface braid groups; see §5.2.2.2.

Finally, for loop braid groups, we have two main choices for the parameter Z : one may take for Z a set of $k \geq 1$ unordered points in the interior of \mathbb{D}^3 , or else an embedded k -component unlink. In the latter case we may either consider this to be an *oriented* unlink or an *unoriented* unlink. Quillen's bracket construction provides categories $\mathfrak{UL}\beta$ and $\mathfrak{UL}\beta'$ having respectively the loop braid groups and extended loop braid groups as automorphism groups.

Theorem 1.3 (see Propositions 5.40, 5.42 and 5.43) *Choosing the quotients by the Γ_2 -term and taking homology in degree k , the above procedure defines functors*

$$\begin{array}{ll} \mathfrak{L}_1(1, \mathcal{L}\beta) : \mathfrak{UL}\beta \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}]} & \mathfrak{L}_1(1, \mathcal{L}\beta') : \mathfrak{UL}\beta' \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}/2]} \\ \mathfrak{L}_k(k, \mathcal{L}\beta) : \mathfrak{UL}\beta \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z} \oplus (\mathbb{Z}/2)^2]} & \mathfrak{L}_k(k, \mathcal{L}\beta') : \mathfrak{UL}\beta' \longrightarrow \text{Mod}_{\mathbb{Z}[(\mathbb{Z}/2)^2]} \\ \mathfrak{L}_1(U_1, \mathcal{L}\beta) : \mathfrak{UL}\beta \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z} \oplus (\mathbb{Z}/2)^2]} & \mathfrak{L}_1(U_1, \mathcal{L}\beta') : \mathfrak{UL}\beta' \longrightarrow \text{Mod}_{\mathbb{Z}[(\mathbb{Z}/2)^3]} \\ \mathfrak{L}_k(U_k, \mathcal{L}\beta) : \mathfrak{UL}\beta \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z} \oplus (\mathbb{Z}/2)^4]} & \mathfrak{L}_k(U_k, \mathcal{L}\beta') : \mathfrak{UL}\beta' \longrightarrow \text{Mod}_{\mathbb{Z}[(\mathbb{Z}/2)^5]} \\ \mathfrak{L}_1^+(U_1, \mathcal{L}\beta) : \mathfrak{UL}\beta \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}^2]} & \mathfrak{L}_1^+(U_1, \mathcal{L}\beta') : \mathfrak{UL}\beta' \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z} \oplus (\mathbb{Z}/2)]} \\ \mathfrak{L}_k^+(U_k, \mathcal{L}\beta) : \mathfrak{UL}\beta \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}^3 \oplus (\mathbb{Z}/2)]} & \mathfrak{L}_k^+(U_k, \mathcal{L}\beta') : \mathfrak{UL}\beta' \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}^2 \oplus (\mathbb{Z}/2)^2]}, \end{array}$$

where $k \geq 2$.

Moreover, using non-stopping results for lower central series of *partitioned tripartite welded braid groups* from [DPS21], we obtain *infinite towers of representations* when we consider all $\ell \geq 2$ in the cases $Z \in \{2, U_2, U_3\}$ (that is, a configuration of two points or an unlink with either two or three components).

In §7, we compute explicitly the matrices of the representations encoded by the top two functors, $\mathfrak{L}_1(1, \mathcal{L}\beta)$ and $\mathfrak{L}_1(1, \mathcal{L}\beta')$. These extend the *reduced Burau representations* of the classical braid groups to \mathbf{LB}_n and \mathbf{LB}'_n respectively. The extension to \mathbf{LB}_n is straightforward, and was already introduced by Vershinin in [Ver01, §4] by assigning explicit matrices to generators. The extension to \mathbf{LB}'_n is more indirect: we must first reduce the ground ring $\mathbb{Z}[\mathbb{Z}]$ to $R = \mathbb{Z}[\mathbb{Z}/2]$, then extend the underlying R -module R^{n-1} of the reduced Burau representation to a larger, non-free R -module and then extend the action of \mathbf{LB}_n on R^{n-1} to an action of \mathbf{LB}'_n on this larger module. This extension (in particular the matrices defining it) does not seem algebraically obvious, but it arises naturally as part of our general topological construction. The variant of our construction using *reduced* homology also defines extensions of the *unreduced* Burau representations to \mathbf{LB}_n and \mathbf{LB}'_n ; these are more straightforward again. See §7 for the full details and calculations.

Constructions of representations for mapping class groups. The second type of groups to which we apply our construction are *mapping class groups* of manifolds. For a smooth manifold M , the mapping class group $\text{MCG}(M)$ of M is the group of isotopy classes of diffeomorphisms of M fixing its boundary pointwise. More generally, if Z is a closed submanifold of M , we write $\text{MCG}(M, Z)$ for the group of isotopy classes of M fixing its boundary pointwise and sending Z onto itself. We will focus on the case where M is a compact, connected surface with one boundary-component. Quillen's bracket construction provides suitable categories \mathfrak{UM}_2^+ and \mathfrak{UM}_2^- having the mapping class groups of orientable surfaces $\{\Sigma_{g,1}\}_{g \geq 0}$ and non-orientable surfaces together with the disc $\{\mathcal{N}_{h,1}\}_{h \geq 0}$ respectively as automorphism groups. (These are subcategories of $\pi_0(\mathfrak{UD}_2)$.)

We consider two ways to define functors $F : \mathfrak{UD}_d \rightarrow \text{Cov}_\bullet$, which will then give us homological representations of \mathfrak{UM}_2^+ and \mathfrak{UM}_2^- via diagram (1.1). The first one is the construction given in §5.1, which we have already described above in the setting of motion groups. The idea for mapping class groups is similar, using the natural action (up to homotopy) of $\text{MCG}(M)$ on the space of embeddings modulo diffeomorphisms of Z into the interior of M , whose fundamental group is the motion group $\text{Mot}_Z(M)$. We then take a characteristic quotient of this group, typically induced by its lower central series, to lift this action to the corresponding covering space.

In particular, for orientable surfaces, this principle recovers the following families of representations:

Theorem 1.4 (see §5.4.1.1 and Theorem 5.51) *Taking Z to be a single point and using reduced homology in degree 1, choosing the quotients by the Γ_ℓ -term for each $\ell \geq 1$, the above procedure defines a global homological functor $\mathfrak{Mag}_\ell: \mathfrak{UM}_2^+ \rightarrow \text{Mod}_\bullet$ encoding the level- ℓ Magnus representations introduced by Suzuki [Suz05, §3].*

Taking Z to be a set of $k \geq 1$ ordered points and removing a puncture from the boundary of each surface, taking Borel-Moore homology in degree k and choosing the quotients by the Γ_1 -term, the above procedure defines a global homological functor $\mathfrak{Mot}_k: \mathfrak{UM}_2^+ \rightarrow \text{Mod}_\bullet$ encoding the Moriyama representations introduced by Moriyama [Mor07].

Taking Z to be a set of $k \geq 1$ unordered points, taking homology in degree k and choosing the quotients by the Γ_3 -term, the above procedure defines a global homological functor $\mathfrak{Ban}_k: \mathfrak{UM}_2^+ \rightarrow \text{Mod}_\bullet$. Restricting to the Chillingworth subgroups of mapping class groups, the induced representation for the surface $\Sigma_{g,1}$ becomes an untwisted representation over the group-ring of $\mathbf{B}_k(\Sigma_{g,1})/\Gamma_3$.

The name of the functor \mathfrak{Ban}_k follows from the suggestion of [BGG11] that the ideas developed to construct that functor were already explored by Christian Blanchet but have not been published.

However, the kind of construction of the above theorem is not optimal: as is highlighted in the above results, it typically only gives *twisted* representations. One has to restrict to smaller subgroups of mapping class groups so as to get untwisted representations. For this reason, we give a second construction of a functor $F: \mathfrak{UD}_d \rightarrow \text{Cov}_\bullet$ in §5.3 that is better adapted to mapping class groups in the sense that it more often gives *untwisted* representations when we plug it into diagram (1.1). (The precise meaning of this is given by Proposition 5.48.)

The idea is again to use the natural action (up to homotopy) of $\text{MCG}(M)$ on the space of embeddings modulo diffeomorphisms of Z into the interior of M , whose fundamental group is $\text{Mot}_Z(M)$. However, instead of directly taking lower central quotients of this group to choose coverings, we instead construct a 6-term diagram, similar to the one for motion groups, and choose the quotient ϕ indirectly via the lower central quotients of the bigger mapping class group $\text{MCG}(M, Z)$:

$$\begin{array}{ccccccc}
1 & \longrightarrow & \text{Mot}_Z(M) & \longrightarrow & \text{MCG}(M, Z) & \longrightarrow & \text{MCG}(M) \longrightarrow 1 \\
& & \downarrow \phi & & \downarrow & & \downarrow \\
1 & \longrightarrow & Q & \longrightarrow & \frac{\text{MCG}(M, Z)}{\Gamma_\ell(\text{MCG}(M, Z))} & \longrightarrow & \frac{\text{MCG}(M)}{\Gamma_\ell(\text{MCG}(M))} \longrightarrow 1.
\end{array}$$

Here, the top row consists of the split surjection $\text{MCG}(M, Z) \twoheadrightarrow \text{MCG}(M)$ given by forgetting the condition that diffeomorphisms must send Z onto itself (as before, we always require M to have non-empty boundary; the splitting uses this assumption), whose kernel may be identified with $\text{Mot}_Z(M)$.

This second method produces representations of mapping class groups that appear to be new, as far as the authors are aware. For instance, when $M = S$ is a surface, we take for Z a set of $k \geq 1$ unordered points and define:

Theorem 1.5 (see Propositions 5.54 and 5.55) *Choosing the quotients by the Γ_2 -term and taking homology in degree k , the above procedure defines global homological functors:*

- $\mathfrak{L}_k(\Gamma_{-,1}): \mathfrak{UM}_2^+ \rightarrow \text{Mod}_{\mathbb{Z}[Q_k]}$ where Q_1 is trivial and $Q_k = \mathbb{Z}/2\mathbb{Z}$ for $k \geq 2$.
- $\mathfrak{L}_k(\mathcal{N}_{-,1}): \mathfrak{UM}_2^- \rightarrow \text{Mod}_{\mathbb{Z}[Q_k]}$ where $Q_k = (\mathbb{Z}/2\mathbb{Z})^2$ for $k \geq 3$.

Iterative constructions. Long and Moody [Lon94] have introduced machinery which constructs a representation of \mathbf{B}_n from a representation of \mathbf{B}_{n+1} . For instance it recovers the Burau representation [Bur35] from a one-dimensional representation. This method and its variants have been studied with a functorial point of view in [Sou19b] for braid groups and then generalised in [Sou19a] for general families of groups.

Our construction summarised in diagram (1.1) may also be automated and turned into machinery detailed in diagram (2.15). Namely, for a family of groups $\{G_n\}_{n \in \mathbb{N}}$ with an appropriate category \mathcal{C}_G as above, suppose that we have a continuous functor $F: \mathcal{C}_G \rightarrow \mathbf{Cov}_\bullet$ (for example, constructed as in §5.1 or §5.3). Let $r: \mathbf{Cov}_\bullet \rightarrow \mathbf{Grp}$ be the functor that sends a covering to its deck transformation group and suppose that we also have a continuous functor $\chi: (r \circ F) \times \mathcal{C}_G \rightarrow \mathcal{C}_G$, where \times is the Grothendieck construction (cf. Definition 2.28). Writing $\mathbf{Fct}(\mathcal{C}, \mathcal{D})$ for the category of functors $\mathcal{C} \rightarrow \mathcal{D}$, we have:

Theorem 1.6 (see Propositions 2.33, 6.4 and Theorem 6.2) *For each $i \geq 0$, the above data defines an endofunctor $\Lambda_i(F; \chi): \mathbf{Fct}(\mathcal{C}_G, \mathbf{Mod}_\bullet) \rightarrow \mathbf{Fct}(\mathcal{C}_G, \mathbf{Mod}_\bullet)$ such that, for any $V: \mathcal{C}_G \rightarrow \mathbf{Mod}_\bullet$,*

$$\Lambda_i(F; \chi)(V) = L_i(F; V_{r \circ F}^\chi): \mathcal{C}_G \rightarrow \mathbf{Mod}_\bullet$$

where $V_{r \circ F}^\chi$ is a certain canonical functor induced from V . The endofunctor $\Lambda_i(F; \chi)$ is called the homological representations iterative functor associated to F and χ .

For $i = 1$, if $r \circ F: \mathcal{C}_G \rightarrow \mathbf{Grp}$ takes values in the full subcategory of free groups, then $\Lambda_1(F; \chi)$ recovers a generalised Long-Moody functor introduced in [Sou19a].

In particular, we prove in §6.2 that any family of low-dimensional complex irreducible representations of braid groups is encoded by a global homological representation using this iterative procedure.

Perspectives. A natural question for the representations arising from the above constructions is to determine which of them are *irreducible*, which are *indecomposable* and how they decompose if they are decomposable. In addition, keeping our functorial point of view, notions of polynomiality may be introduced for functors from the discrete categories considered here to module categories: we may investigate whether the above functorial representations are *polynomial*. These questions are beyond the scope of the current paper but will be addressed in the sequel paper [PS21].

Furthermore, among the underlying prospects motivating our study stands the question of which motion groups and mapping class groups are *linear* — in the sense that they act faithfully on a finite-dimensional vector space. Indeed, this question remains wide open in the vast majority of cases. Classical braid groups provide a famous positive answer to this question by Bigelow and Krammer [Big01; Kra02], but there are also families with a negative answer, such as the automorphisms of free groups — which may be viewed mapping class groups of certain 3-manifolds — by Formanek and Procesi [FP92]. The key representations that have provided positive answers have all been homological representations. This suggests that a systematic treatment of homological constructions that work for all motion groups and all mapping class groups is an important and natural avenue for future investigations on the linearity of these families of groups.

Acknowledgements. The authors would like to thank Paolo Bellingeri, Tara Brendle, Oscar Randal-Williams, Antoine Touzé, Christine Vespa and Emmanuel Wagner for illuminating discussions and questions. In particular, they thank Antoine Touzé for pointing out the reference [EN02]. They would also like to thank Oscar Randal-Williams for inviting the first author to the University of Cambridge in November 2019, where significant progress on the present article was made.

Outline. The paper is organised as follows. In §2, we introduce in detail the various tools to define general homological functors as summarised in the diagram (1.1), *assuming given* a functor F with values in covering spaces. In §3–§4, we introduce the appropriate categorical framework to deal with all motion groups and mapping class groups simultaneously. In §5.1 and §5.3, we give two general topological constructions of functors F taking values in covering spaces — combined with all of the previous sections, this completes our general construction of *global homological representations* of motion groups and mapping class groups. We then apply this construction to motion groups (in §5.2) and mapping class groups (in §5.4) in dimensions 2 and 3. We study the general procedure to iteratively construct homological functors in §6. Finally, in §7 we focus in a more concrete way on Burau representations of the loop-braid groups, with explicit calculations.

Contents

§1. Introduction	1
§2. The general construction	9
§2.1 Overview	10
§2.2 Categories	11
§2.3 From bicoverings to bundles of bimodules	13
§2.4 Fibrewise tensor product	15
§2.5 Twisted homology	15
§2.6 The general construction	16
§2.7 An iterative version	19
§3. Topological categories of decorated manifolds	21
§3.1 A topological enrichment of the Quillen bracket construction	21
§3.2 Topological groupoids of decorated manifolds	25
§3.3 The Serre fibration condition	28
§3.4 Semi-monoidal categories and semicategories	32
§4. Topological categories for families of groups	33
§4.1 Braidings and symmetries	33
§4.2 Quillen bracket categories of manifolds	34
§4.3 Split short exact sequences	36
§4.4 Mapping class groups of surfaces	40
§4.5 Surface braid groups	40
§4.6 Loop braid groups	41
§5. Global homological representations	43
§5.1 Global functors for motion groups	43
§5.2 Applications for motion groups	55
§5.3 Global functors for mapping class groups	63
§5.4 Applications for mapping class groups of surfaces	65
§6. Iterative constructions	70
§6.1 Connection to the Long-Moody functors	70
§6.2 Recovering the Tong-Yang-Ma representations	71
§7. Reduced Burau representations for loop braid groups	72
References	78

2. The general construction

As explained in the Introduction, the construction of global homological representations consists of two main parts:

1. constructing continuous functors $\mathfrak{M}\mathcal{D}_d \rightarrow \bullet\text{Cov}_\bullet$,
2. constructing functors $\pi_0(\mathfrak{M}\mathcal{D}_d) \rightarrow \bullet\text{Mod}_\bullet$ given a continuous functor $\mathfrak{M}\mathcal{D}_d \rightarrow \bullet\text{Cov}_\bullet$,

where $\bullet\text{Cov}_\bullet$ and $\bullet\text{Mod}_\bullet$ respectively denote categories of spaces equipped with bicoverings and bimodules over \mathbb{A} -algebras; *cf.* Definitions 2.2 and 2.3. Here, $\pi_0(\mathcal{C})$ is obtained from \mathcal{C} by applying the functor π_0 to all morphism spaces. In this section, we deal with step (2), producing a homological representation of a category starting from a functor from a topological version of that category to covering spaces. More precisely, assuming that we are given a continuous functor $F: \mathcal{C} \rightarrow \bullet\text{Cov}_\bullet$ from any topologically-enriched category \mathcal{C} , we use twisted homology to obtain an induced functor $L_i(F): \pi_0(\mathcal{C}) \rightarrow \bullet\text{Mod}_\bullet$. More generally, this construction may itself be *twisted* by another functor $V: \mathcal{C} \rightarrow \bullet\text{Mod}_\bullet$ that is compatible with F , resulting in an induced functor $L_i(F; V): \pi_0(\mathcal{C}) \rightarrow \bullet\text{Mod}_\bullet$. This generalisation allows step (2) of the construction to be *iterated*, so that a continuous functor $F: \mathcal{C} \rightarrow \text{Cov}_\bullet$ induces an endofunctor for the functor categories

$$\mathbf{Fct}(\mathcal{C}, \text{Mod}_\bullet) \longrightarrow \mathbf{Fct}(\mathcal{C}, \text{Mod}_\bullet).$$

We begin with an overview of the construction in §2.1, and give precise definitions of all of the categories involved in §2.2. The three main steps of the construction are then described in detail

in §2.3–§2.5. This is put together into the general construction (in its non-iterative version) in §2.6, and the iterative version is described in §2.7. Throughout this section, we fix an associative, unital ring \mathbb{A} . The reader may prefer to assume that $\mathbb{A} = \mathbb{Z}$, since this will be the case in all of our examples later.

2.1. Overview

We first give a brief overview of the construction of a *homological representation* of the discrete category $\pi_0(\mathcal{C})$ starting from a functor from a topologically-enriched category \mathcal{C} to covering spaces. There are four versions of this, where the first three are increasingly general, and the fourth is an “iterative” version of the third. Full details of the constructions are given in the remainder of this section, and summarised in §2.6 and §2.7.

(1.) In the first version of the construction, we fix an \mathbb{A} -algebra S and a group G . The inputs for the construction consist of an $(\mathbb{A}[G], S)$ -bimodule V and a continuous functor F taking values in the category whose objects are spaces X equipped with a quotient of $\pi_1(X)$ onto G . Such quotients correspond to regular coverings of X with deck transformation group G , which gives this category its name, Cov_G . The output is the functor $L_i(F; V): \pi_0(\mathcal{C}) \rightarrow \text{Mod}_S$.

The construction may be summarised in the following diagram (with inputs in red and outputs in blue):

$$\begin{array}{ccccccc}
 \mathcal{C} & \xrightarrow{F} & \text{Cov}_G & \xrightarrow{\text{Lift}} & \text{Top}_{\mathbb{A}[G]} & \xrightarrow{- \otimes_{\mathbb{A}[G]} V} & \text{Top}_S & \xrightarrow{H_i} & \text{Mod}_S \\
 \downarrow & & & & & & & & \\
 \pi_0(\mathcal{C}) & & & & & & & \xrightarrow{L_i(F; V)} &
 \end{array} \tag{2.1}$$

and consists in composing F with three functors:

- The functor Lift first takes a space X equipped with a quotient $\pi_1(X) \twoheadrightarrow G$ to the corresponding regular covering of X (this amounts to a certain *lifting* construction for morphisms, whence the name). A regular covering with deck transformation group G is the same thing as a principal G -bundle, and thus in particular a bundle of G -sets. The second step of this functor is to replace each fibre with the free \mathbb{A} -module that it generates, obtaining a bundle of $\mathbb{A}[G]$ -modules. Top_R is the category of spaces equipped with bundles of R -modules. See §2.3 for more details.
- The second functor takes the fibrewise tensor product with the $(\mathbb{A}[G], S)$ -bimodule V , producing a bundle of S -modules; cf. §2.4.
- Finally, H_i is simply the functor encoding twisted homology (in degree i) over S .

The categories and functors in diagram (2.1) are all *topologically-enriched*, but the right-hand category Mod_S is in fact a *discrete* category. Hence the composition $\mathcal{C} \rightarrow \text{Mod}_S$ factors uniquely through the canonical functor $\mathcal{C} \rightarrow \pi_0(\mathcal{C})$ via a functor $\pi_0(\mathcal{C}) \rightarrow \text{Mod}_S$, which is the output of the construction.

(2.) More generally, we may begin with a bimodule V as above and a functor F into the category whose objects are spaces X equipped with a pair of quotients of $\pi_1(X)$, which may be thought of as a category of *bicoverings* of X . The construction is analogous to the above, going via bundles of bimodules to output a functor $L_i(F; V): \pi_0(\mathcal{C}) \rightarrow \bullet\text{Mod}_S$, where $\bullet\text{Mod}_S$ is a certain category – defined below – of (R, S) -bimodules, where R is allowed to vary. This construction is summarised in the following diagram, which is a copy of diagram (2.11) below:

$$\begin{array}{ccccccc}
 \mathcal{C} & \xrightarrow{F} & \bullet\text{Cov}_G & \xrightarrow{\text{Lift}} & \bullet\text{Top}_{\mathbb{A}[G]} & \xrightarrow{- \otimes_{\mathbb{A}[G]} V} & \bullet\text{Top}_S & \xrightarrow{H_i} & \bullet\text{Mod}_S \\
 \downarrow & & & & & & & & \\
 \pi_0(\mathcal{C}) & & & & & & & \xrightarrow{L_i(F; V)} &
 \end{array}$$

(3.) This is in turn a special case of our first general construction, which is summarised in diagram (2.10) below. The main point is that, in the general construction, the bimodule V is also allowed to vary, in the sense that it is replaced by a functor $V: \mathcal{C} \rightarrow \bullet\text{Mod}\bullet$ to the category of bimodules that is *compatible* with the functor $F: \mathcal{C} \rightarrow \bullet\text{Cov}\bullet$ in a specific sense; *cf.* Condition 2.19. This recovers case (2.) above, i.e. diagram (2.11), exactly when V is a constant functor. See §2.6 for more details.

(4.) Now that the input V has been generalised to a functor $\mathcal{C} \rightarrow \bullet\text{Mod}\bullet$, one may wonder whether the construction, for fixed F and i , may be iterated (bearing in mind the fact that continuous functors $\mathcal{C} \rightarrow \bullet\text{Mod}\bullet$ are in bijective correspondence with functors $\pi_0(\mathcal{C}) \rightarrow \bullet\text{Mod}\bullet$). However, this does not work as formulated above. In order to apply the construction above, the functor V is required to satisfy the compatibility condition 2.19 (depending on F). That V satisfies Condition 2.19 does not imply that $L_i(F; V)$ satisfies Condition 2.19, so we cannot apply the construction $L_i(F; -)$ again to $L_i(F; V)$. To solve this issue, we give a modified version of the construction that may be iterated indefinitely, summarised in diagram (2.15), amounting to an endofunctor for the functor categories $\mathbf{Fct}(\mathcal{C}, \text{Mod}\bullet) \rightarrow \mathbf{Fct}(\mathcal{C}, \text{Mod}\bullet)$; *cf.* Proposition 2.31. See §2.7 for more details.

2.2. Categories

In this subsection, we define the different categories involved in the construction.

Remark 2.1 In fact, all of the categories that we define are *topologically-enriched* categories – in other words their morphism sets are equipped with topologies such that composition is continuous – and all functors are *continuous* functors. We will therefore implicitly write *category* to mean topologically-enriched category and *functor* to mean continuous functor. The morphism spaces of the categories defined in this section are all equipped with topologies derived in an obvious way from the *compact-open topology* for sets of maps between topological spaces.

Definition 2.2 (*The category of biconverings.*) The category $\bullet\text{Cov}\bullet$ is the category of *spaces equipped with biconverings*. An object of $\bullet\text{Cov}\bullet$ is a path-connected, based space X admitting a universal covering (i.e., locally path-connected and semi-locally simply-connected), equipped with a pair of surjective homomorphisms

$$\phi_1: \pi_1(X) \longrightarrow Q_1 \quad \phi_2: \pi_1(X) \longrightarrow Q_2$$

such that the induced homomorphism $(\phi_1 \times \phi_2) \circ \Delta: \pi_1(X) \rightarrow Q_1 \times Q_2$ is also surjective. Via the correspondence between path-connected, regular coverings of X and normal subgroups of $\pi_1(X)$, this is the same as a pair of normal subgroups N_1, N_2 of $\pi_1(X)$ (corresponding to regular coverings $X^{N_1} \rightarrow X$ and $X^{N_2} \rightarrow X$) such that the square

$$\begin{array}{ccc} X^{N_1 \cap N_2} & \longrightarrow & X^{N_2} \\ \downarrow \lrcorner & & \downarrow \\ X^{N_1} & \longrightarrow & X \end{array}$$

is a pullback square. A morphism in $\bullet\text{Cov}\bullet$ from (X, ϕ_1, ϕ_2) to (X', ϕ'_1, ϕ'_2) is a based, continuous map $f: X \rightarrow X'$ such that the induced homomorphism $\pi_1(f)$ sends $\ker(\phi_1)$ into $\ker(\phi'_1)$ and $\ker(\phi_2)$ into $\ker(\phi'_2)$. This implies that there are unique homomorphisms $\alpha_1: Q_1 \rightarrow Q'_1$ and $\alpha_2: Q_2 \rightarrow Q'_2$ such that

$$\phi'_1 \circ \pi_1(f) = \alpha_1 \circ \phi_1 \quad \text{and} \quad \phi'_2 \circ \pi_1(f) = \alpha_2 \circ \phi_2.$$

If G is a group, the category ${}_G\text{Cov}\bullet$ is the subcategory of $\bullet\text{Cov}\bullet$ on those objects (X, ϕ_1, ϕ_2) such that $Q_1 = G$ and those morphisms f such that the induced homomorphism α_1 mentioned above is equal to id_G . Similarly, we have a subcategory $\bullet\text{Cov}_G$. If G is the trivial group, we drop it from the notation, and write $\text{Cov}\bullet$ and $\bullet\text{Cov}$ respectively for these subcategories of $\bullet\text{Cov}\bullet$. Note that $\bullet\text{Cov}$ and $\text{Cov}\bullet$ are abstractly isomorphic, but not equal as subcategories of $\bullet\text{Cov}\bullet$. As another variant, we write $\text{Cov}_G^{\text{tw}} \subset \text{Cov}\bullet$ for the *full* subcategory on those objects (X, ϕ_1, ϕ_2) such that $Q_1 = G$ (this is generally larger than Cov_G).

Definition 2.3 (*The category of bimodules.*) The category $\bullet\text{Mod}\bullet$ is the category of *bimodules over \mathbb{A} -algebras*. An object of $\bullet\text{Mod}\bullet$ is a pair of associative, unital \mathbb{A} -algebras (R, S) together with an (R, S) -bimodule V . A morphism from (R, S, V) to (R', S', V') is a pair of \mathbb{A} -algebra homomorphisms $\varphi: R \rightarrow R'$ and $\psi: S \rightarrow S'$ preserving units, together with a morphism of (R, S) -bimodules $\theta: V \rightarrow (\varphi, \psi)^*(V')$.

For any \mathbb{A} -algebra R , we have a subcategory ${}_R\text{Mod}\bullet \subset \bullet\text{Mod}\bullet$ on those objects (R', S', V') where $R' = R$ and those morphisms (φ, ψ, θ) where $\varphi = \text{id}_R$ (note that this is generally not a full subcategory). If moreover R is the trivial \mathbb{A} -algebra, we drop it from the notation, and write $\text{Mod}\bullet$ for ${}_R\text{Mod}\bullet$. Similarly, we have $\bullet\text{Mod}_S$ and ${}_R\text{Mod}_S$ for \mathbb{A} -algebras R and S . Note that ${}_R\text{Mod}_S$ is just the category of (R, S) -bimodules, as usually defined. Furthermore, for a fixed \mathbb{A} -algebra S , we write $\text{Mod}_S^{\text{tw}} \subset \text{Mod}\bullet$ for the *full* subcategory on those objects (S', V') where $S' = S$ (this is generally larger than Mod_S).

Definition 2.4 (*Bundles of G -sets and of R -modules.*) For a path-connected space X , a bundle of left G -sets over X is a fibre bundle over X with fibre a left G -set T and structure group $\text{Aut}_G(T)$, the automorphism group of T as a left G -set. See, for example, [Ste51, §2] for the definition of fibre bundle with specified fibre and structure group. In general, for a space X , a bundle of left G -sets over X is a bundle of left G -sets over each of its path-components. Bundles of left R -modules, (R, S) -bimodules and other algebraic structures are defined similarly. *However, we will shortly replace this definition with a slightly different one that is equivalent for our purposes — see Definition 2.5 and Remark 2.6 below.*

If X is a space with the property that each path-component admits a universal covering (equivalently, X is locally path-connected and semi-locally simply-connected), then a bundle of left G -sets over X is equivalent to a functor from the fundamental groupoid $\pi_{\leq 1}(X)$ to the category ${}_G\text{Set}$ of left G -sets. More precisely, for any space X , a bundle of left G -sets over X (which is in particular a covering of X , since it is a bundle with discrete fibres) determines a functor $\pi_{\leq 1}(X) \rightarrow {}_G\text{Set}$ via unique path-lifting. If X satisfies the above conditions, the induced functor from the category of bundles of left G -sets over X to the functor category $\mathbf{Fct}(\pi_{\leq 1}(X), {}_G\text{Set})$ is an equivalence. An exactly analogous statement holds for bundles of R -modules, (R, S) -bimodules and other algebraic structures — if the base space X is locally path-connected and semi-locally simply-connected, then these correspond equivalently to functors from the fundamental groupoid $\pi_{\leq 1}(X)$ to the appropriate category of algebraic structures.

Definition 2.5 (*Bundles of G -sets and of R -modules, replacing Definition 2.4.*) For any space X , a bundle of left G -sets over X is a functor

$$\pi_{\leq 1}(X) \longrightarrow {}_G\text{Set},$$

where $\pi_{\leq 1}(X)$ is the fundamental groupoid of X and ${}_G\text{Set}$ is the category of left G -sets. Similarly, a bundle of left R -modules over X is a functor from $\pi_{\leq 1}(X)$ to the category of left R -modules and a bundle of (R, S) -bimodules over X is a functor from $\pi_{\leq 1}(X)$ to the category of (R, S) -bimodules.

Remark 2.6 Definitions 2.4 and 2.5 agree whenever X is locally path-connected and semi-locally simply-connected. In all of our examples, the base space X will have these properties, so we will freely pass between these two viewpoints of bundles of algebraic structures over X .

Definition 2.7 (*The category of bundles of bimodules.*) The category $\bullet\text{Top}\bullet$ is the category of *bundles of bimodules over \mathbb{A} -algebras*. An object of $\bullet\text{Top}\bullet$ is a space X together with a pair of associative, unital \mathbb{A} -algebras (R, S) and a bundle of (R, S) -bimodules over X , meaning a functor $\xi: \pi_{\leq 1}(X) \rightarrow {}_R\text{Mod}_S$ (cf. Remark 2.6), where $\pi_{\leq 1}(X)$ is the fundamental groupoid of X . A morphism from (X, R, S, ξ) to (X', R', S', ξ') is a continuous map $f: X \rightarrow X'$, two \mathbb{A} -algebra homomorphisms $\varphi: R \rightarrow R'$ and $\psi: S \rightarrow S'$ preserving units, and an endofunctor $F: {}_R\text{Mod}_S \rightarrow {}_{R'}\text{Mod}_{S'}$ such that $(\varphi, \psi)^* \circ \xi' \circ \pi_{\leq 1}(f) = F \circ \xi$, where $(\varphi, \psi)^*: {}_{R'}\text{Mod}_{S'} \rightarrow {}_R\text{Mod}_S$ is the restriction functor induced by φ and ψ .

For any \mathbb{A} -algebra R , we have a subcategory ${}_R\text{Top}\bullet \subset \bullet\text{Top}\bullet$ on those objects (X', R', S', ξ') where $R' = R$ and those morphisms (f, φ, ψ) where $\varphi = \text{id}_R$ (note that this is generally not a full

subcategory). If moreover R is the trivial \mathbb{A} -algebra, we drop it from the notation, and write \mathbf{Top}_\bullet for ${}_R\mathbf{Top}_\bullet$. Similarly, we have $\bullet\mathbf{Top}_S$ and ${}_R\mathbf{Top}_S$ for \mathbb{A} -algebras R and S . For a fixed \mathbb{A} -algebra S , we write $\mathbf{Top}_S^{\text{tw}} \subset \mathbf{Top}_\bullet$ for the *full* subcategory on those objects (X', S', ξ') where $S' = S$ (this is generally larger than \mathbf{Top}_S).

Via the correspondence of Remark 2.6 between bundles of bimodules over X and functors out of $\pi_{\leq 1}(X)$, the endofunctor F above corresponds to a bundle map, covering f , from ξ to $(\varphi, \psi)^*(\xi')$.

Note also that $\bullet\mathbf{Mod}_\bullet$ is equivalent to the full subcategory of $\bullet\mathbf{Top}_\bullet$ on those objects whose underlying space is a point.

Notation 2.8 Writing $\mathbb{A}\text{-Alg}$ for the category of associative, unital \mathbb{A} -algebras and \mathbf{Grp} for the category of groups, there are obvious forgetful functors

$$\bullet\mathbf{Cov}_\bullet \begin{array}{c} \xrightarrow{\ell} \\ \xrightarrow[r]{} \end{array} \mathbf{Grp} \qquad \bullet\mathbf{Mod}_\bullet \subset \bullet\mathbf{Top}_\bullet \begin{array}{c} \xrightarrow{\ell} \\ \xrightarrow[r]{} \end{array} \mathbb{A}\text{-Alg}$$

that remember just the left (respectively right) underlying group or \mathbb{A} -algebra. For example, an object (X, R, S, ξ) of $\bullet\mathbf{Top}_\bullet$ is sent under ℓ to R and under r to S .

Definition 2.9 Write $\bullet\mathbf{Top}_\bullet\mathbf{Mod}_\bullet$ for the pullback of the forgetful functors $\ell: \bullet\mathbf{Mod}_\bullet \rightarrow \mathbb{A}\text{-Alg}$ and $r: \bullet\mathbf{Top}_\bullet \rightarrow \mathbb{A}\text{-Alg}$:

$$\begin{array}{ccc} \bullet\mathbf{Top}_\bullet\mathbf{Mod}_\bullet & \longrightarrow & \bullet\mathbf{Mod}_\bullet \\ \downarrow \lrcorner & & \downarrow \ell \\ \bullet\mathbf{Top}_\bullet & \xrightarrow[r]{} & \mathbb{A}\text{-Alg}. \end{array} \quad (2.2)$$

In other words, an object of $\bullet\mathbf{Top}_\bullet\mathbf{Mod}_\bullet$ consists of a space X , a triple of \mathbb{A} -algebras (R, S, T) , a bundle of (R, S) -bimodules over X and an (S, T) -bimodule V .

2.3. From bicoverings to bundles of bimodules

Proposition 2.10 *There is a natural continuous functor*

$$\text{Lift}: \bullet\mathbf{Cov}_\bullet \longrightarrow \bullet\mathbf{Top}_\bullet, \quad (2.3)$$

taking bicoverings to bundles of bimodules, such that the squares

$$\begin{array}{ccc} \bullet\mathbf{Cov}_\bullet & \xrightarrow{\text{Lift}} & \bullet\mathbf{Top}_\bullet \\ \ell \downarrow & & \downarrow \ell \\ \mathbf{Grp} & \xrightarrow{\mathbb{A}[(-)^{\text{op}}]} & \mathbb{A}\text{-Alg} \end{array} \qquad \begin{array}{ccc} \bullet\mathbf{Cov}_\bullet & \xrightarrow{\text{Lift}} & \bullet\mathbf{Top}_\bullet \\ r \downarrow & & \downarrow r \\ \mathbf{Grp} & \xrightarrow{\mathbb{A}[-]} & \mathbb{A}\text{-Alg} \end{array} \quad (2.4)$$

commute, where $\mathbb{A}[-]: \mathbf{Grp} \rightarrow \mathbb{A}\text{-Alg}$ takes a group to its group \mathbb{A} -algebra and $(-)^{\text{op}}: \mathbf{Grp} \rightarrow \mathbf{Grp}$ takes a group to its opposite.

Proof. On objects, this is defined as follows. Let $(X, x_0, \phi_1: \pi_1(X, x_0) \rightarrow Q_1, \phi_2: \pi_1(X, x_0) \rightarrow Q_2)$ be an object of $\bullet\mathbf{Cov}_\bullet$. The normal subgroup

$$K := \ker(\phi_1) \cap \ker(\phi_2) = \ker((\phi_1 \times \phi_2) \circ \Delta) \triangleleft \pi_1(X, x_0) \quad (2.5)$$

corresponds to a regular covering of X with deck transformation group $Q := Q_1 \times Q_2$. In order to specify a particular regular covering of X , rather than just an *isomorphism class* of such, we will be slightly more careful. Start with the universal covering \tilde{X} of X ; more specifically, the standard model for \tilde{X} whose underlying set consists of endpoint-preserving homotopy classes of paths in X starting at x_0 . This is equipped with an action of $\pi_1(X, x_0)$; take the quotient of \tilde{X} by the action

of the subgroup K . We denote this regular covering by $\xi_K: X^K = \tilde{X}/K \rightarrow X$. Whether deck transformations act on the left or the right is an arbitrary convention. We will consider them to act on the *right*, since this agrees with the typical convention that the structure group of a principal bundle (of which regular coverings are examples) acts on the total space on the *right*. This is a bundle of right Q -sets over X (whose fibres all happen to be isomorphic to Q itself). Since Q is the product $Q_1 \times Q_2$, we may equally view ξ_K as a bundle of (Q_1^{op}, Q_2) -bisets over X , where a (G, H) -biset is a set equipped with a left G -action and a compatible right H -action.

Now replace each fibre of ξ_K with the free \mathbb{A} -module generated by that fibre; this forms a bundle of $(\mathbb{A}[Q_1^{\text{op}}], \mathbb{A}[Q_2])$ -bimodules over X . The operation of “*taking free \mathbb{A} -modules fibrewise*” is simplest to describe by viewing bundles of (G, H) -bisets over X as functors $\pi_{\leq 1}(X) \rightarrow {}_G\text{Set}_H$ to the category of (G, H) -bisets, and similarly bundles of (R, S) -bimodules over X as functors $\pi_{\leq 1}(X) \rightarrow {}_R\text{Mod}_S$ to the category of (R, S) -bimodules. In this viewpoint, the operation is simply post-composition with the free functor $\mathbb{A}[-]: Q_1^{\text{op}}\text{Set}_{Q_2} \rightarrow \mathbb{A}[Q_1^{\text{op}}]\text{Mod}_{\mathbb{A}[Q_2]}$. Denote the resulting bundle of bimodules by $\mathbb{A}_{\text{fib}}[\xi_K]: \mathbb{A}_{\text{fib}}[X^K] \rightarrow X$. This defines Lift on objects:

$$\text{Lift}(X, x_0, \phi_1, \phi_2) = (X, \mathbb{A}[Q_1^{\text{op}}], \mathbb{A}[Q_2], \mathbb{A}_{\text{fib}}[\xi_K]).$$

In order to define Lift on morphisms, we first note that, although we did not need it to define the functor on objects, the regular covering $\xi_K: X^K \rightarrow X$ associated to (X, x_0, ϕ_1, ϕ_2) comes equipped with a particular choice of basepoint of X^K , covering the basepoint x_0 of X . This is because the standard construction of the universal cover \tilde{X} has a canonical basepoint (namely the constant path at x_0), and therefore so does its quotient X^K . Let us denote this basepoint by $\tilde{x}_0 \in X^K$.

Now suppose we have a morphism $(X, \phi_1, \phi_2) \rightarrow (Y, \phi'_1, \phi'_2)$ in $\bullet\text{Cov}_\bullet$, that is, a continuous map $f: X \rightarrow Y$ such that $f(x_0) = y_0$, $f_*(\ker(\phi_1)) \subseteq \ker(\phi'_1)$ and $f_*(\ker(\phi_2)) \subseteq \ker(\phi'_2)$. We recall from Definition 2.2 that this determines certain homomorphisms $\alpha_1: Q_1 \rightarrow Q'_1$ and $\alpha_2: Q_2 \rightarrow Q'_2$, which determine \mathbb{A} -algebra homomorphisms $\mathbb{A}[\alpha_1^{\text{op}}]: \mathbb{A}[Q_1^{\text{op}}] \rightarrow \mathbb{A}[(Q'_1)^{\text{op}}]$ and $\mathbb{A}[\alpha_2]: \mathbb{A}[Q_2] \rightarrow \mathbb{A}[Q'_2]$. Let

$$K = \ker(\phi_1) \cap \ker(\phi_2) \quad L = \ker(\phi'_1) \cap \ker(\phi'_2)$$

and write $\xi_K: X^K \rightarrow X$ and $\xi_L: Y^L \rightarrow Y$ for the corresponding regular covering spaces. By covering space theory, for each point $\tilde{y} \in \xi_L^{-1}(y_0)$, there is a unique continuous map $X^K \rightarrow Y^L$ that lifts the composition $f \circ \xi_K: X^K \rightarrow Y$ and that takes \tilde{x}_0 to \tilde{y} . We therefore obtain a uniquely-determined lift

$$\tilde{f}: X^K \longrightarrow Y^L$$

by requiring $\tilde{f}(\tilde{x}_0) = \tilde{y}_0$. Extending this map \mathbb{A} -linearly in each fibre results in a map

$$\mathbb{A}_{\text{fib}}[\tilde{f}]: \mathbb{A}_{\text{fib}}[X^K] \longrightarrow \mathbb{A}_{\text{fib}}[Y^L]$$

of bundles of \mathbb{A} -modules. Finally, one may check that $\mathbb{A}_{\text{fib}}[\tilde{f}]$ is a map of bundles of $(\mathbb{A}[Q_1^{\text{op}}], \mathbb{A}[Q_2])$ -bimodules, covering f , from $\mathbb{A}_{\text{fib}}[\xi_K]$ to $\mathbb{A}_{\text{fib}}[\xi_L]$, where the latter is given the structure of a bundle of $(\mathbb{A}[Q_1^{\text{op}}], \mathbb{A}[Q_2])$ -bimodules via $\mathbb{A}[\alpha_1^{\text{op}}]$ and $\mathbb{A}[\alpha_2]$. This uses the interpretation of morphisms of $\bullet\text{Top}_\bullet$ from Remark 2.6. Hence we may define

$$\text{Lift}(f) = (f, \mathbb{A}[\alpha_1^{\text{op}}], \mathbb{A}[\alpha_2], \mathbb{A}_{\text{fib}}[\tilde{f}]). \quad \square$$

Remark 2.11 (*Modified lifting.*) As a brief aside, we mention a non-functorial modification of the lifting functor, defined as follows. We recall that an input object for the lifting functor consists of a space X and a jointly surjective pair of maps $\phi_1: \pi_1(X) \rightarrow Q_1$ and $\phi_2: \pi_1(X) \rightarrow Q_2$, and the output is the tuple $(\mathbb{A}[Q_1], \mathbb{A}[Q_2], X, \xi)$, where ξ is a certain bundle of $(\mathbb{A}[Q_1], \mathbb{A}[Q_2])$ -bimodules over X . Via the ring homomorphism $\mathbb{A}[\pi_1(X)] \rightarrow \mathbb{A}[Q_2]$, we may consider ξ to be a bundle of $(\mathbb{A}[Q_1], \mathbb{A}[\pi_1(X)])$ -bimodules over X . Define:

$$\text{Lift}'(X, \phi_1, \phi_2) = (\mathbb{A}[Q_1], \mathbb{A}[\pi_1(X)], X, \xi).$$

This defines a function on the objects $\text{ob}(\bullet\text{Cov}_\bullet) \rightarrow \text{ob}(\bullet\text{Top}_\bullet)$. This definition cannot, however, be extended to morphisms in general, so this function of object sets cannot be extended to a modified lift *functor*. Nevertheless, this alternative “*modified lifting function*” at the level of objects can be useful for constructing Markov functions (which need not be representations) on the classical braid groups. See for example [Con18; BC18].

2.4. Fibrewise tensor product

In this short subsection, we define the “*fibrewise tensor product*” functor

$$\otimes: \bullet\text{Top}_\bullet\text{Mod}_\bullet \longrightarrow \bullet\text{Top}_\bullet. \quad (2.6)$$

Notation 2.12 Recall from Definition 2.7 that, for two \mathbb{A} -algebra homomorphisms $\alpha: R \rightarrow R'$ and $\beta: S \rightarrow S'$, we denote the corresponding restriction functor ${}_{R'}\text{Mod}_{S'} \rightarrow {}_R\text{Mod}_S$ by $(\alpha, \beta)^*$.

Definition 2.13 We define the functor (2.6) as follows.

- (*On objects.*) We recall that an object of $\bullet\text{Top}_\bullet\text{Mod}_\bullet$ consists of a space X , three \mathbb{A} -algebras R, S, T , an (S, T) -bimodule V and a bundle $\xi: \pi_{\leq 1}(X) \rightarrow {}_R\text{Mod}_S$ of (R, S) -bimodules over X . Define its image under (2.6) to be the following object of $\bullet\text{Top}_\bullet$:

$$(X, R, T, \pi_{\leq 1}(X) \xrightarrow{\xi} {}_R\text{Mod}_S \xrightarrow{-\otimes_S V} {}_R\text{Mod}_T).$$

- (*On morphisms.*) A morphism of $\bullet\text{Top}_\bullet\text{Mod}_\bullet$ from (X, R, S, T, V, ξ) to $(X', R', S', T', V', \xi')$ consists of a continuous map $f: X \rightarrow X'$, \mathbb{A} -algebra homomorphisms $\alpha: R \rightarrow R'$, $\beta: S \rightarrow S'$ and $\gamma: T \rightarrow T'$, a homomorphism $\theta: V \rightarrow (\beta, \gamma)^*V'$ of (S, T) -bimodules and a natural transformation $\tau: \xi \Rightarrow (\alpha, \beta)^* \circ \xi'$. Define its image under (2.6) to be the morphism

$$(f, \alpha, \gamma, \hat{\tau})$$

of $\bullet\text{Top}_\bullet$, where $\hat{\tau}: (-\otimes_S V) \circ \xi \Rightarrow (\alpha, \gamma)^* \circ (-\otimes_{S'} V') \circ \xi'$ is the natural transformation defined as follows. First, note that, for any (R, S') -bimodule A and (S', T) -bimodule B , there is a canonical homomorphism of (R, T) -bimodules

$$\beta^* A \otimes_S \beta^* B \longrightarrow A \otimes_{S'} B,$$

which is an isomorphism if $\beta: S \rightarrow S'$ is surjective. Using this fact, we define $\hat{\tau}$ on the object x of $\pi_{\leq 1}(X)$ by

$$\begin{aligned} \xi(x) \otimes_S V &\longrightarrow (\alpha, \beta)^*(\xi'(x)) \otimes_S V \\ &\longrightarrow (\alpha, \beta)^*(\xi'(x)) \otimes_S (\beta, \gamma)^*V' \\ &\longrightarrow (\alpha, 1)^*(\xi'(x)) \otimes_{S'} (1, \gamma)^*V' \\ &= (\alpha, \gamma)^*(\xi'(x) \otimes_{S'} V'), \end{aligned}$$

where the first arrow is induced by τ_x , the second is induced by θ and the third is the canonical homomorphism from above.

Remark 2.14 This definition is somewhat formal, but it has a very natural geometric interpretation. If we view a bundle as an actual bundle over X (cf. Remark 2.6) and choose an open cover \mathcal{U} of X together with trivialisations of the bundle over each $U \in \mathcal{U}$, then we may take the tensor product over each U (since the bundle is now trivial over each U and it is obvious how to define this) and then glue these trivial bundles back together again using the same transition functions as for the original bundle.

Notation 2.15 There are forgetful functors $\bullet\text{Top}_\bullet\text{Mod}_\bullet \rightarrow \bullet\text{Top}_\bullet$ and $\bullet\text{Top}_\bullet\text{Mod}_\bullet \rightarrow \bullet\text{Mod}_\bullet$ coming from the pullback square (2.2). For a continuous functor $F: \mathcal{C} \rightarrow \bullet\text{Top}_\bullet\text{Mod}_\bullet$, denote its compositions with these two forgetful functors by

$$F_1: \mathcal{C} \longrightarrow \bullet\text{Top}_\bullet \quad \text{and} \quad F_2: \mathcal{C} \longrightarrow \bullet\text{Mod}_\bullet$$

respectively. With this notation, we define $F_1 \otimes F_2$ by $F_1 \otimes F_2 := \otimes \circ F$.

2.5. Twisted homology

Over a fixed \mathbb{A} -algebra R , one may view local coefficient systems on a space X as bundles of right R -modules over X . In this viewpoint, homology with local coefficients (in any fixed degree $i \geq 0$) is a functor of the form

$$H_i: \text{Top}_R \longrightarrow \text{Mod}_R. \quad (2.7)$$

See, for example [DK01, §5.4] or [Pal18b, §5.1]. The following fact may be proven by directly generalising the usual construction of singular twisted homology, keeping careful track of the variable bimodule structure.

Proposition 2.16 *In any degree $i \geq 0$, homology with local coefficients extends to a functor*

$$H_i: \bullet\text{Top}_\bullet \longrightarrow \bullet\text{Mod}_\bullet \quad (2.8)$$

such that the square

$$\begin{array}{ccc} \bullet\text{Top}_\bullet & \xrightarrow{H_i} & \bullet\text{Mod}_\bullet \\ \uparrow & & \uparrow \\ \text{Top}_R & \xrightarrow{H_i} & \text{Mod}_R \end{array} \quad (2.9)$$

commutes for any \mathbb{A} -algebra R . There are exactly analogous statements for relative homology (on a category of bimodule bundles over pairs of spaces) and Borel-Moore homology (where we must restrict to locally-compact spaces and proper maps). Moreover, there are similar statements for cohomology (including compactly-supported cohomology, where we must again restrict to locally-compact spaces and proper maps), although the domain category is not the opposite of $\bullet\text{Top}_\bullet$, but rather another category $\bullet\text{Top}_\bullet^{\leftarrow}$ defined below.

Definition 2.17 The category $\bullet\text{Top}_\bullet^{\leftarrow}$ has the same objects as $\bullet\text{Top}_\bullet$, but its morphisms are slightly different: they are *contravariant* on the underlying spaces and *covariant* on bimodule bundles (local systems). More precisely, its morphisms are exactly as in Definition 2.7, except that the map f goes in the opposite direction $X' \rightarrow X$ and the compatibility condition is correspondingly changed to $(\varphi, \psi)^* \circ \xi' = F \circ \xi \circ \pi_{\leq 1}(f)$. See also [DK01, Theorem 5.12].

Remark 2.18 (*Interpreting twisted homology as homology of covering spaces.*) The homological representations that we construct will be a composition of functors, ending with (2.8) (or one of its variants, such as twisted Borel-Moore homology). Thus, restricted to each object of the domain, they will give a representation of its automorphism group on the twisted homology of some space X equipped with a local system \mathcal{L} . Typically, \mathcal{L} will correspond to a regular covering $\hat{X} \rightarrow X$ (in fact, it will *always* arise from a regular covering, followed possibly by a fibrewise tensor product). If so, the twisted homology $H_*(X; \mathcal{L})$ is canonically isomorphic to $H_*(\hat{X})$, by Shapiro's lemma for covering spaces, so we may think of the homological representation as an action on the untwisted homology of the covering space \hat{X} . — However, this last remark only applies to (twisted) *ordinary* homology, and not to (twisted) *Borel-Moore* homology, since Shapiro's lemma for covering spaces is generally false for Borel-Moore homology. For example if $\hat{X} \rightarrow X$ is the universal covering $\mathbb{R} \rightarrow S^1$ then $H_1^{BM}(S^1; \mathcal{L}) = 0 \not\cong \mathbb{Z} \cong H_1^{BM}(\mathbb{R})$. See [AP20, §6.1 and §6.2] for further details.

2.6. The general construction

The first version of our general construction is obtained by concatenating the lifting functor of §2.3, the fibrewise tensor product of §2.4 and twisted homology §2.5.

Suppose that we are given a topologically-enriched category \mathcal{C} , which is assumed to be *nice*, meaning that, for each pair of objects (x, y) , the connected components of the morphism space $\mathcal{C}(x, y)$ are path-connected. Note that this condition holds whenever each $\mathcal{C}(x, y)$ is locally path-connected, and it ensures that, for any discrete category \mathcal{D} , any continuous functor $\mathcal{C} \rightarrow \mathcal{D}$ factors (uniquely) through $\mathcal{C} \rightarrow \pi_0(\mathcal{C})$. The input for the construction consists of

- a continuous functor $F: \mathcal{C} \rightarrow \bullet\text{Cov}_\bullet$,
- a continuous functor $V: \mathcal{C} \rightarrow \bullet\text{Mod}_\bullet$,
- a positive integer $i \geq 0$,

satisfying Condition 2.19 below, where we recall (*cf.* Notation 2.8) that ℓ and r denote functors

Remark 2.24 (*Other variants of homology.*) All of the above constructions go through identically if we replace (twisted) ordinary homology with (twisted) Borel-Moore homology, cohomology, compactly-supported cohomology, reduced homology, etc., as long as we modify each category in the construction accordingly. For example, if we wish to use twisted Borel-Moore homology, we must restrict to locally compact spaces and proper maps, so in particular the input functor F must take values in this subcategory of $\bullet\text{Cov}_\bullet$. Similarly, if we wish to use twisted cohomology, we must replace $\bullet\text{Top}_\bullet$ with the “ambivariant” version $\bullet\text{Top}_\bullet^\leftarrow$, as described in Definition 2.17, and similarly for $\bullet\text{Cov}_\bullet$, and F must take values in $\bullet\text{Cov}_\bullet^\leftarrow$ instead of $\bullet\text{Cov}_\bullet$.

Remark 2.25 Another possible simplification is to *omit* the step where we take the fibrewise tensor product with V . Let us write the output of this simplified construction (which depends only on F) by $L_i(F)$. This is actually a special case of the setting where we *do* take the fibrewise tensor product with a particular V_F determined by F ; in other words, we have $L_i(F) = L_i(F; V_F)$. The “trivial” coefficient system V_F may be constructed as follows. We denote by $\text{triv}: \mathbb{A}\text{-Alg} \rightarrow \bullet\text{Mod}_\bullet$ the functor that sends R to the bimodule (R, R, R) . Then V_F is the composition $\text{triv} \circ \mathbb{A}[-] \circ r \circ F$. In this case, the diagram (2.10) simplifies to:

$$\begin{array}{ccccc}
 \mathcal{C} & \xrightarrow{F} & \bullet\text{Cov}_\bullet & \xrightarrow{\text{Lift}} & \bullet\text{Top}_\bullet & \xrightarrow{H_i} & \bullet\text{Mod}_\bullet \\
 \downarrow & & & & & \nearrow & \\
 \pi_0(\mathcal{C}) & & & & & & \\
 & & & & & L_i(F) = L_i(F; V_F) &
 \end{array} \tag{2.12}$$

Remark 2.26 If F takes values in a subcategory of $\bullet\text{Cov}_\bullet$ with some restriction on the left module structure, then $L_i(F; V)$ will take values in the subcategory of $\bullet\text{Mod}_\bullet$ with the corresponding restriction on the left module structure. Similarly, if V takes values in a subcategory of $\bullet\text{Mod}_\bullet$ with some restriction on the right module structure, then $L_i(F; V)$ will take values in the subcategory of $\bullet\text{Mod}_\bullet$ with the same restriction on the right module structure. For example, if V takes values in $\bullet\text{Mod}$ and F takes values in ${}^{\text{tw}}_{\mathbb{A}[G]}\text{Cov}_\bullet$, then $L_i(F; V)$ will take values in ${}^{\text{tw}}_{\mathbb{A}[G]}\text{Mod} (\cong \text{Mod}_{\mathbb{A}[G]}^{\text{tw}})$.

Connected components, path-components and semifunctors. There is a small subtlety related to connected components and path-components in diagram (2.1), and the other diagrams illustrating the constructions. We first describe this at the level of spaces, and then at the level of *topological categories*, i.e. categories enriched over the symmetric monoidal category of topological spaces with the Cartesian product.

If $f: X \rightarrow Y$ is a continuous map with target a discrete space Y , then f must be constant on each connected component of X , hence in particular on each path-component of X , so there is a well-defined induced function $\pi_0(X) \rightarrow Y$, in other words, f factors uniquely as $X \rightarrow \pi_0(X) \rightarrow Y$ in the category of sets. In order for this statement to be true also in the category of spaces, we must either give $\pi_0(X)$ the discrete topology and assume that the path-components of X are open (which is equivalent to saying that its path-components and its connected components coincide), or more generally, without this assumption, give $\pi_0(X)$ the quotient topology induced from X and the surjective function $X \twoheadrightarrow \pi_0(X)$ (this will be the discrete topology if and only if X satisfies the assumption mentioned before).

An exactly analogous discussion holds for a continuous functor $\mathcal{C} \rightarrow \mathcal{D}$ from a topologically-enriched category \mathcal{C} to a discrete category \mathcal{D} , where the corresponding assumption on \mathcal{C} is that each morphism space $\mathcal{C}(c, c')$ has open path-components. In diagram (2.1), we would like $\pi_0(\mathcal{C})$ to be a discrete category – since ultimately we are trying to construct representations of certain discrete categories – so we must make this assumption about \mathcal{C} in order for diagram (2.1) to be strictly correct. In all of our examples, this condition will indeed be satisfied (in fact $\mathcal{C}(c, c')$ will always be locally contractible, which is much stronger).

A second minor subtlety is that, in our examples, \mathcal{C} will typically only be a semi-category (a category without identities), although $\pi_0(\mathcal{C})$ will be a (usual) category. The diagrams above should therefore be interpreted as diagrams of semifunctors. However, the induced arrow from $\pi_0(\mathcal{C})$ is always in the end a *functor*. See §3.4 for why we must allow \mathcal{C} to be a semi-category, and Lemma 5.10 for the statement that the output of the construction is nevertheless a functor.

2.7. An iterative version

We may view the construction above, for a fixed $i \geq 0$ and functor $F: \mathcal{C} \rightarrow \bullet\text{Cov}\bullet$, as sending a functor of the form $\mathcal{C} \rightarrow \bullet\text{Mod}\bullet$ to another functor of the same form. However, this does not imply that we may iterate the procedure, since the input functor $V: \mathcal{C} \rightarrow \bullet\text{Mod}\bullet$ is required to satisfy Condition 2.19 with respect to F . In this subsection, we describe a variant of the construction which may be iterated.

Definition 2.27 Fix a topologically-enriched category \mathcal{C} . Denoting by Cat/\mathcal{C} the slice category of Cat over \mathcal{C} , recall that the *Grothendieck construction* is a functor

$$f^{\mathcal{C}} = \int: \mathbf{Fct}(\mathcal{C}, \text{Cat}) \longrightarrow \text{Cat}/\mathcal{C},$$

that, on objects, takes a functor $F: \mathcal{C} \rightarrow \text{Cat}$ to the functor $\int F \rightarrow \mathcal{C}$, where the objects

$$\text{ob}(\int F) = \{(c, a) \mid c \in \text{ob}(\mathcal{C}), a \in \text{ob}(F(c))\}$$

and a morphism in $\int F$ from (c, a) to (c', a') is a morphism $f: c \rightarrow c'$ in \mathcal{C} together with a morphism $g: a \rightarrow F(f)(a')$ in $F(c)$. The functor $\int F \rightarrow \mathcal{C}$ simply takes (c, a) to c on objects and (f, g) to f on morphisms.

Definition 2.28 For a continuous functor $a: \mathcal{C} \rightarrow \text{Grp}$, we may compose it with the inclusion $i: \text{Grp} \rightarrow \text{Cat}$ into the category of small categories and then apply the (topologically enriched) Grothendieck construction to this functor. Let us denote the resulting topologically enriched category by $a \rtimes \mathcal{C}$. In other words, we define

$$a \rtimes \mathcal{C} = \int^{\mathcal{C}} (i \circ a).$$

Denoting by 0 the functor $\mathcal{C} \rightarrow \text{Grp}$ sending all objects to the trivial group, there exists a unique natural transformation $0 \rightarrow a$. Applying the Grothendieck construction, this induces a section $\mathcal{C} = 0 \rtimes \mathcal{C} \hookrightarrow a \rtimes \mathcal{C}$ of the projection functor $a \rtimes \mathcal{C} \rightarrow \mathcal{C}$ which we denote by s_a . Note that, if \mathcal{C} is a group G (considered as a one-object discrete category), then the functor a is simply a choice of another group H and a G -action on H , and we have $a \rtimes \mathcal{C} = H \rtimes G$. Hence, we may think of the construction $a \rtimes \mathcal{C}$ as a *semi-direct product with many objects*.

Lemma 2.29 Suppose we are given continuous functors

$$a: \mathcal{C} \longrightarrow \text{Grp} \quad \text{and} \quad \chi: a \rtimes \mathcal{C} \longrightarrow \mathcal{C},$$

and write $\bar{\chi}$ for the endofunctor of \mathcal{C} given by composing χ with the natural inclusion $s_a: \mathcal{C} \hookrightarrow a \rtimes \mathcal{C}$. Then any family of right modules parametrised by \mathcal{C} has – after shifting by $\bar{\chi}$ – a canonical bimodule structure determined by a and χ . More precisely, for any functor $V: \mathcal{C} \rightarrow \text{Mod}\bullet$, there is a canonical functor $V_a^{\chi}: \mathcal{C} \rightarrow \bullet\text{Mod}\bullet$ making the following diagram commute:

$$\begin{array}{ccccc}
 & & \text{Grp} & \xrightarrow{\mathbb{A}[-]} & \mathbb{A}\text{-Alg} \\
 & \nearrow a & & & \uparrow \ell \\
 & & & & \bullet\text{Mod}\bullet \\
 & & & \dashrightarrow V_a^{\chi} & \\
 \mathcal{C} & \xrightarrow{\bar{\chi}} & \mathcal{C} & \xrightarrow{V} & \text{Mod}\bullet \\
 & & & & \downarrow \text{forget}
 \end{array} \tag{2.13}$$

Proof. The category $a \rtimes \mathcal{C}$ has the same objects as \mathcal{C} , with morphisms given by

$$\text{Hom}_{a \rtimes \mathcal{C}}(c, c') = \{(\varphi, g) \mid \varphi \in \text{Hom}_{\mathcal{C}}(c, c'), g \in a(c')\}$$

with composition given by $(\varphi', g') \circ (\varphi, g) = (\varphi' \circ \varphi, g'.a(\varphi')(g))$. In particular, the automorphism group $\text{Aut}_{a \rtimes \mathcal{C}}(c)$ splits canonically as the semi-direct product $a(c) \rtimes \text{Aut}_{\mathcal{C}}(c)$. Together with the functors χ and V , noting that $\bar{\chi}(c) = \chi(c)$ on objects, we obtain a homomorphism

$$a(c) \hookrightarrow a(c) \rtimes \text{Aut}_{\mathcal{C}}(c) \cong \text{Aut}_{a \rtimes \mathcal{C}}(c) \xrightarrow{\chi} \text{Aut}_{\mathcal{C}}(\bar{\chi}(c)) \xrightarrow{V} \text{Aut}_{\text{Mod}\bullet}(V(\bar{\chi}(c))),$$

which is natural in c . If we write $V(\bar{\chi}(c)) = (W_c, S_c)$, where S_c is an \mathbb{A} -algebra and W_c is a right S_c -module, this means that we have – naturally in c – a left $a(c)$ -action on W_c that is compatible with the right S_c -action. Extending \mathbb{A} -linearly and writing $R_c = \mathbb{A}[a(c)]$, this means that we have – naturally in c – extended (W_c, S_c) to an object (R_c, W_c, S_c) of $\bullet\text{Mod}\bullet$. This is the functor V_a^χ . \square

Remark 2.30 Note that, if χ is the canonical projection functor, then $\bar{\chi}$ is the identity and, for each object c of \mathcal{C} , the bimodule $V_a^\chi(c)$ is obtained from the right-module $V(c)$ simply by giving it the trivial left-action of $\mathbb{A}[a(c)]$. In our constructions, however, $\bar{\chi}$ will typically not be the identity, but rather a kind of “shift” or “stabilisation” endofunctor of \mathcal{C} .

The iterative version of our general construction is as follows. We first fix the data of

- a continuous functor $F: \mathcal{C} \rightarrow \text{Cov}\bullet$, and
- a continuous functor $\chi: a \times \mathcal{C} \rightarrow \mathcal{C}$,

where

$$a := r \circ F. \quad (2.14)$$

Now suppose we begin (as input) with a functor

$$V: \mathcal{C} \longrightarrow \text{Mod}\bullet.$$

The relation (2.14) implies that $r \circ \text{Lift} \circ F = \mathbb{A}[a]$, and hence that the functor $\text{Lift} \circ F$ and the functor V_a^χ from Lemma 2.29 determine a functor of the form $\mathcal{C} \rightarrow \text{Top}\bullet\text{Mod}\bullet$. This is then composed with the fibrewise tensor product and twisted homology to obtain a functor of the form

$$\mathcal{C} \longrightarrow \text{Mod}\bullet,$$

which is the output of the iterative construction. This may be summarised diagrammatically as follows (the input is red, the rest of diagram (2.13) of Lemma 2.29 is green and the output is blue):

$$\begin{array}{c}
 \begin{array}{ccccc}
 \mathcal{C} & \xrightarrow{F} & \text{Cov}\bullet & \xrightarrow{\text{Lift}} & \text{Top}\bullet \\
 \bar{\chi} \downarrow & \searrow^a & r \downarrow & \searrow^{\mathbb{A}[-]} & \downarrow r \\
 \mathcal{C} & & \text{Grp} & \xrightarrow{\mathbb{A}[-]} & \mathbb{A}\text{-Alg} \\
 V \downarrow & \searrow^{V_a^\chi} & \ell \nearrow & & \\
 \text{Mod}\bullet & \xleftarrow{\text{forget}} & \bullet\text{Mod}\bullet & & \\
 & & & & \uparrow \ulcorner \\
 & & & & \text{Top}\bullet\text{Mod}\bullet \xrightarrow{\otimes} \text{Top}\bullet \xrightarrow{H_i} \text{Mod}\bullet
 \end{array} \\
 \text{A blue arrow } \Lambda_i(F; \chi)(V) \text{ points from } \text{Mod}\bullet \text{ back to } \mathcal{C}.
 \end{array} \quad (2.15)$$

In the notation of Definition 2.21, we may write:

$$V \longmapsto L_i(F; V_{r \circ F}^\chi). \quad (2.16)$$

From an easy check of the definitions, we see that this construction is *natural* in V in the following sense.

Proposition 2.31 (*The iterative construction.*) *Given continuous functors F and χ as above, the assignment (2.16) is functorial in V , in other words, it extends to an endofunctor*

$$\mathbf{Fct}(\mathcal{C}, \text{Mod}\bullet) \longrightarrow \mathbf{Fct}(\mathcal{C}, \text{Mod}\bullet). \quad (2.17)$$

Notation 2.32 Let us denote the endofunctor (2.17) of Proposition 2.31 by $\Lambda_i(F; \chi)$, so

$$\Lambda_i(F; \chi)(V) = L_i(F; V_{r \circ F}^\chi).$$

We will sometimes wish to perform a similar iterative construction when we do not have a continuous functor $F: \mathcal{C} \rightarrow \text{Cov}_\bullet$, but only a functor taking values in the discrete category $\pi_0(\text{Cov}_\bullet)$. In this case we may as well suppose that \mathcal{C} is also a discrete category (since continuous functors $\mathcal{C} \rightarrow \pi_0(\text{Cov}_\bullet)$ correspond bijectively to functors $\pi_0(\mathcal{C}) \rightarrow \pi_0(\text{Cov}_\bullet)$). The construction works equally well with such an F : we may simply apply the functor $\pi_0(\)$ – replacing the morphism spaces in a topological category with their sets of path-components – to the whole diagram (2.15), where we note that the pullback square of topological categories (2.2) remains a pullback square after applying $\pi_0(\)$.

More precisely, let us fix a category \mathcal{C} and the data of

- a functor $F: \mathcal{C} \rightarrow \pi_0(\text{Cov}_\bullet)$, and
- a functor $\chi: a \rtimes \mathcal{C} \rightarrow \mathcal{C}$,

where $a := r \circ F$. Then the diagram obtained from (2.15) by applying the functor $\pi_0(\)$ determines an assignment

$$V \longmapsto \Lambda_i(F; \chi)(V). \quad (2.18)$$

As before, this is natural in V in the following sense.

Proposition 2.33 *Given a category \mathcal{C} and functors F and χ as above, the assignment (2.18) extends to an endofunctor*

$$\Lambda_i(F; \chi): \mathbf{Fct}(\mathcal{C}, \text{Mod}_\bullet) \longrightarrow \mathbf{Fct}(\mathcal{C}, \text{Mod}_\bullet). \quad (2.19)$$

Remark 2.34 (*Other variants of homology.*) As in the previous subsection (on the non-iterative version of the construction), we may replace ordinary (twisted) homology with other flavours, such as Borel-Moore homology, cohomology, etc., as long as we make the appropriate modifications of the intermediate categories (in particular, F will have to take values in the appropriate modification of Cov_\bullet). See Remark 2.24 for more details.

3. Topological categories of decorated manifolds

In this section, we define the categories that will serve as the domain of the “homological representations” that we will construct in §5. They are obtained from certain monoidal groupoids by the *Quillen bracket construction*, an operation that enlarges a given monoidal groupoid to a monoidal category having the original monoidal groupoid as its underlying groupoid. More precisely, we will start with certain *topologically-enriched* monoidal groupoids, so we first describe, in §3.1, a topological enrichment of the Quillen bracket construction and show that it behaves well with respect to the functor π_0 that replaces all morphism spaces with their sets of path-components, subject to a *Serre fibration condition*. In §3.2, we then define the topologically-enriched monoidal groupoids that we wish to consider, and prove that they satisfy this Serre fibration condition.

Informally, the idea is that the domain category used in §5, for a given dimension $d \geq 2$, will be a topologically-enriched category $\mathcal{U}\mathcal{D}_d$ having the property that the automorphism groups of $\pi_0(\mathcal{U}\mathcal{D}_d)$ contain all motion groups and mapping class groups in dimension d . To construct this, we define in §3.2 a topological groupoid \mathcal{D}_d whose automorphism groups are the diffeomorphism groups of all d -dimensional “decorated manifolds”. The topologically-enriched Quillen bracket construction of §3.1 then gives us a topologically-enriched category $\mathcal{U}\mathcal{D}_d$ such that $\pi_0(\mathcal{U}\mathcal{D}_d) \cong \mathcal{U}(\pi_0(\mathcal{D}_d))$. The underlying groupoid of $\pi_0(\mathcal{U}\mathcal{D}_d)$ is therefore $\pi_0(\mathcal{D}_d)$, consisting of all mapping class groups of d -dimensional “decorated manifolds”, which contain all d -dimensional motion groups as normal subgroups.

3.1. A topological enrichment of the Quillen bracket construction

Throughout this section, we fix a *topological* monoidal groupoid $(\mathcal{G}, \natural, 0)$ and a *topological* category (\mathcal{M}, \natural) with a continuous left-action of \mathcal{G} . We use the abbreviation ob to denote the set of objects of a category. We refer the reader to [Mac98] for a complete introduction to the notions of (strict) monoidal categories and modules over them. We recall that, by *topological category*, we mean a

category enriched over the symmetric monoidal category of topological spaces with the Cartesian product.

Definition 3.1 The category $\langle \mathcal{G}, \mathcal{M} \rangle$ is defined to have the same objects as \mathcal{M} , and for objects X, Y of \mathcal{M} , we define $\text{Hom}_{\langle \mathcal{G}, \mathcal{M} \rangle}(X, Y)$ to be the quotient space

$$\left[\bigsqcup_{A \in \text{ob}(\mathcal{G})} \text{Hom}_{\mathcal{M}}(A \natural X, Y) \right] / \sim,$$

where \sim is the equivalence relation given by $(A, \varphi) \sim (A', \varphi')$ if and only if $\varphi = \varphi' \circ (\sigma \natural \text{id}_X)$ for some $\sigma \in \text{Hom}_{\mathcal{G}}(A, A')$. For two morphisms $[A, \varphi] : X \rightarrow Y$ and $[B, \psi] : Y \rightarrow Z$ in $\langle \mathcal{G}, \mathcal{M} \rangle$, the composition is defined by $[B, \psi] \circ [A, \varphi] = [B \natural A, \psi \circ (\text{id}_B \natural \varphi)]$. Note that this may also be written as the colimit

$$\text{Hom}_{\langle \mathcal{G}, \mathcal{M} \rangle}(X, Y) = \text{colim}_{\mathcal{G}} [\text{Hom}_{\mathcal{M}}(- \natural X, Y)].$$

Remark 3.2 There is a canonical faithful functor $c_{\langle \mathcal{G}, \mathcal{M} \rangle} : \mathcal{M} \hookrightarrow \langle \mathcal{G}, \mathcal{M} \rangle$ defined as the identity on objects and sending $\phi \in \text{Hom}_{\mathcal{M}}(X, Y)$ to $[0, \phi]$.

Example 3.3 A topological monoidal groupoid \mathcal{G} has a continuous left-action on itself given by its monoidal structure, so we may always take $\mathcal{M} = \mathcal{G}$ and consider the category $\langle \mathcal{G}, \mathcal{G} \rangle$.

Notation 3.4 As an abbreviation, we denote the category $\langle \mathcal{G}, \mathcal{G} \rangle$ by $\mathfrak{U}\mathcal{G}$.

If the categories \mathcal{G} and \mathcal{M} are both *discrete*, then Definition 3.1 recovers the classical *bracket construction* of Quillen. This is a particular case of a more general construction of [Gra76, p.219]. Assuming in addition that \mathcal{M} is a groupoid (as for all the examples discussed in this paper cf. §4), following mutatis mutandis [RW17, Proposition 1.7], if $(\mathcal{G}, \natural, 0)$ has *no zero divisors* – meaning that $A \natural B \cong 0$ if and only if $A \cong B \cong 0$ for all objects A and B of \mathcal{G} – and if $\text{Aut}_{\mathcal{G}}(0) = \{\text{id}_0\}$, then \mathcal{M} is isomorphic to the maximal subgroupoid of $\langle \mathcal{G}, \mathcal{M} \rangle$ (i.e. the subcategory which has the same objects as $\langle \mathcal{G}, \mathcal{M} \rangle$ and of which the morphisms are the isomorphisms of $\langle \mathcal{G}, \mathcal{M} \rangle$). We record this for future use:

Lemma 3.5 *Suppose that \mathcal{G} and \mathcal{M} are discrete groupoids, that \mathcal{G} has no zero divisors and that its monoidal unit has no non-trivial automorphisms. Then the canonical functor of Remark 3.2 above is an isomorphism from \mathcal{M} onto the maximal subgroupoid of $\langle \mathcal{G}, \mathcal{M} \rangle$.*

Under these assumptions, if \mathcal{M} is a groupoid with non-negative integers as objects and a family of groups $\{G_n\}_{n \in \mathbb{N}}$ as isomorphisms, then Quillen’s bracket construction consists in just “artificially” adding to \mathcal{M} morphisms which go from n to $n + 1$: this justifies the use of this construction as source category to encode *compatible* representations of families of groups.

Remark 3.6 A topological version of Quillen’s bracket construction is mentioned briefly in Remark 2.10 of [Kra19], although there the categories are *topological* in the sense of being categories internal to the category of topological spaces, rather than topologically-enriched categories. Lemma 3.8 below is stated for topologically-enriched categories, but it is likely that it has an analogue for categories internal to the category of topological spaces, in which case Lemma 2.11 of [Kra19] would be a particular case of this analogue.

This construction is of course *functorial* in \mathcal{M} and \mathcal{G} in an appropriate sense. We mention some properties of this functoriality that we will need.

Lemma 3.7 *Let \mathcal{D} be a topological monoidal groupoid and let $\mathcal{G}_1 \subseteq \mathcal{G}_2 \subseteq \mathcal{D}$ and $\mathcal{M}_1 \subseteq \mathcal{M}_2 \subseteq \mathcal{D}$ be subgroupoids such that, for $i = 1, 2$, \mathcal{G}_i is closed under $-\natural-$ and \mathcal{M}_i is closed under $g \natural -$ for each object g of \mathcal{G}_i . Then there is a canonical functor $\langle \mathcal{G}_1, \mathcal{M}_1 \rangle \rightarrow \langle \mathcal{G}_2, \mathcal{M}_2 \rangle$. Moreover,*

- if $\mathcal{G}_1 = \mathcal{G}_2 = \mathcal{G}$, the functor $\langle \mathcal{G}, \mathcal{M}_1 \rangle \rightarrow \langle \mathcal{G}, \mathcal{M}_2 \rangle$ is an inclusion of a subcategory, which is full if the inclusion $\mathcal{M}_1 \subseteq \mathcal{M}_2$ is full;
- if $\mathcal{M}_1 = \mathcal{M}_2 = \mathcal{M}$, the functor $\langle \mathcal{G}_1, \mathcal{M} \rangle \rightarrow \langle \mathcal{G}_2, \mathcal{M} \rangle$ is the identity on objects, and is faithful – thus an inclusion of a subcategory – if the inclusion $\mathcal{G}_1 \subseteq \mathcal{G}_2$ is full.

In particular, there is a canonical functor $\langle \mathcal{G}_1, \mathcal{M}_1 \rangle \rightarrow \mathfrak{UD}$, which is an inclusion of a subcategory if the inclusion $\mathcal{G}_1 \subseteq \mathcal{D}$ is full.

Proof. The objects of $\langle \mathcal{G}_i, \mathcal{M}_i \rangle$ are the objects of \mathcal{M}_i , so we define the functor on objects as the inclusion $\text{ob}(\mathcal{M}_1) \hookrightarrow \text{ob}(\mathcal{M}_2)$. A morphism in $\langle \mathcal{G}_i, \mathcal{M}_i \rangle$ from X to Y is represented by a choice of object A of \mathcal{G}_i and a morphism $A \downarrow X \rightarrow Y$ of \mathcal{M}_i . We may therefore send such a morphism, for $i = 1$, to the morphism, for $i = 2$, represented by the same data, since $\mathcal{G}_1 \subseteq \mathcal{G}_2$ and $\mathcal{M}_1 \subseteq \mathcal{M}_2$. It is straightforward to check that this assignment respects the defining equivalence relation, so induces a continuous map of morphism spaces, and that it also respects composition and identities. The statements in the two bullet points may be verified by unwinding the definition of morphisms in $\langle \mathcal{G}_i, \mathcal{M}_i \rangle$. \square

Lemma 3.8 *Let \mathcal{G} be a topological monoidal groupoid and \mathcal{M} a topological category with a continuous left-action of \mathcal{G} . Assume that, for each object A of \mathcal{G} and each pair of objects X, Y of \mathcal{M} , the quotient map*

$$\text{Hom}_{\mathcal{M}}(A \downarrow X, Y) \longrightarrow \text{Hom}_{\mathcal{M}}(A \downarrow X, Y) / \text{Aut}_{\mathcal{G}}(A) \quad (3.1)$$

is a Serre fibration. Then there is a canonical isomorphism of categories

$$\pi_0(\langle \mathcal{G}, \mathcal{M} \rangle) \cong \langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle. \quad (3.2)$$

Proof. First note that $\pi_0(\langle \mathcal{G}, \mathcal{M} \rangle)$ and $\langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle$ have the same object set, by the definition of the discrete and topologically-enriched Quillen bracket constructions, and the functor π_0 . Specifically, their common object set is $\text{ob}(\mathcal{M})$. It therefore remains to show that, for objects X and Y of \mathcal{M} , there is a natural bijection between $\pi_0(\text{Hom}_{\langle \mathcal{G}, \mathcal{M} \rangle}(X, Y))$ and $\text{Hom}_{\langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle}(X, Y)$. Set

$$\Phi = \bigsqcup_{A \in \text{ob}(\mathcal{G})} \text{Hom}_{\mathcal{M}}(A \downarrow X, Y).$$

Unravelling the definitions, what we need to prove is that there is a natural bijection

$$\pi_0(\Phi / \sim_t) \cong \pi_0(\Phi) / \sim_h,$$

where \sim_t is the equivalence relation given by $(A, \varphi) \sim_t (A', \varphi')$ if and only if there is a morphism $\sigma \in \text{Hom}_{\mathcal{G}}(A, A')$ such that $\varphi = \varphi' \circ (\sigma \downarrow \text{id}_A)$, and \sim_h is the equivalence relation given by $(A, [\varphi]) \sim_h (A', [\varphi'])$ if and only if there is a morphism $\sigma \in \text{Hom}_{\mathcal{G}}(A, A')$ such that $\varphi \simeq \varphi' \circ (\sigma \downarrow \text{id}_A)$. Note that the only difference between these definitions is that the equality is replaced by a homotopy in the definition of \sim_h . As sets, these are both quotients of (the underlying set of) Φ , so we just need to show that, given two elements (A, φ) and (A', φ') of Φ , they have the same image in $\pi_0(\Phi / \sim_t)$ if and only if they have the same image in $\pi_0(\Phi) / \sim_h$.

(a) Suppose first that (A, φ) and (A', φ') have the same image in $\pi_0(\Phi) / \sim_h$. This means that there is a morphism $\sigma \in \text{Hom}_{\mathcal{G}}(A, A')$ and a path $\gamma: [0, 1] \rightarrow \text{Hom}_{\mathcal{M}}(A \downarrow X, Y) \subseteq \Phi$ with $\gamma(0) = (A, \varphi)$ and $\gamma(1) = (A', \varphi' \circ (\sigma \downarrow \text{id}_A))$. Composing with the projection $\Phi \rightarrow \Phi / \sim_t$ and writing $[-]_t$ for the equivalence classes with respect to \sim_t , we obtain a path in Φ / \sim_t from $[(A, \varphi)]_t$ to $[(A', \varphi' \circ (\sigma \downarrow \text{id}_A))]_t = [(A', \varphi')]_t$. Hence (A, φ) and (A', φ') have the same image in $\pi_0(\Phi / \sim_t)$.

(b) To prove the converse, we first make an assumption, which we will justify later. Namely, we assume that quotient map $q: \Phi \rightarrow \Phi / \sim_t$ is a Serre fibration. Now assume that (A, φ) and (A', φ') have the same image in $\pi_0(\Phi / \sim_t)$, so there is a path $\delta: [0, 1] \rightarrow \Phi / \sim_t$ with $\delta(0) = [(A, \varphi)]_t$ and $\delta(1) = [(A', \varphi')]_t$. By our assumption that q is a Serre fibration, we may lift this to a path $\varepsilon: [0, 1] \rightarrow \Phi$ with $\varepsilon(0) = (A, \varphi)$ and $\varepsilon(1) \sim_t (A', \varphi')$. Its image $\varepsilon([0, 1])$ is path-connected, so it must lie in $\text{Hom}_{\mathcal{M}}(A \downarrow X, Y) \subseteq \Phi$. Hence we have a path $\varepsilon: [0, 1] \rightarrow \text{Hom}_{\mathcal{M}}(A \downarrow X, Y)$ with $\varepsilon(0) = (A, \varphi)$ and $\varepsilon(1) = (A, \varphi'') \sim_t (A', \varphi')$, for some $\varphi'' \in \text{Hom}_{\mathcal{M}}(A \downarrow X, Y)$. The relation $(A, \varphi'') \sim_t (A', \varphi')$ means that there is a morphism $\sigma \in \text{Hom}_{\mathcal{G}}(A, A')$ such that $\varphi'' = \varphi' \circ (\sigma \downarrow \text{id}_A)$. Hence ε is a homotopy witnessing that $\varphi \simeq \varphi' \circ (\sigma \downarrow \text{id}_A)$, so we have shown that $(A, [\varphi]) \sim_h (A', [\varphi'])$, in other words, (A, φ) and (A', φ') have the same image in $\pi_0(\Phi) / \sim_h$.

(c) It now just remains to prove our earlier assumption that q is a Serre fibration. Directly from the definition, one may easily verify the following two facts:

- $\bigsqcup_i f_i: \bigsqcup_i E_i \rightarrow B$ is a Serre fibration if and only if each $f_i: E_i \rightarrow B$ is a Serre fibration.
- $f: E \rightarrow B$ is a Serre fibration if and only if $f(E)$ is a union of path-components of B and $f: E \rightarrow f(E)$ is a Serre fibration.

It therefore suffices to prove that

- (i) $q(\mathrm{Hom}_{\mathcal{M}}(A \natural X, Y))$ is a union of path-components of Φ/\sim_t for each $A \in \mathrm{ob}(\mathcal{G})$,
- (ii) $\mathrm{Hom}_{\mathcal{M}}(A \natural X, Y) \rightarrow q(\mathrm{Hom}_{\mathcal{M}}(A \natural X, Y))$ is a Serre fibration for each $A \in \mathrm{ob}(\mathcal{G})$.

Let us partition $\mathrm{ob}(\mathcal{G})$ into equivalence classes \mathcal{O}_α , under the equivalence relation where two objects A, A' of \mathcal{G} are equivalent if and only if there is a morphism $A \rightarrow A'$ in \mathcal{G} . This is an equivalence relation since \mathcal{G} is a groupoid. We may then write $\Phi = \bigsqcup_\alpha \Phi_\alpha$, where

$$\Phi_\alpha = \bigsqcup_{A \in \mathcal{O}_\alpha} \mathrm{Hom}_{\mathcal{M}}(A \natural X, Y).$$

The equivalence relation \sim_t on Φ clearly preserves the topological disjoint union $\bigsqcup_\alpha \Phi_\alpha$, so we have

$$\Phi/\sim_t = \bigsqcup_\alpha (\Phi_\alpha/\sim_t).$$

Also note that, for any two objects $A, A' \in \mathcal{O}_\alpha$ (for fixed α), we have $q(\mathrm{Hom}_{\mathcal{M}}(A \natural X, Y)) = q(\mathrm{Hom}_{\mathcal{M}}(A' \natural X, Y))$. So, if we make a choice of object $A_\alpha \in \mathcal{O}_\alpha$ for each α , we have a decomposition of Φ/\sim_t as a topological disjoint union:

$$\Phi/\sim_t = \bigsqcup_\alpha q(\mathrm{Hom}_{\mathcal{M}}(A_\alpha \natural X, Y)).$$

This immediately implies point (i) above.

For point (ii), note that two elements $\varphi, \varphi' \in \mathrm{Hom}_{\mathcal{M}}(A \natural X, Y)$ have the same image under q if and only if they are \sim_t -equivalent, which is equivalent to saying that they lie in the same orbit of the $\mathrm{Aut}_{\mathcal{G}}(A)$ -action on $\mathrm{Hom}_{\mathcal{M}}(A \natural X, Y)$. Hence the map

$$q_A: \mathrm{Hom}_{\mathcal{M}}(A \natural X, Y) \longrightarrow q(\mathrm{Hom}_{\mathcal{M}}(A \natural X, Y)) \quad (3.3)$$

is isomorphic to (3.1), at least on underlying sets. If we can show that they are isomorphic also as continuous maps of spaces, then we will be done, since we know by hypothesis that (3.1) is a Serre fibration. Since (3.1) and (3.3) are surjective continuous maps with the same domain and the same point-fibres, and we know moreover that (3.1) is a quotient map, it suffices to prove that (3.3) is also a quotient map.

Let $U \subseteq q(\mathrm{Hom}_{\mathcal{M}}(A \natural X, Y))$ be a subset such that $q_A^{-1}(U)$ is open in $\mathrm{Hom}_{\mathcal{M}}(A \natural X, Y)$. We need to show that U is open in $q(\mathrm{Hom}_{\mathcal{M}}(A \natural X, Y))$. To see this, let $A \in \mathcal{O}_\alpha$ and note that, by the fact discussed above that the equivalence relation \sim_t preserves the decomposition of Φ into a topological disjoint union, the restriction

$$q_\alpha = q|_{\Phi_\alpha}: \Phi_\alpha \longrightarrow q(\Phi_\alpha) = q(\mathrm{Hom}_{\mathcal{M}}(A \natural X, Y))$$

is a quotient map. So it suffices to show that $q_\alpha^{-1}(U)$ is open in Φ_α . Now, from the definitions, we observe the following description of the subset

$$q_\alpha^{-1}(U) \subseteq \Phi_\alpha = \bigsqcup_{A' \in \mathcal{O}_\alpha} \mathrm{Hom}_{\mathcal{M}}(A' \natural X, Y).$$

For each object $A' \in \mathcal{O}_\alpha$, choose an isomorphism $\sigma_{A'}: A' \rightarrow A$ in \mathcal{G} . This induces a homeomorphism

$$\Upsilon_{A'} = - \circ (\sigma_{A'} \natural \mathrm{id}): \mathrm{Hom}_{\mathcal{M}}(A \natural X, Y) \longrightarrow \mathrm{Hom}_{\mathcal{M}}(A' \natural X, Y).$$

Then we have

$$q_\alpha^{-1}(U) = \bigsqcup_{A' \in \mathcal{O}_\alpha} \Upsilon_{A'}(q_A^{-1}(U)).$$

Since $q_A^{-1}(U)$ is open in $\mathrm{Hom}_{\mathcal{M}}(A \natural X, Y)$, it follows that $\Upsilon_{A'}(q_A^{-1}(U))$ is open in $\mathrm{Hom}_{\mathcal{M}}(A' \natural X, Y)$ for each $A' \in \mathcal{O}_\alpha$. Thus $q_\alpha^{-1}(U)$ is open in Φ_α , as required. \square




Figure 3.1 An illustration of the notation for the solid cylinder \mathbb{B}_t^d from Notation 3.9 for $d = 3$. Its lower boundary $\partial_\ell \mathbb{B}_t^3$ is coloured yellow, its base $b\mathbb{B}_t^3$ is yellow-green and the codimension-2 stratum $\partial b\mathbb{B}_t^3 = \partial \mathbb{D}^2 \times \{0\}$ is blue.




Figure 3.2 Two decorated manifolds and their boundary connected sum.

3.2. Topological groupoids of decorated manifolds

Fix an integer $d \geq 2$. First, we define the notion of *decorated manifolds* and their morphisms. The idea is that the groups $\pi_0(\text{Diff}(-))$ of decorated manifolds will contain all motion groups as normal subgroups; cf. Remark 4.24.

Notation 3.9 (*Solid cylinders.*) We denote by \mathbb{D}^{d-1} the closed unit $(d-1)$ -dimensional disc in \mathbb{R}^{d-1} . For a real number $t > 0$, we will write $\mathbb{B}_t^d = \mathbb{D}^{d-1} \times [0, t]$ for the *solid d -dimensional cylinder of height t* . We will also write

$$\partial_\ell \mathbb{B}_t^d = (\partial \mathbb{D}^{d-1} \times [0, t]) \cup (\mathbb{D}^{d-1} \times \{0\})$$

and call this the *lower boundary* of \mathbb{B}_t^d , as well as $b\mathbb{B}_t^d = \mathbb{D}^{d-1} \times \{0\}$ and call this the *base* of \mathbb{B}_t^d . This is illustrated in Figure 3.1.

Definition 3.10 (*Boundary-cylinders.*) Let M be a smooth d -manifold. A *boundary-cylinder* for M is a topological embedding $e: \mathbb{B}_1^d \hookrightarrow M$ such that $e^{-1}(\partial M) = \partial_\ell \mathbb{B}_1^d$ and e is a smooth embedding except on the $(d-2)$ -sphere $\partial b\mathbb{B}_1^d$. Two boundary-cylinders e, e' are *equivalent* if they are equal when restricted to $\mathbb{B}_\epsilon^d \subseteq \mathbb{B}_1^d$ for some $\epsilon > 0$. An equivalence class of boundary-cylinders is called a *boundary-cylinder germ*.

Definition 3.11 (*Decorated manifolds.*) A *decorated manifold* is a smooth d -manifold M , equipped with a closed submanifold $A \subset \text{int}(M)$ and a pair (e_1, e_2) of boundary-cylinder germs for $M \setminus A$ such that $e_1(b\mathbb{B}_1^d)$ and $e_2(b\mathbb{B}_1^d)$ are disjoint. See Figure 3.2 for a schematic picture.

Definition 3.12 (*Morphisms of decorated manifolds*) A morphism of decorated manifolds from (M, A, e_1, e_2) to (M', A', e'_1, e'_2) is a smooth, proper (preimages of compact subspaces are compact) map $\varphi: M \rightarrow M'$ such that $\varphi(A) \subseteq A'$ and such that, for some $\epsilon > 0$ and for each $i \in \{1, 2\}$, we have $\varphi(e_i(\mathbb{B}_\epsilon^d)) = e'_i(\mathbb{B}_\epsilon^d)$ and the composition $(e'_i)^{-1} \circ \varphi \circ e_i: \mathbb{B}_\epsilon^d \rightarrow \mathbb{B}_\epsilon^d$ is the identity. Write $C_{\text{dec}}^\infty(M, M')$ for the set of such morphisms, where by abuse of notation we are abbreviating (M, A, e_1, e_2) to M and (M', A', e'_1, e'_2) to M' .

Definition 3.13 (*Morphism spaces.*) The set $C_{\text{dec}}^\infty(M, M')$ is topologised as a colimit of Whitney topologies, as follows. First choose representative boundary cylinders for the boundary-cylinder germs e_i and e'_i . This ensures that the condition in Definition 3.12 makes sense for a fixed $\epsilon \in (0, 1)$, not just for an unspecified $\epsilon \in (0, 1)$ that is quantified over.

Now fix $\epsilon \in (0, 1)$ and write $C_{\text{dec}, \epsilon}^\infty(M, M')$ for the subset of $C_{\text{dec}}^\infty(M, M')$ where the condition in Definition 3.12 holds for this fixed ϵ . Equip each subset $C_{\text{dec}, \epsilon}^\infty(M, M')$ with the subspace topology induced from the smooth Whitney topology on the set $C^\infty(M, M')$ of all smooth maps from M to M' ; for details of the Whitney topology, see for example [Hir76, Chapter 2]. As a set, $C_{\text{dec}}^\infty(M, M')$ is the union of $C_{\text{dec}, \epsilon}^\infty(M, M')$ over all choices of $\epsilon \in (0, 1)$; we equip $C_{\text{dec}}^\infty(M, M')$ with the colimit topology induced by the increasing filtration $\{C_{\text{dec}, \epsilon}^\infty(M, M')\}_{\epsilon \in (0, 1)}$.

Different choices of representative boundary-cylinders for the boundary-cylinder germs e_i and e'_i will result in different filtrations. However, any two such filtrations are cofinal in each other, so the colimit topology induced on $C_{\text{dec}}^\infty(M, M')$ does not depend on this choice.

Note also that this topology may differ from the subspace topology that inherited directly from the Whitney topology on $C^\infty(M, M')$ (the colimit topology may be finer). However, these two topologies on $C_{\text{dec}}^\infty(M, M')$ are weakly equivalent. In particular, they have the same π_0 .

Definition 3.14 (*Boundary connected sum.*) Let (M, A, e_1, e_2) and (M', A', e'_1, e'_2) be two decorated d -manifolds. Define

$$M \natural M' = (M \sqcup M') / \sim,$$

where \sim is the equivalence relation generated by $e_2(x, 0) \sim e'_1(x, 0)$ for all $(x, 0) \in b\mathbb{B}_1^d$. We give this a smooth structure as follows. There are obvious topological embeddings

$$M \setminus e_2(b\mathbb{B}_1^d) \hookrightarrow M \natural M' \quad \text{and} \quad M' \setminus e'_1(b\mathbb{B}_1^d) \hookrightarrow M \natural M',$$

and another topological embedding

$$\mathbb{D}^{d-1} \times [-1, 1] \hookrightarrow M \natural M'$$

given by $(x, t) \mapsto e_2(x, -t)$ for $t \leq 0$ and $(x, t) \mapsto e'_1(x, t)$ for $t \geq 0$, where we have implicitly chosen representative boundary-cylinders for the boundary-cylinder germs e_2 and e'_1 . We define a smooth structure on $M \natural M'$ by declaring that these are both *smooth* embeddings.

This is a well-defined smooth structure, since:

- the smooth structures induced by these embeddings are compatible on intersections, due to the fact that boundary-cylinders are *smooth* embeddings away from $\partial b\mathbb{B}_1^d$;
- the smooth structure of $M \natural M'$ is determined, except on $e_2(b\mathbb{B}_1^d) = e'_1(b\mathbb{B}_1^d)$, by the smooth structures of M and M' . The embedding of $\mathbb{D}^{d-1} \times [-1, 1]$ induced by (boundary-cylinders representing the boundary-cylinder germs) e_2 and e'_1 is therefore only required to extend this smooth structure to $e_2(b\mathbb{B}_1^d) = e'_1(b\mathbb{B}_1^d)$. As a result, the smooth structure does not depend on the choice of representative boundary-cylinders, but only their germs.

Finally, we define:

$$(M, A, e_1, e_2) \natural (M', A', e'_1, e'_2) = (M \natural M', A \sqcup A', e_1, e'_2).$$

See Figure 3.2 for a schematic illustration.

Remark 3.15 The usual definition of boundary connected sum of two smooth manifolds M, M' depends on a choice of embedded disc in the boundary of each manifold, and a method of “straightening corners” after gluing these discs together. Up to diffeomorphism, the resulting smooth manifold $M \natural M'$ depends only on the choice of a boundary-component of M and of M' , and orientations of these if they are orientable (this is a result of Palais’ *Disc theorem* [Pal60a, Theorem B and Corollary 1] and the existence of collar neighbourhoods). However, in order for \natural to induce a well-defined monoidal structure on some category of manifolds with boundary (which we will do just below), it must be well-defined on the nose, not just up to diffeomorphism (since objects are manifolds, not diffeomorphism-classes of manifolds). The definition of decorated manifolds is designed so that these additional choices are *built in*, and no additional choices are required in Definition 3.14 above.

Definition 3.16 (*Decorated manifold categories.*) Let $\mathcal{D}ec_d$ be the topological category defined as follows. Its objects are all *decorated manifolds* (M, A, e_1, e_2) of dimension d , as in Definition 3.11. The space of morphisms from $M = (M, A, e_1, e_2)$ to $M' = (M', A', e'_1, e'_2)$ is the space $C_{\text{dec}}^\infty(M, M')$ defined in Definition 3.12 and topologised in Definition 3.13. Note that composition is continuous in this topology, since it is a colimit of Whitney topologies and composition of smooth, proper maps is continuous in the Whitney topology.

Let \mathcal{D}_d be the underlying topological groupoid of $\mathcal{D}ec_d$. In other words, its objects are all decorated manifolds of dimension d and its morphisms are those morphisms $(M, A, e_1, e_2) \rightarrow (M', A', e'_1, e'_2)$ of decorated manifolds whose underlying smooth map $\varphi: M \rightarrow M'$ is a diffeomorphism and $\varphi(A) = A'$.

The boundary connected sum of Definition 3.14 induces a semi-monoidal structure on $\mathcal{D}ec_d$, and hence on \mathcal{D}_d :

Definition 3.17 (*Semi-monoidal structure.*) We define the functor

$$\natural: \mathcal{D}ec_d \times \mathcal{D}ec_d \longrightarrow \mathcal{D}ec_d$$

on objects via the boundary connected sum of Definition 3.14. Now suppose we are given morphisms $\varphi: (L, A, e_1, e_2) \rightarrow (L', A', e'_1, e'_2)$ and $\psi: (M, B, f_1, f_2) \rightarrow (M', B', f'_1, f'_2)$ in $\mathcal{D}ec_d$. By definition, these are smooth, proper maps $L \rightarrow L'$ and $M \rightarrow M'$ that take A and B into A' and B' respectively, and are compatible with the given boundary-cylinder germs. This compatibility implies that they glue to a well-defined, smooth map $L \natural M \rightarrow L' \natural M'$, which is moreover a morphism

$$(L, A, e_1, e_2) \natural (M, B, f_1, f_2) \longrightarrow (L', A', e'_1, e'_2) \natural (M', B', f'_1, f'_2)$$

of $\mathcal{D}ec_d$. It is then easily checked that this gives $\mathcal{D}ec_d$ the structure of a topological semi-monoidal groupoid; cf. §3.4.

Remark 3.18 (*Monoidal and semi-monoidal structures.*) This is a somewhat technical side remark to explain certain choices in our definitions. The semi-monoidal structure \natural on $\mathcal{D}ec_d$ does not have a unit, in other words, it is not a monoidal structure, since there is no *natural* way of identifying $M \natural \mathbb{B}_1^d$ with M for all M (although they are of course non-naturally diffeomorphic). This non-naturality is essentially because decorated manifolds come equipped with *germs* of boundary-cylinders (and morphisms preserve *germs* of boundary-cylinders), rather than actual boundary-cylinders.

One possible way to fix this issue would be to redefine decorated manifolds to be equipped with boundary-cylinders (not just germs of such) and morphisms of decorated manifolds to preserve these boundary-cylinders. Then, one would be able to define the boundary connected sum $M \natural M'$ of two decorated manifolds M and M' by identifying $e_2(\mathbb{B}_1^d)$ with $e'_1(\mathbb{B}_1^d)$ via the homeomorphism $e_2(x, t) \mapsto e'_1(x, 1-t)$ – in other words, by gluing together whole boundary-cylinders, instead of just their bases. Under this definition, there are obvious natural isomorphisms $M \natural \mathbb{B}_1^d \cong M \cong \mathbb{B}_1^d \natural M$, so the topological category defined in this way would be monoidal, not just semi-monoidal. However, the proof of Proposition 3.24 below, which tells us that the Serre fibration hypothesis of Lemma 3.8 is satisfied for subgroupoids of \mathcal{D}_d (cf. Lemma 4.4), depends crucially on the fact that morphisms of decorated manifolds are only required to preserve *germs* of boundary-cylinders, rather than boundary-cylinders themselves. Thus, this alternative definition would fix the lack of units in the monoidal structure at the expense of breaking a key property that we will need.

Another possible fix would be simply to formally adjoin a unit object to $\mathcal{D}ec_d$ by taking the disjoint union with the trivial category on one object 1 and extending \natural by defining $c \natural 1 = 1 \natural c = c$ for any object c and $\text{id}_1 \natural \text{id}_1 = \text{id}_1$. However, this is not a very natural thing to do, since the discrete semi-monoidal category $\pi_0(\mathcal{D}ec_d)$ *does* have a unit object (see Lemma 3.20 below). Adjoining a formal unit object to $\mathcal{D}ec_d$ would correspond, on π_0 , to forgetting this “natural” unit object and adjoining a new, formal unit object instead.

Instead of trying to force $\mathcal{D}ec_d$ to have a unit object, we will simply work with it as a topological *semi-monoidal* category. As explained in §3.4 below, the end result of this lack of units will be that the output of our general construction will be a *semi-functor* from whichever category \mathcal{C} we are interested in to the category $\bullet\text{Mod}\bullet$. In each example where we apply this, one may trivially check that this semi-functor does in fact preserve identities, and is therefore a functor.

Finally we can analogously define a topological groupoid of *oriented* decorated manifolds:

Definition 3.19 Let \mathcal{D}_d^+ denote the topological groupoid whose objects are decorated d -manifolds (M, A, e_1, e_2) together with an orientation of $A \subset \text{int}(M)$, and whose morphisms are diffeomorphisms φ as in Definition 3.16 such that the restriction $\varphi|_A: A \rightarrow A'$ is an orientation-preserving diffeomorphism. The boundary connected sum for such decorated d -manifolds is defined as in Definition 3.14, with the orientation for $A \sqcup B$ being induced from those of A and B . This then extends, just as in Definition 3.17, to a structure of a topological semi-monoidal groupoid on \mathcal{D}_d^+ .

One may of course similarly define \mathcal{D}_d^θ for any type of tangential structure $\theta: X \rightarrow BO$, equipping $A \subset \text{int}(M)$ with a θ -structure and requiring this to be preserved by morphisms φ . Applying π_0 to all morphism spaces, we may consider the discrete groupoids $\pi_0(\mathcal{D}_d)$ and $\pi_0(\mathcal{D}_d^+)$. The following lemma may be easily checked from the definitions.

Lemma 3.20 *The discrete groupoids $\pi_0(\mathcal{D}_d)$ and $\pi_0(\mathcal{D}_d^+)$ inherit well-defined semi-monoidal structures from those on \mathcal{D}_d and \mathcal{D}_d^+ . Moreover, these semi-monoidal structures are monoidal, with unit object given in each case by the solid cylinder $(\mathbb{B}_1^d, \emptyset, \text{id}, r)$, where $r: \mathbb{B}_1^d \rightarrow \mathbb{B}_1^d$ is the reflection $(x, t) \mapsto (x, 1 - t)$.*

More generally, if $\mathcal{G} \subseteq \mathcal{D}_d$ is any subgroupoid closed under the semi-monoidal structure and containing the solid cylinder $(\mathbb{B}_1^d, \emptyset, \text{id}, r)$, then the semi-monoidal structure inherited by $\pi_0(\mathcal{G})$ is monoidal. Similarly for subgroupoids $\mathcal{G} \subseteq \mathcal{D}_d^+$.

3.3. The Serre fibration condition

We now prove a technical result that will imply that the condition (3.1) is true in a very general setting, covering all of our examples.

Definition 3.21 (*Decorated diffeomorphisms.*) A *diffeomorphism* of decorated manifolds (or a *decorated diffeomorphism*) is a morphism of decorated manifolds (cf. Definition 3.12) that admits an inverse. Write $\text{Diff}_{\text{dec}}(M, N)$ for the space of decorated diffeomorphisms $M \rightarrow N$ and abbreviate $\text{Diff}_{\text{dec}}(M) = \text{Diff}_{\text{dec}}(M, M)$. This is topologised as a subspace of $C_{\text{dec}}^\infty(M, N)$, which is topologised as a colimit of Whitney topologies (cf. Definition 3.13).

Definition 3.22 (*Decorated embeddings.*) Let $M = (M, A)$ and $N = (N, B)$ be decorated manifolds. Define $\text{Emb}_{\text{dec}}(M, N)$ to be the space of smooth, proper embeddings $\varphi: M \hookrightarrow N$, equipped with a germ of an extension φ' to an embedding $\mathbb{B}_1^d \natural M \hookrightarrow N$, such that

- $\varphi(A) \subseteq B$;
- for some $\epsilon > 0$ we have $\varphi(e_2(\mathbb{B}_\epsilon^d)) = e_2'(\mathbb{B}_\epsilon^d)$ and $(e_2')^{-1} \circ \varphi \circ e_2$ is the identity map $\mathbb{B}_\epsilon^d \rightarrow \mathbb{B}_\epsilon^d$, where e_1, e_2 are the boundary-cylinder germs of (M, A) and e_1', e_2' are those of (N, B) . This is similar to Definition 3.12, except that we only require the condition on e_2 , not on e_1 ;
- there is a decorated manifold M' and diffeomorphism of decorated manifolds $\bar{\varphi}: M' \natural M \rightarrow N$ such that $\varphi = \bar{\varphi} \circ \iota_{M, M'}$, where $\iota_{M, M'}$ denotes the canonical embedding of M into $M' \natural M$. This extension of φ to $\bar{\varphi}$ should be compatible with the given germ of an extension φ' of φ .

This space is topologised as a colimit of Whitney topologies, analogously to Definition 3.13.

Definition 3.23 For decorated manifolds L, M, N , denote by $\text{Emb}_{\text{dec}}(M, N)_L$ the subspace of $\text{Emb}_{\text{dec}}(M, N)$ of those embeddings for which we may take $M' = L$ in the third point of Definition 3.22. Note that $\text{Emb}_{\text{dec}}(M, N)$ decomposes as a topological disjoint union:

$$\text{Emb}_{\text{dec}}(M, N) \cong \bigsqcup_L \text{Emb}_{\text{dec}}(M, N)_L, \quad (3.4)$$

where the disjoint union runs over representatives of isomorphism classes of decorated manifolds.

Let L and M be decorated manifolds. There is a continuous right action of $\text{Diff}_{\text{dec}}(L)$ on $\text{Diff}_{\text{dec}}(L \natural M)$ given by $\varphi \cdot \psi = \varphi \circ (\psi \natural \text{id}_M)$, and hence a quotient map

$$\Psi: \text{Diff}_{\text{dec}}(L \natural M) \longrightarrow \text{Diff}_{\text{dec}}(L \natural M) / \text{Diff}_{\text{dec}}(L). \quad (3.5)$$




Figure 3.3 The boundary connected sum $L\sharp M$ from the proof of Proposition 3.24.

Proposition 3.24 *The quotient map (3.5) is a Serre fibration. There is a homeomorphism between its codomain and $\text{Emb}_{\text{dec}}(M, L\sharp M)_L$, induced by the restriction map.*

Remark 3.25 This is related to results of Cerf [Cer61, Corollaire 2, §II.2.2.2, page 294], Palais [Pal60b, Theorem B] and Lima [Lim63], but we were not able to find an instance of their results that covers exactly the setting that we need here. We therefore give a complete proof of Proposition 3.24 below, using as an input two key results of Cerf and Palais, namely Lemme II.2.1.2 (page 291) of [Cer61] and [Pal60b, Theorem A].

Proof of Proposition 3.24. The decorated manifolds $L = (L, A, e_1, e_2)$ and $M = (M, B, e'_1, e'_2)$ come equipped with germs e_1, e_2, e'_1, e'_2 of boundary cylinders; let us once and for all choose representative boundary cylinders for these germs, and denote them by the same symbols, by abuse of notation.

For $\epsilon \in (0, 1)$, let $\text{Diff}_{\text{dec}, \epsilon}(L\sharp M)$ denote the group of self-diffeomorphisms of $L\sharp M$ sending $A \sqcup B$ onto itself and restricting to the identity on $e_1(\mathbb{B}_\epsilon^d)$ and on $e'_2(\mathbb{B}_\epsilon^d)$. If we give this the Whitney topology, then

$$\text{Diff}_{\text{dec}}(L\sharp M) \cong \text{colim}_{\epsilon \rightarrow 0}(\text{Diff}_{\text{dec}, \epsilon}(L\sharp M)), \quad (3.6)$$

by our definition of morphisms of decorated manifolds (*cf.* Definition 3.12) and the topology on morphism spaces (*cf.* Definition 3.13). Similarly, for each $\epsilon, t \in (0, 1)$, let $\text{Diff}_{\text{dec}, \epsilon, t}(L)$ denote the subgroup of $\text{Diff}_{\text{dec}, \epsilon}(L\sharp M)$ consisting of diffeomorphisms that restrict to the identity on the submanifold $M_t = M \cup e_2(\mathbb{B}_t^d)$ of $L\sharp M$ pictured in Figure 3.3. We have a quotient map

$$\Psi_{\epsilon, t}: \text{Diff}_{\text{dec}, \epsilon}(L\sharp M) \longrightarrow \text{Diff}_{\text{dec}, \epsilon}(L\sharp M) / \text{Diff}_{\text{dec}, \epsilon, t}(L).$$

For any $\epsilon, \epsilon', t, t' \in (0, 1)$ with $\epsilon' \leq \epsilon$ and $t' \leq t$ there are natural maps

$$\text{Diff}_{\text{dec}, \epsilon}(L\sharp M) / \text{Diff}_{\text{dec}, \epsilon, t}(L) \longrightarrow \text{Diff}_{\text{dec}, \epsilon'}(L\sharp M) / \text{Diff}_{\text{dec}, \epsilon', t'}(L),$$

so we may take the directed colimit of the maps $\Psi_{\epsilon, t}$ to obtain

$$\text{colim}_{\epsilon, t \rightarrow 0}(\Psi_{\epsilon, t}): \text{Diff}_{\text{dec}}(L\sharp M) \longrightarrow \text{colim}_{\epsilon, t \rightarrow 0}(\text{Diff}_{\text{dec}, \epsilon}(L\sharp M) / \text{Diff}_{\text{dec}, \epsilon, t}(L)),$$

where we have used the identification (3.6) in the domain. Since each $\Psi_{\epsilon, t}$ is a quotient map, it follows from general facts about colimits in the category of topological spaces that $\text{colim}_{\epsilon, t \rightarrow 0}(\Psi_{\epsilon, t})$ is also a quotient map. The map

$$\Psi: \text{Diff}_{\text{dec}}(L\sharp M) \longrightarrow \text{Diff}_{\text{dec}}(L\sharp M) / \text{Diff}_{\text{dec}}(L),$$

i.e., the map (3.5) that we would like to show is a Serre fibration, is also a quotient map, with the same domain. Since M_t is a cofinal family of neighbourhoods of M in $L\sharp M$, two diffeomorphisms of $\text{Diff}_{\text{dec}}(L\sharp M)$ have the same image under Ψ if and only if they have the same image under $\text{colim}_{\epsilon, t \rightarrow 0}(\Psi_{\epsilon, t})$. As they are quotient maps of the same space, it follows that $\Psi \cong \text{colim}_{\epsilon, t \rightarrow 0}(\Psi_{\epsilon, t})$.

We will prove below that each $\Psi_{\epsilon, t}$ is a fibre bundle (and hence a Serre fibration), and then deduce that Ψ is a Serre fibration using the following general fact.

- (*) Any filtered colimit of based Serre fibrations between compactly-generated weak-Hausdorff spaces is again a Serre fibration.

For a reference for this fact, see Proposition 1.2.3.5(1) of [TV08], which states that a filtered colimit of fibrations is a fibration in any compactly generated model category. The classical model category of based compactly-generated weak-Hausdorff spaces, with its Quillen model structure in which the fibrations are the Serre fibrations, is compactly generated; see for example Proposition 6.3 of [MMSS01].

To apply (*) in our situation, first note that we are taking a directed colimit, which is in particular a filtered colimit. We then need to check that the diffeomorphism groups $\text{Diff}_{\text{dec},\epsilon}(L\downarrow M)$ and their quotients are compactly-generated weak-Hausdorff spaces. Diffeomorphism groups of manifolds, in the Whitney topology, are always first-countable and Hausdorff, and thus compactly-generated and weak-Hausdorff. Moreover, the property of being compactly-generated is preserved when taking quotients. The property of being weak Hausdorff is *not* preserved when taking quotients; however, in the process of proving that each $\Psi_{\epsilon,t}$ is a fibre bundle below, we will also show that its target space $\text{Diff}_{\text{dec},\epsilon}(L\downarrow M)/\text{Diff}_{\text{dec},\epsilon,t}(L)$ is Hausdorff.

It therefore remains to show that each $\Psi_{\epsilon,t}$ is a fibre bundle (and its target space is Hausdorff). Write

$$\text{Emb}_{\text{dec},\epsilon}(M_t, L\downarrow M)_L$$

for the space of smooth, proper embeddings $\varphi: M_t \rightarrow L\downarrow M$ such that $\varphi(B) \subseteq A \sqcup B$, the restriction of φ to $e'_2(\mathbb{B}_\epsilon^d)$ is the identity and there exists $\bar{\varphi} \in \text{Diff}_{\text{dec},\epsilon}(L\downarrow M)$ such that $\varphi = \bar{\varphi} \circ \iota$, where ι is the inclusion of M_t into $L\downarrow M$. Then

$$\text{Emb}_{\text{dec}}(M, L\downarrow M)_L \cong \text{colim}_{\epsilon,t \rightarrow 0} (\text{Emb}_{\text{dec},\epsilon}(M_t, L\downarrow M)_L), \quad (3.7)$$

see Definitions 3.22 and 3.23. There is a restriction map

$$\Phi_{\epsilon,t}: \text{Diff}_{\text{dec},\epsilon}(L\downarrow M) \longrightarrow \text{Emb}_{\text{dec},\epsilon}(M_t, L\downarrow M)_L,$$

which is equivariant with respect to the left-action of $\text{Diff}_{\text{dec},\epsilon}(L\downarrow M)$ by post-composition. This factors through the quotient map $\Psi_{\epsilon,t}$, so we have an induced map

$$\begin{array}{ccc} \text{Diff}_{\text{dec},\epsilon}(L\downarrow M) & \xrightarrow{\Phi_{\epsilon,t}} & \text{Emb}_{\text{dec},\epsilon}(M_t, L\downarrow M)_L \\ \Psi_{\epsilon,t} \downarrow & & \nearrow \widehat{\Phi}_{\epsilon,t} \\ \text{Diff}_{\text{dec},\epsilon}(L\downarrow M)/\text{Diff}_{\text{dec},\epsilon,t}(L) & & \end{array}$$

By definition of the right-hand embedding space, the map $\Phi_{\epsilon,t}$ is surjective, and hence so is the induced map $\widehat{\Phi}_{\epsilon,t}$. Moreover, if two diffeomorphisms of $\text{Diff}_{\text{dec},\epsilon}(L\downarrow M)$ have the same image under $\Phi_{\epsilon,t}$, their difference lies in $\text{Diff}_{\text{dec},\epsilon,t}(L)$, so the induced map $\widehat{\Phi}_{\epsilon,t}$ is also injective. We will prove in the next paragraphs that:

- (**) The map $\Phi_{\epsilon,t}$ is a fibre bundle.

In particular, it is a quotient map, since surjective fibre bundles are always quotient maps. Thus the induced map $\widehat{\Phi}_{\epsilon,t}$ must be a homeomorphism. This implies:

- The map $\Psi_{\epsilon,t}$ is also a fibre bundle, hence a Serre fibration.
- Its target space is homeomorphic to the embedding space $\text{Emb}_{\text{dec},\epsilon}(M_t, L\downarrow M)_L$, which we have given the Whitney topology, so it is Hausdorff.
- We also obtain the second statement of the proposition:

$$\begin{aligned} \text{Diff}_{\text{dec}}(L\downarrow M)/\text{Diff}_{\text{dec}}(L) &\cong \text{colim}_{\epsilon,t \rightarrow 0} (\text{Diff}_{\text{dec},\epsilon}(L\downarrow M)/\text{Diff}_{\text{dec},\epsilon,t}(L)) \\ &\cong \text{colim}_{\epsilon,t \rightarrow 0} (\text{Emb}_{\text{dec},\epsilon}(M_t, L\downarrow M)_L) \\ &\cong \text{Emb}_{\text{dec}}(M, L\downarrow M)_L. \end{aligned}$$

Here we combine the identification $\Psi \cong \operatorname{colim}_{\epsilon, t \rightarrow 0}(\Psi_{\epsilon, t})$ with the colimit of $\widehat{\Phi}_{\epsilon, t}$ and (3.7).

It therefore remains just to prove statement (**), that $\Phi_{\epsilon, t}$ is a fibre bundle. Since it is equivariant with respect to the left-action of $\operatorname{Diff}_{\operatorname{dec}, \epsilon}(L\sharp M)$, it suffices to prove that the action of $\operatorname{Diff}_{\operatorname{dec}, \epsilon}(L\sharp M)$ on $\operatorname{Emb}_{\operatorname{dec}, \epsilon}(M_t, L\sharp M)_L$ is *locally retractile* (\equiv *admits local cross-sections*). This is because, by [Pal60b, Theorem A], any G -equivariant map into a G -locally retractile space is a fibre bundle.

Thus, we have to prove the following statement: given an embedding $e \in \operatorname{Emb}_{\operatorname{dec}, \epsilon}(M_t, L\sharp M)_L$, we may find an open neighbourhood \mathcal{U} of e and a continuous map $\gamma: \mathcal{U} \rightarrow \operatorname{Diff}_{\operatorname{dec}, \epsilon}(L\sharp M)$ such that $\gamma(e) = \operatorname{id}$ and $\gamma(f) \circ e = f$ for any $f \in \mathcal{U}$. Note that, since $\operatorname{Diff}_{\operatorname{dec}, \epsilon}(L\sharp M)$ acts transitively on $\operatorname{Emb}_{\operatorname{dec}, \epsilon}(M_t, L\sharp M)_L$ (because $\Phi_{\epsilon, t}$ is both equivariant and surjective), it suffices to prove this for just one such e , which we take to be the inclusion $M_t \hookrightarrow L\sharp M$.

To prove this, we apply a result of Cerf [Cer61, Lemme II.2.1.2, page 291], which we first recall. Let X be a manifold-with-corners. This means in particular that X has a stratification into *faces* (for example, if X is a connected manifold with boundary, but no higher-codimension corners, then its set of faces is $\pi_0(\partial X) \sqcup \{X\}$). Each point $x \in X$ may lie in many faces, but it has a unique *smallest* face (according to inclusion) in which it lies, which we denote by $F_X(x)$. Now if Y is any submanifold-with-corners of X , we define

$$C_{\operatorname{face}}^{\infty}(Y, X) = \{\text{smooth maps } \varphi: Y \rightarrow X \text{ such that } F_X(\varphi(x)) = F_X(x) \text{ for each } x \in Y\},$$

equipped with the Whitney topology. The *Extension Lemma* II.2.1.2 of [Cer61] says that, if Y is closed in X and V is any neighbourhood of Y in X , then the restriction map

$$C_{\operatorname{face}}^{\infty}(X, X) \longrightarrow C_{\operatorname{face}}^{\infty}(Y, X)$$

admits a section s defined on an open neighbourhood \mathcal{V} of the inclusion in $C_{\operatorname{face}}^{\infty}(Y, X)$, such that $s(\operatorname{incl}) = \operatorname{id}$ and $s(f)(x) = x$ for all $f \in \mathcal{V}$ and $x \in X \setminus V$.

Step 1. Let us write $\partial_{\bullet}L$ for the union of all boundary components of L except for the one that intersects the image of e_2 ; see Figure 3.3. Note that $\partial_{\bullet}L$ may or may not intersect the image of e_1 . Then there is a canonical identification:

$$\pi_0(\partial(L\sharp M)) \cong \pi_0(\partial_{\bullet}L) \sqcup \pi_0(\partial M). \quad (3.8)$$

This is necessarily asymmetric in L and M . Each embedding $f \in \operatorname{Emb}_{\operatorname{dec}, \epsilon}(M_t, L\sharp M)_L$ extends to a diffeomorphism of $L\sharp M$, so it induces an injection $f_{\partial}: \pi_0(\partial M) \rightarrow \pi_0(\partial(L\sharp M))$. In particular, if f is the inclusion, then f_{∂} is also the inclusion, under the identification (3.8). In addition, we know that f sends B into $A \sqcup B$, so it also induces a map $f_{\sharp}: \pi_0(B) \rightarrow \pi_0(A) \sqcup \pi_0(B)$, which must be an injection since A and B are closed manifolds and f is an embedding. The function $f \mapsto (f_{\partial}, f_{\sharp})$ is locally constant, so its fibres are open. Let \mathcal{U}' be the open subset of $\operatorname{Emb}_{\operatorname{dec}, \epsilon}(M_t, L\sharp M)_L$ consisting of all f such that f_{∂} is the inclusion and $f_{\sharp}(\pi_0(B)) = \pi_0(B)$. Note that the second condition implies that $f(B) = B$, since f is an embedding and B is a closed manifold.

Step 2. Write $M_{\epsilon, t} = e_1(\mathbb{B}_{\epsilon}^d) \sqcup M_t$ (pictured in Figure 3.4). Let

$$\gamma': \operatorname{Emb}_{\operatorname{dec}, \epsilon}(M_t, L\sharp M)_L \longrightarrow C^{\infty}(M_{\epsilon, t}, L\sharp M)$$

be the continuous map that extends a given embedding $M_t \hookrightarrow L\sharp M$ to a smooth map $M_{\epsilon, t} \rightarrow L\sharp M$ by defining it to be the identity on $e_1(\mathbb{B}_{\epsilon}^d)$. Note that this may fail to be injective, so it is just a smooth map, not necessarily an embedding. Also observe that, if f lies in the open subset \mathcal{U}' from Step 1, then $\gamma'(f)$ lies in the subspace $C_{\operatorname{face}}^{\infty}(M_{\epsilon, t}, L\sharp M)$, since it takes points of $\operatorname{int}(L\sharp M) \cap M_{\epsilon, t}$ into $\operatorname{int}(L\sharp M)$ and, for any boundary-component P of $L\sharp M$, it takes $P \cap M_{\epsilon, t}$ into P (this uses the fact that $f_{\partial} = \operatorname{id}$). Restricting γ' to \mathcal{U}' , we therefore have a continuous map

$$\gamma': \mathcal{U}' \longrightarrow C_{\operatorname{face}}^{\infty}(M_{\epsilon, t}, L\sharp M)$$

such that $\gamma'(\operatorname{incl}) = \operatorname{incl}$ and $\gamma'(f)|_{M_t} = f$ for all $f \in \mathcal{U}'$.

Step 3. Now set $X = L\sharp M$ and $Y = M_{\epsilon, t}$ in the Extension Lemma of Cerf above, and choose V to be any open neighbourhood of $M_{\epsilon, t}$ in $L\sharp M$ that is disjoint from the submanifold $A \subset \operatorname{int}(L)$.




Figure 3.4 The submanifold $M_{\epsilon,t}$ (shaded in green) of $L\mathfrak{h}M$ from the proof of Proposition 3.24.

Composing the local section s obtained from the Extension Lemma with γ' , we have a continuous map

$$\gamma'' = s \circ \gamma' : \mathcal{U}'' = (\gamma')^{-1}(\mathcal{V}) \longrightarrow C_{\text{face}}^\infty(L\mathfrak{h}M, L\mathfrak{h}M)$$

such that $\gamma''(\text{incl}) = \text{id}$ and for any $f \in \mathcal{U}''$ we have $\gamma''(f)|_{M_t} = f$ and $\gamma''(f)(A) = A$. Moreover, by construction, we also know that $\gamma''(f)(B) = B$ and $\gamma''(f)(x) = x$ for all $x \in e_1(\mathbb{B}_\epsilon^d) \sqcup e'_2(\mathbb{B}_\epsilon^d)$.

Step 4. Finally, note that $\text{Diff}(L\mathfrak{h}M)$ is open in $C^\infty(L\mathfrak{h}M, L\mathfrak{h}M)$, so

$$\mathcal{U} = (\gamma'')^{-1}(C_{\text{face}}^\infty(L\mathfrak{h}M, L\mathfrak{h}M) \cap \text{Diff}(L\mathfrak{h}M))$$

is an open neighbourhood of the inclusion in $\text{Emb}_{\text{dec},\epsilon}(M_t, L\mathfrak{h}M)_L$. For each $f \in \mathcal{U}$, the diffeomorphism $\gamma''(f)$ of $L\mathfrak{h}M$ fixes each point of $e_1(\mathbb{B}_\epsilon^d) \sqcup e'_2(\mathbb{B}_\epsilon^d)$ and sends $A \sqcup B$ onto itself, so it is an element of $\text{Diff}_{\text{dec},\epsilon}(L\mathfrak{h}M)$. So we have a continuous map

$$\gamma = \gamma''|_{\mathcal{U}} : \mathcal{U} \longrightarrow \text{Diff}_{\text{dec},\epsilon}(L\mathfrak{h}M)$$

such that $\gamma(\text{incl}) = \text{id}$ and, for all $f \in \mathcal{U}$, we have $\gamma(f) \circ \text{incl} = \gamma(f)|_{M_t} = f$.

Summary. The 4-step construction above may be summarised in the following diagram:

$$\begin{array}{ccccc}
\text{Emb}_{\text{dec},\epsilon}(M_t, L\mathfrak{h}M) & \xrightarrow{\gamma'} & C^\infty(M_{\epsilon,t}, L\mathfrak{h}M) & & \\
\cup & & \cup & & \\
\mathcal{U}' & \longrightarrow & C_{\text{face}}^\infty(M_{\epsilon,t}, L\mathfrak{h}M) & \longleftarrow & C_{\text{face}}^\infty(L\mathfrak{h}M, L\mathfrak{h}M) \\
\cup & & \cup & & \uparrow \\
\mathcal{U}'' & \longrightarrow & \mathcal{V} & \xrightarrow{s} & \\
\cup & & & & \downarrow \\
\mathcal{U} & \xrightarrow{\gamma} & C_{\text{face}}^\infty(L\mathfrak{h}M, L\mathfrak{h}M) \cap \text{Diff}(L\mathfrak{h}M) & \subseteq & \text{Diff}(L\mathfrak{h}M),
\end{array}$$

where the construction of γ ensures that its image lies in $\text{Diff}_{\text{dec},\epsilon}(L\mathfrak{h}M) \subseteq \text{Diff}(L\mathfrak{h}M)$. \square

Remark 3.26 We note that all of the above may be adapted to the setting where the closed submanifolds $A \subset \text{int}(L)$ and $B \subset \text{int}(M)$ are equipped with orientations, and all morphisms of decorated manifolds are required to preserve these orientations. Proposition 3.24 generalises immediately to this setting.

3.4. Semi-monoidal categories and semicategories

All of the examples of categories \mathcal{C}_\circ for which we would like topologically to construct representations will be of the form $\langle \mathcal{G}_\circ, \mathcal{M}_\circ \rangle$, where \mathcal{G}_\circ is a braided monoidal groupoid and \mathcal{M}_\circ is a groupoid with a left-action of \mathcal{G}_\circ . We would therefore like to find a topological monoidal groupoid \mathcal{G} and a topological groupoid \mathcal{M} with a left-action of \mathcal{G} , satisfying condition (3.1) of Lemma 3.8 and such that $\pi_0(\mathcal{G}) \cong \mathcal{G}_\circ$ and $\pi_0(\mathcal{M}) \cong \mathcal{M}_\circ$. Given this, a continuous functor $\mathcal{C} := \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \bullet\text{Cov}\bullet$ will induce a functor

$$\mathcal{C}_\circ = \langle \mathcal{G}_\circ, \mathcal{M}_\circ \rangle \cong \pi_0(\langle \mathcal{G}, \mathcal{M} \rangle) = \pi_0(\mathcal{C}) \longrightarrow \bullet\text{Mod}\bullet,$$

via the construction summarised in the diagram (2.10). Note that there is no need for \mathcal{G} to be braided, since this structure is not needed in order to form the topological Quillen bracket

construction, or for Lemma 3.8. In fact, it will be convenient for our examples to drop even more structure from \mathcal{G} , and assume only that it is a *semi-monoidal category*. We recall that this is defined analogously to a monoidal category, but without any of the structure or conditions involving left or right units. This is because we will be able to lift the monoidal structure of \mathcal{G}_\circ to an associative binary operation on \mathcal{G} (more precisely, a binary operation admitting an associator that satisfies the pentagon condition), but we will not be able to ensure that this lifted operation is *unital*, without losing condition (3.1) of Lemma 3.8; cf. Remark 3.18.

Definition 3.27 If \mathcal{G} is a topological semi-monoidal groupoid and \mathcal{M} is a topological category with a continuous left-action of \mathcal{G} , then Definition 3.1 generalises directly to this setting, and produces a semicategory $\langle \mathcal{G}, \mathcal{M} \rangle$. The associator of \mathcal{G} is used to define composition in $\langle \mathcal{G}, \mathcal{M} \rangle$ and the pentagon condition for the associator implies associativity of this composition.

Lemma 3.28 *Let \mathcal{G} be a topological semi-monoidal groupoid and \mathcal{M} is a topological category with a continuous left-action of \mathcal{G} , satisfying the condition of Lemma 3.8. Then there is a canonical isomorphism of semicategories*

$$\pi_0(\langle \mathcal{G}, \mathcal{M} \rangle) \cong \langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle.$$

Using this definition and lemma it will therefore suffice, in our examples, to find a topological *semi-monoidal groupoid* \mathcal{G} , such that $\pi_0(\mathcal{G}) \cong \mathcal{G}_\circ$ as semi-monoidal groupoids and which satisfies condition (3.1) of Lemma 3.8. Then $\langle \mathcal{G}, \mathcal{M} \rangle$ is a topological *semicategory*, and we will construct, geometrically in §5.1, continuous semifunctors $\langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \bullet\text{Cov}\bullet$. Via Lemma 3.28 and the construction of §2.6 summarised in diagram (2.10), we then obtain a *semifunctor*

$$\langle \mathcal{G}_\circ, \mathcal{M}_\circ \rangle \longrightarrow \bullet\text{Mod}\bullet.$$

The source and target of this semifunctor are both categories (since \mathcal{G}_\circ is a monoidal groupoid, not just a semi-monoidal groupoid), and so one may ask whether this semifunctor is in fact a functor, and in all of our examples it will be trivial to verify that it does in fact preserve identities, and is therefore a functor.

Caveat 3.29 In practice, in the remainder of this paper, we will omit mention of this subtlety about a lack of units and identities at the topological level, to avoid unnecessary extra complications. Formally, however, one should modify the construction as described above.

4. Topological categories for families of groups

In this section, we discuss in more detail the topological groupoids \mathcal{D}_d of “decorated manifolds” introduced in the previous section. In §4.1, we discuss their properties with respect to braidings and symmetries. In §4.2, we give an explicit description of the morphism spaces of “Quillen bracket categories of manifolds” in terms of embedding spaces. In §4.3, we establish several split homotopy fibration sequences, and hence split short exact sequences on π_1 and π_0 of embedding spaces and diffeomorphism groups, and use these to identify motion groups with “braided mapping class groups”. In sections 4.4–4.6, we then describe, and recall the key properties of, the relevant subgroupoids of \mathcal{D}_2 and \mathcal{D}_3 that we will be especially interested in.

4.1. Braidings and symmetries

If we restrict attention to those decorated manifolds whose two boundary-cylinder germs lie on the same boundary-component, which is a sphere, then there is a natural braiding at the level of π_0 , which is symmetric if $d \geq 3$.

Definition 4.1 We say that a decorated manifold (M, A, e_1, e_2) has *spherical preferred boundary* if the embedded $(d-1)$ -discs $e_1(b\mathbb{B}_1^d)$ and $e_2(b\mathbb{B}_1^d)$ lie on the same boundary-component $\partial_0 M$ of M , and moreover $\partial_0 M \cong S^{d-1}$. Write $\mathcal{D}_d^{\text{sph}}$ for the full subgroupoid of \mathcal{D}_d on decorated manifolds with spherical preferred boundary. Define $\mathcal{D}_d^{+, \text{sph}} \subseteq \mathcal{D}_d^+$ similarly.

Definition 4.2 An inclusion of topologically-enriched categories $\mathcal{C} \subseteq \mathcal{D}$ is called *0-full* if, for each pair of objects c, c' of \mathcal{C} , the subspace $\mathcal{C}(c, c') \subseteq \mathcal{D}(c, c')$ is a union of path-components. Note that a 0-full inclusion $\mathcal{C} \subseteq \mathcal{D}$ induces an inclusion $\pi_0(\mathcal{C}) \subseteq \pi_0(\mathcal{D})$. In fact, subcategories of $\pi_0(\mathcal{D})$ correspond bijectively to 0-full subcategories of \mathcal{D} .

Lemma 4.3 *The subgroupoid $\pi_0(\mathcal{D}_d^{\text{sph}})$ of $\pi_0(\mathcal{D}_d)$ is closed under \natural , so it is a monoidal groupoid. Moreover, $\pi_0(\mathcal{D}_d^{\text{sph}})$ is braided if $d = 2$ and symmetric if $d \geq 3$.*

It follows that, if $\mathcal{G} \subseteq \mathcal{D}_d^{\text{sph}}$ is a 0-full subgroupoid that is closed under \natural and contains the braiding morphisms – described in the proof below – then $\pi_0(\mathcal{G})$ is braided monoidal, and symmetric monoidal if $d \geq 3$. The same statements hold when $\mathcal{D}_d^{\text{sph}}$ is replaced by $\mathcal{D}_d^{\text{sph},+}$.

Proof. The first statement is clear, since the connected sum of two spheres is again a sphere. The construction of the braiding morphisms is exactly analogous to Figure 2 on page 609 of [RW17], with the minor difference that they use two intervals, embedded in a circle boundary-component, intersecting at a point, whereas we use two disjoint discs in a spherical boundary-component. If the dimension d is at least 3, one may use the extra dimension to isotope the square of this braiding morphism to the identity, so it is a symmetry for $\pi_0(\mathcal{D}_d^{\text{sph}})$. \square

4.2. Quillen bracket categories of manifolds

Fix $d \geq 2$. Let $\mathcal{G} \subseteq \mathcal{D}_d$ be a full subgroupoid that is closed under \natural and let $\mathcal{M} \subseteq \mathcal{D}_d$ be a 0-full subgroupoid that is closed under the action of \mathcal{G} through \natural . Alternatively, we allow \mathcal{D}_d^+ in place of \mathcal{D}_d . This implies that \mathcal{G} is a semi-monoidal groupoid with an action on \mathcal{M} , and we may form the Quillen bracket construction $\langle \mathcal{G}, \mathcal{M} \rangle$, which is a topological semicategory; cf. §3.1 and §3.4. It is called a *Quillen bracket category of manifolds*.

Lemma 4.4 *The Serre fibration condition (3.1) of Lemma 3.8 is satisfied for this \mathcal{G} and \mathcal{M} , and hence we have*

$$\pi_0(\langle \mathcal{G}, \mathcal{M} \rangle) \cong \langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle.$$

Proof. If \mathcal{M} is a full subgroupoid of \mathcal{D}_d , this follows directly from Proposition 3.24 (and Remark 3.26 for \mathcal{D}_d^+) together with Lemma 3.8. If \mathcal{M} only satisfies the weaker property of being a 0-full subgroupoid of \mathcal{D}_d , it follows from these results together with Lemma 4.5 below. \square

Lemma 4.5 *Let X be a space with a continuous right-action of a topological group G such that the projection $X \rightarrow X/G$ is a Serre fibration. Let $X_0 \subseteq X$ be a union of path-components such that the G -action sends X_0 into itself. Then the projection $X_0 \rightarrow X_0/G$ is also a Serre fibration.*

Proof. More generally, by considering lifting diagrams, one may prove that, in the following square:

$$\begin{array}{ccc} A & \xrightarrow{a} & B \\ g \downarrow & & \downarrow f \\ C & \xrightarrow{b} & D \end{array}$$

if f is a Serre fibration, a is an inclusion of a union of path-components, and b is injective, then g is also a Serre fibration. In our setting, a is the inclusion $X_0 \hookrightarrow X$, which is assumed to be a union of path-components, and b is the induced map $X_0/G \rightarrow X/G$, which is injective. \square

Definition 4.6 (*Decorated diffeomorphisms, revisited.*) For a decorated manifold $M = (M, A)$, recall from Definition 3.21 that the topological group $\text{Diff}_{\text{dec}}(M)$ of decorated diffeomorphisms of M is simply the automorphism group of M in the topologically-enriched groupoid \mathcal{D}_d . If M lies in a subgroupoid $\mathcal{H} \subseteq \mathcal{D}_d$, write $\text{Diff}_{\mathcal{H}}(M) \subseteq \text{Diff}_{\text{dec}}(M)$ for the subgroup of automorphisms of M in \mathcal{H} . Note that this is an equality if $\mathcal{H} \subseteq \mathcal{D}_d$ is full.

Definition 4.7 (*Decorated embeddings, revisited.*) For decorated manifolds $M = (M, A)$ and $N = (N, B)$, recall from Definition 3.22 the space $\text{Emb}_{\text{dec}}(M, N)$ of decorated embeddings. If M and N lie in $\mathcal{M} \subseteq \mathcal{D}_d$, define $\text{Emb}_{\langle \mathcal{G}, \mathcal{M} \rangle}(M, N) \subseteq \text{Emb}_{\text{dec}}(M, N)$ to be the subspace where, in the third condition of Definition 3.22, the decorated manifold M' lies in \mathcal{G} and the decorated diffeomorphism $\bar{\varphi}$ lies in \mathcal{M} . Note that this second condition is automatic if $\mathcal{M} \subseteq \mathcal{D}_d$ is full. For an object L of \mathcal{G} , we also write $\text{Emb}_{\langle \mathcal{G}, \mathcal{M} \rangle}(M, N)_L$ for the subspace of $\text{Emb}_{\langle \mathcal{G}, \mathcal{M} \rangle}(M, N)$ where we may take $M' = L$ in Definition 3.22 (and $\bar{\varphi}$ lies in \mathcal{M}). Similarly to (3.4), the space $\text{Emb}_{\langle \mathcal{G}, \mathcal{M} \rangle}(M, N)$ decomposes as a topological disjoint union:

$$\text{Emb}_{\langle \mathcal{G}, \mathcal{M} \rangle}(M, N) \cong \bigsqcup_L \text{Emb}_{\langle \mathcal{G}, \mathcal{M} \rangle}(M, N)_L, \quad (4.1)$$

where the disjoint union runs over representatives of isomorphism classes of objects L of \mathcal{G} .

Proposition 4.8 *For any Quillen bracket category of manifolds $\langle \mathcal{G}, \mathcal{M} \rangle$, its morphism spaces may be identified as follows:*

$$\langle \mathcal{G}, \mathcal{M} \rangle(M, N) \cong \text{Emb}_{\langle \mathcal{G}, \mathcal{M} \rangle}(M, N).$$

Remark 4.9 When $\mathcal{M} \subseteq \mathcal{D}_d$ is full, this says that morphisms in $\langle \mathcal{G}, \mathcal{M} \rangle$ are embeddings of decorated manifolds such that the complement of the image of the embedding is an object of \mathcal{G} . In particular the space of morphisms is non-empty if and only if there exists an object M' of \mathcal{G} such that $M' \natural M$ is diffeomorphic to N as decorated manifolds.

Proof of Proposition 4.8. The space $\langle \mathcal{G}, \mathcal{M} \rangle(M, N)$ is described in Definition 3.1. In the notation of the proof of Lemma 3.8 (setting $X = M$ and $Y = N$), this is the quotient space Φ / \sim_t . In that proof, it is shown that this splits as the topological disjoint union of certain spaces denoted $q(\text{Hom}_{\mathcal{M}}(Z \natural X, Y))$, as Z runs over representatives of isomorphism classes of objects of \mathcal{G} . It is also proved that this space is homeomorphic to the quotient space $\text{Hom}_{\mathcal{M}}(Z \natural X, Y) / \text{Aut}_{\mathcal{G}}(Z)$. We therefore have the following homeomorphism, where the disjoint union runs over representatives Z of isomorphism classes of objects of \mathcal{G} :

$$\langle \mathcal{G}, \mathcal{M} \rangle(M, N) \cong \bigsqcup_Z \text{Hom}_{\mathcal{M}}(Z \natural X, Y) / \text{Aut}_{\mathcal{G}}(Z).$$

Since \mathcal{M} is a groupoid, the space $\text{Hom}_{\mathcal{M}}(Z \natural X, Y) / \text{Aut}_{\mathcal{G}}(Z)$ is empty unless $Z \natural X$ is isomorphic to Y in \mathcal{M} , in which case we may rewrite it as $\text{Aut}_{\mathcal{M}}(Z \natural X) / \text{Aut}_{\mathcal{G}}(Z) = \text{Diff}_{\mathcal{M}}(L \natural M) / \text{Diff}_{\text{dec}}(L)$, using the notation of Definition 4.7 and setting $L = Z$ (recall that $\mathcal{G} \subseteq \mathcal{D}_d$ is full). The second part of Proposition 3.24 says that the restriction map induces a homeomorphism

$$\text{Diff}_{\text{dec}}(L \natural M) / \text{Diff}_{\text{dec}}(L) \cong \text{Emb}_{\text{dec}}(M, L \natural M)_L,$$

and one may easily see that this sends the subspace $\text{Diff}_{\mathcal{M}}(L \natural M) / \text{Diff}_{\text{dec}}(L)$ homeomorphically onto the subspace $\text{Emb}_{\langle \mathcal{G}, \mathcal{M} \rangle}(M, L \natural M)_L$. Putting this all together, we have a homeomorphism

$$\langle \mathcal{G}, \mathcal{M} \rangle(M, N) \cong \bigsqcup_L \text{Emb}_{\langle \mathcal{G}, \mathcal{M} \rangle}(M, L \natural M)_L, \quad (4.2)$$

where the disjoint union runs over representatives L of isomorphism classes of objects of \mathcal{G} such that $L \natural M$ is isomorphic to N in \mathcal{M} .

Finally, consider the topological decomposition (4.1), where the disjoint union is indexed by representatives L of isomorphism classes of all objects of \mathcal{G} . If $L \natural M \not\cong N$ in \mathcal{M} , the corresponding term is empty, whereas if $L \natural M \cong N$ in \mathcal{M} , we may rewrite the corresponding term by replacing N with $L \natural M$, to obtain:

$$\text{Emb}_{\langle \mathcal{G}, \mathcal{M} \rangle}(M, N) \cong \bigsqcup_L \text{Emb}_{\langle \mathcal{G}, \mathcal{M} \rangle}(M, L \natural M)_L, \quad (4.3)$$

where the disjoint union now runs over representatives L of isomorphism classes of objects of \mathcal{G} such that $L \natural M$ is isomorphic to N in \mathcal{M} . Combining (4.2) and (4.3), we obtain the desired result. \square

Remark 4.10 Suppose that *cancellation* holds for our chosen subgroupoids \mathcal{G} and \mathcal{M} — meaning that $L \natural M \cong L' \natural M$ in \mathcal{M} implies $L \cong L'$ in \mathcal{G} for objects L, L' of \mathcal{G} and M of \mathcal{M} . Then the disjoint union (4.2) is taken either over the empty set or a set of size one, so we have:

$$\langle \mathcal{G}, \mathcal{M} \rangle(M, N) \cong \begin{cases} \text{Emb}_{(\mathcal{G}, \mathcal{M})}(M, L \natural M)_L & \text{if } \exists L \in \mathcal{G} \text{ such that } L \natural M \cong N \text{ in } \mathcal{M}; \\ \emptyset & \text{otherwise.} \end{cases}$$

All of our examples of \mathcal{G} and \mathcal{M} in §4.4–§4.6 satisfy cancellation. This follows from the classification of compact surfaces in all of our examples in dimension 2. In the examples of §4.6, it follows from the fact that the groupoid of finite sets and bijections under disjoint union satisfies cancellation, since the objects of the groupoids in the examples of §4.6 are determined up to isomorphism by the number of copies of \mathbb{S}^1 in the submanifold $A \subset M$.

4.3. Split short exact sequences

We now establish several split homotopy fibration sequences whose associated split short exact sequences will be used in our construction. In particular, the split short exact sequences (4.6) and (4.9) below will be used in the two versions of our general construction of global homological representations in §5.1 and §5.3 respectively. In addition, the split short exact sequence (4.9) implies that any motion group is a *braided mapping class group* – see Proposition 4.23 – in particular, a normal subgroup of a mapping class group.

Fix a closed submanifold $Z \subset \mathbb{R}^d$ and an open subgroup $G \leq \text{Diff}(Z)$. Note that, since $\text{Diff}(Z)$ is locally path-connected, this corresponds to a choice of subgroup of $\pi_0(\text{Diff}(Z))$.

4.3.1. The first short exact sequence.

Definition 4.11 For smooth manifolds X and Y , let us write $\mathcal{E}(X, Y) = \text{Emb}(X, Y)/\text{Diff}(X)$. For a subgroup $G \leq \text{Diff}(X)$, we also write $\mathcal{E}_G(X, Y) = \text{Emb}(X, Y)/G$.

Lemma 4.12 *For any decorated manifold $(M, A) \in \mathcal{D}_d$ there is a homotopy fibration sequence*

$$\mathcal{E}_G(Z, \overset{\circ}{M} \setminus A) \longrightarrow \mathcal{E}_{\text{Diff}(A) \times G}(A \sqcup Z, \overset{\circ}{M}) \overset{\leftarrow \text{---}}{\longrightarrow} \mathcal{E}(A, \overset{\circ}{M}), \quad (4.4)$$

in which the second map admits a section up to homotopy, as pictured.

Proof. The map $\mathcal{E}_{\text{Diff}(A) \times G}(A \sqcup Z, \overset{\circ}{M}) \rightarrow \mathcal{E}(A, \overset{\circ}{M})$ that forgets the embedding (modulo G) of Z is equivariant with respect to the left action of the topological group $\text{Diff}_c(\overset{\circ}{M})$ of compactly-supported diffeomorphisms of $\overset{\circ}{M}$. By [Pal18a, Proposition 4.15], the action of $\text{Diff}_c(\overset{\circ}{M})$ on $\mathcal{E}(A, \overset{\circ}{M})$ is locally retractile, i.e., it admits local sections. Thus, by [Pal60b, Theorem A], the map

$$\mathcal{E}_{\text{Diff}(A) \times G}(A \sqcup Z, \overset{\circ}{M}) \longrightarrow \mathcal{E}(A, \overset{\circ}{M}) \quad (4.5)$$

is a fibre bundle, in particular a Serre fibration. Write incl. for the inclusion of A into $\overset{\circ}{M}$ and $[\text{incl.}]$ for its $\text{Diff}(A)$ -orbit; this is a natural basepoint for $\mathcal{E}(A, \overset{\circ}{M})$. The point-set fibre of (4.5) over $[\text{incl.}] \in \mathcal{E}(A, \overset{\circ}{M})$ is clearly equal to $\mathcal{E}_G(Z, \overset{\circ}{M} \setminus A)$, so (4.4) is a fibration sequence.

We construct a section-up-to-homotopy as follows. Let $[\varphi] \in \mathcal{E}(A, \overset{\circ}{M})$. This determines an embedding $A \sqcup Z \hookrightarrow \overset{\circ}{M} \sqcup \mathbb{R}^d$ modulo the action of $\text{Diff}(A)$, since Z is given as a submanifold of \mathbb{R}^d . We recall from Definition 3.11 that a decorated manifold comes equipped with two *germs* of boundary-cylinders; we choose one of these and then choose a boundary-cylinder representing the given germ. Using this boundary-cylinders of M , we construct an embedding $\overset{\circ}{M} \sqcup \mathbb{R}^d \hookrightarrow \overset{\circ}{M}$ whose restriction to $\overset{\circ}{M}$ is isotopic to the identity. Composing this with the embedding above gives us an embedding $\varphi' : A \sqcup Z \hookrightarrow \overset{\circ}{M}$ modulo the action of $\text{Diff}(A)$. The desired splitting is then given by $[\varphi] \mapsto [\varphi']$. This is a section up to homotopy because, forgetting Z , φ' is isotopic to φ . \square

Corollary 4.13 *For any decorated manifold $(M, A) \in \mathcal{D}_d$ there is a split short exact sequence*

$$1 \longrightarrow \pi_1(\mathcal{E}_G(Z, \mathring{M} \setminus A)) \longrightarrow \pi_1(\mathcal{E}_{\text{Diff}(A) \times G}(A \sqcup Z, \mathring{M})) \xrightarrow{\leftarrow \text{---}} \pi_1(\mathcal{E}(A, \mathring{M})) \longrightarrow 1, \quad (4.6)$$

where the basepoint of each space of embeddings modulo diffeomorphisms is given by the inclusion.

We may, alternatively, consider spaces of embeddings whose image is allowed to lie either in the interior of M or in its boundary (but it must lie wholly in one or the other).

Definition 4.14 For $(M, A) \in \mathcal{D}_d$ and $Z \subset \mathbb{R}^d$ and $G \leq \text{Diff}(Z)$ as above, we write $\text{Emb}'(Z, \mathring{M} \setminus A)$ for the subspace of smooth maps $Z \rightarrow \mathring{M} \setminus A$ that are either smooth embeddings of Z into $\mathring{M} \setminus A$ or smooth embeddings of Z into ∂M . Write $\mathcal{E}'_G(Z, \mathring{M} \setminus A) = \text{Emb}'(Z, \mathring{M} \setminus A)/G$. Similarly, write $\text{Emb}'(A \sqcup Z, M)$ for the subspace of smooth maps $A \sqcup Z \rightarrow M$ that are either embeddings of $A \sqcup Z$ into M or disjoint unions of an embedding of A into \mathring{M} together with an embedding of Z into ∂M . Write $\mathcal{E}'_{\text{Diff}(A) \times G}(A \sqcup Z, M) = \text{Emb}'(A \sqcup Z, M)/(\text{Diff}(A) \times G)$.

Lemma 4.15 *The natural inclusions*

$$\mathcal{E}_G(Z, \mathring{M} \setminus A) \hookrightarrow \mathcal{E}'_G(Z, \mathring{M} \setminus A) \quad \text{and} \quad \mathcal{E}_{\text{Diff}(A) \times G}(A \sqcup Z, \mathring{M}) \hookrightarrow \mathcal{E}'_{\text{Diff}(A) \times G}(A \sqcup Z, M)$$

are homotopy equivalences. Hence we may rewrite the split short exact sequence (4.6) above as

$$1 \longrightarrow \pi_1(\mathcal{E}'_G(Z, \mathring{M} \setminus A)) \longrightarrow \pi_1(\mathcal{E}'_{\text{Diff}(A) \times G}(A \sqcup Z, M)) \xrightarrow{\leftarrow \text{---}} \pi_1(\mathcal{E}(A, \mathring{M})) \longrightarrow 1. \quad (4.7)$$

Proof. In each case, a deformation retraction may be defined by post-composing embeddings with an isotopy of self-embeddings of M starting from the identity and “shrinking” a collar neighbourhood of its boundary. \square

4.3.2. The second short exact sequence.

We now construct the second key split short exact sequence (sequence (4.9) below). We first need some notation.

Definition 4.16 For a decorated manifold $(M, A) = (M, A, e_1, e_2) \in \mathcal{D}_d$, recall that we write $\text{Diff}_{\text{dec}}(M, A)$ for its automorphism group in \mathcal{D}_d , in other words, the self-diffeomorphisms of M that send A onto itself and that are compatible with the boundary-cylinder germs e_1 and e_2 . If we have a decomposition $A = A_1 \sqcup A_2$, we now also write $\text{Diff}_{\text{dec}}(M, A_1, A_2)$ for the subgroup of $\text{Diff}_{\text{dec}}(M, A)$ of those diffeomorphisms that preserve this decomposition, in other words, that send A_1 onto itself and A_2 onto itself. Note that this is a finite covering of $\text{Diff}_{\text{dec}}(M, A)$.

Notation 4.17 We identify \mathbb{R}^d with the interior of the solid cylinder \mathbb{B}_1^d in a standard way, so the choice of closed submanifold $Z \subset \mathbb{R}^d$ determines a decorated manifold $(\mathbb{B}_1^d, Z) = (\mathbb{B}_1^d, Z, \text{id}, r)$, where r is the reflection of the solid cylinder $\mathbb{B}_1^d = \mathbb{D}^{d-1} \times [0, 1]$ in its second coordinate. For any other decorated manifold $(M, A) \in \mathcal{D}_d$, we may therefore consider the boundary connected sum $(M, A) \natural (\mathbb{B}_1^d, Z)$, which we denote by an abuse of notation by $(M, A \sqcup Z)$.

Lemma 4.18 *For any decorated manifold $(M, A) \in \mathcal{D}_d$ there is a homotopy fibration sequence*

$$\text{Diff}_{\text{dec}}(M, A, Z) \xrightarrow{\leftarrow \text{---}} \text{Diff}_{\text{dec}}(M, A) \longrightarrow \mathcal{E}(Z, \mathring{M} \setminus A), \quad (4.8)$$

in which the first map admits a section up to homotopy, as pictured.

Proof. The right-hand map above is equivariant with respect to the left action of $\text{Diff}_c(\mathring{M} \setminus A)$. By [Pal18a, Proposition 4.15], its action on $\mathcal{E}(Z, \mathring{M} \setminus A)$ is locally retractile, so by [Pal60b, Theorem A], the right-hand map above is a fibre bundle. Its point-set fibre over the basepoint [incl.] is clearly equal to $\text{Diff}_{\text{dec}}(M, A, Z)$. The section up to homotopy is constructed similarly to the proof of Lemma 4.12, using a self-embedding $M \hookrightarrow M$ that restricts to the identity on $A \subset M$, is isotopic to the identity and whose image is disjoint from Z (when considered as a submanifold of M via a chosen boundary-cylinder). \square

Corollary 4.19 *For any decorated manifold $(M, A) \in \mathcal{D}_d$ there is a split short exact sequence*

$$1 \longrightarrow \pi_1(\mathcal{E}(Z, \mathring{M} \setminus A)) \longrightarrow \pi_0(\mathrm{Diff}_{\mathrm{dec}}(M, A, Z)) \xrightarrow{\kappa \dashrightarrow} \pi_0(\mathrm{Diff}_{\mathrm{dec}}(M, A)) \longrightarrow 1, \quad (4.9)$$

where the basepoint of the space $\mathcal{E}(Z, \mathring{M} \setminus A)$ is given by the inclusion.

The split homotopy fibration sequence (4.8) may be generalised as follows.

Lemma 4.20 *For any decorated manifold $(M, A) \in \mathcal{D}_d$ and any open subgroup $H \leq \mathrm{Diff}_{\mathrm{dec}}(M, A)$, the split homotopy fibration sequence (4.8) restricts to a split homotopy fibration sequence*

$$\mathrm{Diff}_{\mathrm{dec}}(M, A, Z) \cap H \xrightarrow{\kappa \dashrightarrow} H \longrightarrow \mathcal{E}(Z, \mathring{M} \setminus A), \quad (4.10)$$

and hence a corresponding split short exact sequence.

Proof. Since H is an open subgroup of a topological group, it is also closed and thus a union of path-components. The restriction map $\mathrm{Diff}_{\mathrm{dec}}(M, A) \rightarrow \mathcal{E}(Z, \mathring{M} \setminus A)$ is a fibre bundle, hence a Serre fibration, by the proof of Lemma 4.18. In general, if $E \rightarrow B$ is a Serre fibration and $E_0 \subseteq E$ is a union of path-components, then the restriction $E_0 \rightarrow B$ is also a Serre fibration. This establishes the homotopy fibration sequence. To see that the section up to homotopy of (4.8) restricts to give a section up to homotopy of (4.10), one again just has to use the fact that H is a union of path-components of $\mathrm{Diff}_{\mathrm{dec}}(M, A)$. \square

4.3.3. Motion groups are braided mapping class groups.

Definition 4.21 The *braided diffeomorphism group* $\mathrm{Diff}_{\mathrm{dec}}^{\mathrm{br}}(M, A)$ of a decorated manifold (M, A) is the kernel of the natural homomorphism

$$\mathrm{Diff}_{\mathrm{dec}}(M, A) \longrightarrow \pi_0(\mathrm{Diff}_{\mathrm{dec}}(M, \emptyset)),$$

in other words, the subgroup of diffeomorphisms of (M, A) that become isotopic to the identity after forgetting A .

Definition 4.22 Given a closed manifold Z and an embedding $Z \hookrightarrow \mathring{M}$, the corresponding *motion group* $\mathrm{Mot}_Z(M)$ is the fundamental group $\pi_1(\mathcal{E}(Z, \mathring{M}))$.

Corollary 4.19 implies the following identification of motion groups with π_0 of braided diffeomorphism groups, in other words, *motion groups are braided mapping class groups*.

Proposition 4.23 *For any closed submanifold $Z \subset \mathbb{R}^d$, we have a canonical isomorphism*

$$\mathrm{Mot}_Z(M) = \pi_1(\mathcal{E}(Z, \mathring{M})) \cong \pi_0(\mathrm{Diff}_{\mathrm{dec}}^{\mathrm{br}}(M, Z)).$$

Proof. This follows immediately from Corollary 4.19 in the special case where $A = \emptyset$. \square

Remark 4.24 (*Globality*) In particular, Proposition 4.23 tells us that all motion groups are normal subgroups of the appropriate $\pi_0(\mathrm{Diff}(-))$'s of decorated manifolds. Also, more obviously, mapping class groups of d -manifolds are clearly examples of $\pi_0(\mathrm{Diff}(-))$'s of decorated manifolds, where we take A to be the empty submanifold. Thus *the discrete groupoid $\pi_0(\mathcal{D}_d)$ contains all motion groups and all mapping class groups of d -dimensional manifolds*.

The homological representations that we construct in §5.1 and §5.3 below are functors of the form $\mathfrak{U}\mathcal{D}_d \rightarrow \bullet\mathrm{Mod}\bullet$. Since the target category is discrete, they factor through $\pi_0(\mathfrak{U}\mathcal{D}_d) \cong \mathfrak{U}(\pi_0(\mathcal{D}_d))$, whose underlying groupoid is $\pi_0(\mathcal{D}_d)$. Each such homological representation therefore restricts to homological representations of all motion groups and mapping class groups in dimension d . This is why we refer to these as *global* homological representations.

4.3.4. Actions of (braided) mapping class groups.

The following lemma will be important for the proof of the “ Q -stability lemma” in §5.1.

Lemma 4.25 *There is a natural action of $\text{Diff}_{\text{dec}}(M, A)$ on $\mathcal{E}(Z, \mathring{M} \setminus A)$ given by post-composition, which induces an action of $\pi_0(\text{Diff}_{\text{dec}}(M, A))$ on $\pi_1(\mathcal{E}(Z, \mathring{M} \setminus A))$. This action agrees with the one coming from the splitting in the short exact sequence (4.9).*

Similarly, there is a natural action of $\text{Diff}_{\text{dec}}(M, A)$ on $\mathcal{E}_G(Z, \mathring{M} \setminus A)$ given by post-composition, which induces an action of

$$\pi_1(\mathcal{E}(A, \mathring{M})) = \pi_0(\text{Diff}_{\text{dec}}^{\text{br}}(M, A)) \leq \pi_0(\text{Diff}_{\text{dec}}(M, A)) \quad (4.11)$$

on $\pi_1(\mathcal{E}_G(Z, \mathring{M} \setminus A))$. This action agrees with the one coming from the splitting in the short exact sequence (4.6).

Proof. We first of all note that there is a slight generalisation of the split short exact sequence (4.9), as follows. Lemma 4.18 generalises (with the same proof) to the setting where, for an open subgroup $G \leq \text{Diff}(Z)$, the base space of the fibration is replaced with $\mathcal{E}_G(Z, \mathring{M} \setminus A)$ and the fibre is replaced with $\text{Diff}_{\text{dec}}(M, A, Z|G)$, the subgroup of $\varphi \in \text{Diff}_{\text{dec}}(M, A, Z)$ such that $\varphi|_Z \in G$. The corresponding split short exact sequence is the middle row of the following diagram.

$$\begin{array}{ccccccc}
 & & 1 & & 1 & & 1 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 1 & \longrightarrow & \pi_1(\mathcal{E}_G(Z, \mathring{M} \setminus A)) & \longrightarrow & \pi_1(\mathcal{E}_{\text{Diff}(A) \times G}(A \sqcup Z, \mathring{M})) & \longrightarrow & \pi_1(\mathcal{E}(A, \mathring{M})) \longrightarrow 1 \\
 & & \text{id} \downarrow & & \downarrow & & \downarrow \\
 1 & \longrightarrow & \pi_1(\mathcal{E}_G(Z, \mathring{M} \setminus A)) & \longrightarrow & \pi_0(\text{Diff}_{\text{dec}}(M, A, Z|G)) & \longrightarrow & \pi_0(\text{Diff}_{\text{dec}}(M, A)) \longrightarrow 1 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 1 & \longrightarrow & 1 & \longrightarrow & \pi_0(\text{Diff}_{\text{dec}}(M)) & \xrightarrow{\text{id}} & \pi_0(\text{Diff}_{\text{dec}}(M)) \longrightarrow 1 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 1 & & 1 & & 1
 \end{array} \quad (4.12)$$

Moreover, in this diagram:

- the top row is the split short exact sequence (4.6),
- the right-hand column is an instance of (4.9) (with $A = \emptyset$ and replacing $Z \mapsto A$),
- the middle column is an instance of the middle row (with $A = \emptyset$ and replacing $Z \mapsto A \sqcup Z$ and $G \mapsto \text{Diff}(A) \times G$)

In particular, the top-right vertical map is the inclusion (4.11). The top-right square of (4.12) also commutes when the horizontal arrows are replaced by the given splittings. Thus both parts of the lemma (for (4.6) and for (4.9)) will follow once we prove the analogue for the middle row of (4.12).

To see this, let us describe geometrically the action of $\varphi \in \pi_0(\text{Diff}_{\text{dec}}(M, A))$ on $\pi_1(\mathcal{E}_G(Z, \mathring{M} \setminus A))$ that arises from the split short exact sequence forming the middle row of (4.12). By definition, φ is an isotopy class of diffeomorphisms of M fixing (pointwise) a neighbourhood of its two boundary-cylinders and fixing A setwise. First choose an identification of M with $M \natural \mathbb{B}_1^d$ so that $A \subseteq M$ and $Z \subseteq \mathbb{B}_1^d$. The section sends φ to the diffeomorphism φ' of $M \natural \mathbb{B}_1^d$ that acts by φ on M and by the identity on \mathbb{B}_1^d . The diffeomorphism φ' then acts on $\pi_1(\mathcal{E}_G(Z, \mathring{M} \setminus A))$ by conjugation, where the latter group is viewed as a subgroup of $\pi_0(\text{Diff}_{\text{dec}}(M, A, Z|G))$ via the connecting homomorphism of the long exact sequence, which may be viewed geometrically as sending a loop of embeddings of Z to the corresponding *submanifold-pushing* diffeomorphism of M (named by analogy with the well-known *point-pushing* diffeomorphism when Z is zero-dimensional). But using the submanifold-pushing construction to view $\pi_1(\mathcal{E}_G(Z, \mathring{M} \setminus A))$ as a subgroup of $\pi_0(\text{Diff}_{\text{dec}}(M, A, Z|G))$ and then

acting by conjugation is equivalent to acting directly on $\pi_1(\mathcal{E}_G(Z, \overset{\circ}{M} \setminus A))$ through the obvious action of $\text{Diff}_{\text{dec}}(M, A, Z|G)$ on $\mathcal{E}_G(Z, \overset{\circ}{M} \setminus A)$. \square

4.4. Mapping class groups of surfaces

Define \mathcal{M}_2^{\natural} to be the full subgroupoid of $\mathcal{D}_2^{\text{spH}}$ on decorated surfaces (S, \emptyset, e_1, e_2) where S is a compact, connected, smooth surface with boundary. This collection of objects is clearly closed under the operation \natural , so it is a semi-monoidal subgroupoid, so the category \mathcal{M}_2^{\natural} inherits a topological semi-monoidal structure from \mathcal{D}_2 by Lemma 3.20. In the general notation of this section, this first example is $\mathcal{M} = \mathcal{G} = \mathcal{M}_2^{\natural}$.

We denote by \mathcal{M}_2 the groupoid $\pi_0(\mathcal{M}_2^{\natural})$. We recall that for diffeomorphisms of surfaces, the condition of fixing (a neighbourhood of) an interval in a boundary-component is equivalent to fixing two (neighbourhoods of) intervals in that boundary-component. Therefore \mathcal{M}_2 can be described as the groupoid of decorated surfaces (S, I) where S is a surface as above, equipped with a parametrised interval $I: [-1, 1] \hookrightarrow \partial_0 S$ in the boundary *decorating* one boundary component denoted by $\partial_0 S$. The *decorated surfaces groupoid* \mathcal{M}_2 is already introduced in [RW17, §5.6]. When there is no ambiguity, we omit the parametrised interval I from the notation.

The morphisms in \mathcal{M}_2 are the isotopy classes of diffeomorphisms of surfaces which restrict to the identity on a neighbourhood of their parametrised intervals I (or equivalently diffeomorphisms which fixes pointwise the boundary component $\partial_0 S$). The non-distinguished boundary components may be freely moved by the mapping classes. The automorphism group of S forms the *mapping class group* of S , which we denote by $\pi_0 \text{Diff}_I(S)$. When the surface S is orientable, the diffeomorphisms then automatically preserve that orientation as they restrict to the identity on a neighbourhood of I (the orientation on S being induced by that of I). By Lemma 4.3, the groupoid \mathcal{M}_2 has a braided monoidal structure induced by gluing, which is already considered in [RW17, §5.6.1]. Hence we may apply Quillen's bracket construction: by Lemma 4.4, we deduce that as semicategories $\pi_0(\mathcal{U}\mathcal{M}_2^{\natural}) \cong \mathcal{U}\mathcal{M}_2$.

Notation 4.26 We denote by \mathbb{D}^2 the unit 2-disc. Let $\Sigma_{0,1}^1$ denote the cylinder $\mathbb{S}^1 \times [0, 1]$ (which can be thought of as the disc \mathbb{D}^2 with a smaller disc in the interior which is removed), $\Sigma_{1,1}$ denote the torus with one boundary component ($\mathbb{S}^1 \times \mathbb{S}^1 \setminus \text{Int}(\mathbb{D}^2)$) and $\mathcal{N}_{1,1}$ denote a Möbius band. For S an object of the groupoid \mathcal{M}_2 , by the classification of surfaces, there exist $g, s, h \in \mathbb{N}$ such that there is a diffeomorphism $S \simeq (\natural_s \Sigma_{0,1}^1) \natural_g (\natural_h \Sigma_{1,1}) \natural_h (\natural_h \mathcal{N}_{1,1})$.

If $h = 0$, then g and s are unique, we denote by $\Sigma_{g,1}^s$ the boundary connected sum $(\Sigma_{0,1}^1)^{\natural_s} \natural_g \Sigma_{1,1}^{\natural_g}$ and by $\mathbf{\Gamma}_{g,1}^s$ the mapping class group $\pi_0 \text{Diff}_I(\Sigma_{g,1}^s)$. If $g = 0$, we denote by $\mathcal{N}_{h,1}^s$ the boundary connected sum $(\Sigma_{0,1}^1)^{\natural_s} \natural_h \mathcal{N}_{1,1}^{\natural_h}$ and by $\mathcal{N}_{h,1}^s$ the mapping class group $\pi_0 \text{Diff}_I(\mathcal{N}_{h,1}^s)$. In both cases, when $s = 0$, we omit it most of the time from the notation.

4.5. Surface braid groups

Let S be a compact, connected, smooth surface with a chosen boundary-component $\partial_0 S$. For each non-negative integer k , we denote by \underline{k} a closed submanifold of S consisting in k distinct points of the interior of S . Let $\mathcal{B}r^S$ be the subgroupoid of $\mathcal{D}_2^{\text{spH}}$ with objects all decorated surfaces $(S', \underline{n}, e_1, e_2)$ such that n is any non-negative integer and there is a diffeomorphism $S \cong S'$ taking $\partial_0 S$ onto $\partial_0 S'$; the morphisms of $\mathcal{B}r^S$ are given by the subgroups of the braided diffeomorphisms of Definition 4.21 and denoted by $\text{Diff}_{\text{dec}}^{\text{br}}(S, \underline{n})$. If $S = \mathbb{D}^2$, this collection of objects is closed under the operation \natural , so $\mathcal{B}r^{\mathbb{D}^2}$ is a monoidal subgroupoid of \mathcal{D}_2 by Lemma 3.20. Also, the groupoid $\mathcal{B}r^S$ is closed under the left-action of $\mathcal{B}r^{\mathbb{D}^2}$ via \natural . Hence, in the general notation of this section, we take $\mathcal{G} = \mathcal{B}r^{\mathbb{D}^2}$ and $\mathcal{M} = \mathcal{B}r^S$.

In general, $\mathcal{B}r^S$ is not a full subgroupoid of $\mathcal{D}_2^{\text{spH}}$, since there may be diffeomorphisms of S which restrict to the identity on the set \underline{n} that are not isotopic to the identity. However, the special case of $\mathcal{B}r^{\mathbb{D}^2}$ is a full subgroupoid, since *all* diffeomorphisms of \mathbb{D}^2 fixing a pair of disjoint intervals in the boundary are isotopic to the identity. In other words, the diffeomorphism group $\text{Diff}_{I \sqcup I}(\mathbb{D}^2)$ is path-connected, as we show in the following lemma.

Lemma 4.27 *Let $I \sqcup I$ be a pair of disjoint closed intervals in the boundary of the disc \mathbb{D}^2 and write $\text{Diff}_{I \sqcup I}(\mathbb{D}^2)$ for the group of diffeomorphisms of \mathbb{D}^2 that fix $I \sqcup I$ pointwise. Then $\text{Diff}_{I \sqcup I}(\mathbb{D}^2)$ is weakly contractible, in particular path-connected.*

Proof. By [Cer61, §II.2.2.2, Corollaire 2] (see also [Pal60b; Lim63]), the map

$$\text{Diff}_{I \sqcup I}(\mathbb{D}^2) \longrightarrow \text{Diff}_\partial(I \sqcup I) \cong (\text{Diff}_\partial(I))^2$$

that remembers just the action of a diffeomorphism restricted to the *complementary* pair of intervals $\partial\mathbb{D}^2 \setminus (I \sqcup I) \cong I \sqcup I$ is a fibre bundle. Its fibre over the identity is $\text{Diff}_\partial(\mathbb{D}^2)$, the group of diffeomorphisms of \mathbb{D}^2 fixing all of $\partial\mathbb{D}^2$ pointwise. The diffeomorphism group $\text{Diff}_\partial(I)$ is easily seen to be contractible, and the diffeomorphism group $\text{Diff}_\partial(\mathbb{D}^2)$ was shown to be contractible by Smale [Sma59]. The long exact sequence of the fibre bundle above then implies that $\text{Diff}_{I \sqcup I}(\mathbb{D}^2)$ is weakly contractible. \square

Notation 4.28 For simplicity, we denote the decorated surface $(\mathbb{D}^2, \underline{n})$ for each non-negative integer n by \mathbb{D}_n and called the *n -th marked 2-disc*, and the decorated surface (S, \underline{n}) by $S^{(n)}$. If $n = 0$ i.e. $\underline{n} = \emptyset$, then we simply denote $(S, \underline{0})$ by S .

The groupoid $\pi_0(\mathcal{B}r^{\mathbb{D}^2})$ is clearly isomorphic to the *braid groupoid* β which objects are non-negative integers and automorphism groups are the *classical braid groups* $\{\mathbf{B}_n\}_{n \in \mathbb{N}}$; cf. [Mac98, Chapter XI, §4] for instance. Let β^S be the groupoid $\pi_0(\mathcal{B}r^S)$. For all non-negative integers n , its automorphism groups is the surface braid group $\pi_0(\text{Diff}_{\text{dec}}^{\text{br}}(S^{(n)}))$ of S , that we denote by $\mathbf{B}_n(S)$.

By Lemma 4.3, the groupoid β inherits the braided monoidal structure of \mathcal{M}_2 which also induces a left β -module structure on $\pi_0(\mathcal{B}r^S)$. Hence we define Quillen's bracket constructions $\mathcal{U}\mathcal{B}r^{\mathbb{D}^2}$, $\langle \mathcal{B}r^{\mathbb{D}^2}, \mathcal{B}r^S \rangle$, $\mathcal{U}\beta$ and $\langle \beta, \beta^S \rangle$. By Lemma 4.4, we deduce that as semicategories $\pi_0(\mathcal{U}\mathcal{B}r^{\mathbb{D}^2}) \cong \mathcal{U}\beta$ and $\pi_0(\langle \mathcal{B}r^{\mathbb{D}^2}, \mathcal{B}r^S \rangle) \cong \langle \beta, \beta^S \rangle$.

Alternatively, braid groups on surfaces may be defined as the fundamental groups of some configuration spaces. We fix non-negative integers n and k , and a surface $S^{(n)}$ as above. The embedding space $\mathcal{E}(\underline{k}, \dot{S} \setminus \underline{n})$ is the *ordered* configuration space of k points in S , and denoted by $F_k(S^{(n)})$ (and $F_k(\mathbb{D}_n)$ if $S = \mathbb{D}^2$). Moreover, the embedding space $F_k(S^{(n)})/\mathfrak{S}_k$, induced by the natural action by permutation of coordinates of the symmetric group on the coordinates, is the *unordered* configuration space of k points in S , and denoted by $C_k(S^{(n)})$. Corollary 4.19 with $M = S$, $A = \emptyset$ and $Z = \underline{n}$ proves that $\mathbf{B}_n(S)$ is isomorphic to the fundamental group of the unordered configuration space $C_n(S)$. The group $\pi_0(\text{Diff}_{\text{dec}}^{\text{br}}(S, \underline{k}, \underline{n}))$ is called the *partitioned (k, n) -braid group* $\mathbf{B}_{k,n}(S)$.

4.6. Loop braid groups

We now focus on the families of extended and non-extended loop braid groups. Their definitions are recalled here and we refer to [Dam17] for a complete and unified introduction to these groups.

Loop braid groups may be defined in terms of *motion groups* of circles in a 3-disc. This is the setting that we shall use to construct suitable *topological* categories for the loop braid groups.

Notation 4.29 We denote by \mathbb{D}^3 the unit 3-disc. For each non-negative integer n , we denote by $\underline{n}\mathbb{S}^1$ a closed submanifold of \mathbb{D}^3 consisting in a collection of n disjoint, unknotted, oriented circles, that form a trivial link of n components in the interior of \mathbb{D}^3 . Whenever a map acts on $\underline{n}\mathbb{S}^1$, the notation $\underline{n}\mathbb{S}_+^1$ indicates that of the orientation of the circles is preserved by the map. For simplicity, we denote the decorated manifold $(\mathbb{D}^3, \underline{n}\mathbb{S}^1)$ by \mathbb{D}_n^3 and $(\mathbb{D}^3, \underline{0}\mathbb{S}^1)$ by \mathbb{D}^3 .

Let $\text{Diff}_\partial(\mathbb{D}_n^3)$ be the group of self-diffeomorphisms of \mathbb{D}^3 that fix $\partial\mathbb{D}^3$ pointwise and fix $\underline{n}\mathbb{S}^1$ as a subset. The *extended loop braid group* \mathbf{LB}'_n is the group of isotopy classes of $\text{Diff}_\partial(\mathbb{D}^3, \underline{n}\mathbb{S}^1)$. Let $\text{Diff}_\partial(\mathbb{D}^3, \underline{n}\mathbb{S}_+^1)$ be the subgroup of $\text{Diff}_\partial(\mathbb{D}^3, \underline{n}\mathbb{S}^1)$ of elements that also preserve the orientation of $\underline{n}\mathbb{S}^1$. The *(non-extended) loop braid group* \mathbf{LB}_n is the group of isotopy classes of $\text{Diff}_\partial(\mathbb{D}^3, \underline{n}\mathbb{S}_+^1)$. Finally, let $\text{Diff}_\partial(\mathbb{D}^3, \underline{n}\mathbb{S}_*^1)$ be the subgroup of $\text{Diff}_\partial(\mathbb{D}^3, \underline{n}\mathbb{S}_+^1)$ of elements that send each connected component of $\underline{n}\mathbb{S}^1$ to itself. The *pure loop braid group* \mathbf{LP}_n is the group of isotopy classes of $\text{Diff}_\partial(\mathbb{D}^3, \underline{n}\mathbb{S}_*^1)$.

We now set a categorical framework for handling these families of groups. Let \mathcal{LB}' be the full subgroupoid of \mathcal{D}_3 on those decorated manifolds (M, A, e_1, e_2) such that M is diffeomorphic to the 3-disc and A is diffeomorphic to a disjoint union of finitely many circles, embedded into the interior of M as an *unlink*. This is clearly closed under the operation \natural , so it inherits a topological semi-monoidal structure from \mathcal{D}_3 by Lemma 3.20. In the general notation of this section, we set $\mathcal{G} = \mathcal{M} = \mathcal{LB}'$.

Let \mathcal{LB} be the full subgroupoid of \mathcal{D}_3^+ on those decorated manifolds (M, A, e_1, e_2) such that M is diffeomorphic to the 3-disc and A is diffeomorphic to a disjoint union of finitely many circles, embedded into the interior of M as an *unlink*. Again, since \mathcal{LB} is clearly closed under the operation \natural , it inherits a topological semi-monoidal structure from \mathcal{D}_3^+ by Lemma 3.20. In the general notation of this section, we consider $\mathcal{G} = \mathcal{M} = \mathcal{LB}$. Note that both \mathcal{LB} and \mathcal{LB}' are subcategories of $\mathcal{D}_3^{\text{sph}}$.

We respectively denote by \mathcal{LB}' and \mathcal{LB} their discrete versions $\pi_0(\mathcal{LB}')$ and $\pi_0(\mathcal{LB})$. Let us show that there are isomorphisms $\text{Aut}_{\mathcal{LB}'}(\mathbb{D}_n^3) \cong \mathbf{LB}'_n$ and $\text{Aut}_{\mathcal{LB}}(\mathbb{D}_n^3) \cong \mathbf{LB}_n$. We focus on the extended loop braid groups case, the other one following mutatis mutandis. The automorphism group of \mathbb{D}_n^3 in \mathcal{LB}' is $\pi_0(\text{Diff}_{\text{dec}}(\mathbb{D}_n^3))$ where $\text{Diff}_{\text{dec}}(\mathbb{D}_n^3)$ is the topological group of diffeomorphisms of \mathbb{D}^3 that send the embedded the n -components unlink onto itself and that restrict to the identity on a neighbourhood of two disjoint 2-discs in $\partial\mathbb{D}^3$. The condition of fixing a *neighbourhood of two discs* in the boundary is clearly homotopy equivalent to fixing just the two discs. It therefore suffices to show that:

Lemma 4.30 *Let M be a 3-manifold with a spherical boundary-component $\partial_0 M$. Then, for isotopy classes of diffeomorphisms, fixing two disjoint 2-discs in $\partial_0 M$ is equivalent to fixing all of $\partial_0 M$.*

Proof. Let $\text{Diff}(M, \partial_0 M)$ be the group of diffeomorphisms of M that send $\partial_0 M$ to itself. The restriction map $\text{Diff}(M, \partial_0 M) \rightarrow \text{Diff}(\partial_0 M) = \text{Diff}(\mathbb{S}^2)$ is a fibre bundle, by [Cer61, Corollaire 2, §II.2.2.2, page 294]. Hence its restriction $\text{Diff}_{\mathbb{D}^2 \sqcup \mathbb{D}^2}(M) \rightarrow \text{Diff}_{\mathbb{D}^2 \sqcup \mathbb{D}^2}(\mathbb{S}^2) = \text{Diff}_{\partial C}(C)$ is also a fibre bundle, where the subscript $\mathbb{D}^2 \sqcup \mathbb{D}^2$ means that diffeomorphisms must restrict to the identity on a given pair of disjoint discs in $\partial_0 M = \mathbb{S}^2$, and C is the 2-dimensional cylinder $\mathbb{S}^1 \times [0, 1]$. The fibre is $\text{Diff}_{\partial_0 M}(M)$ and we obtain an exact sequence

$$\cdots \rightarrow \pi_1(\text{Diff}_{\partial C}(C)) \rightarrow \pi_0(\text{Diff}_{\partial_0 M}(M)) \xrightarrow{(*)} \pi_0(\text{Diff}_{\mathbb{D}^2 \sqcup \mathbb{D}^2}(M)) \rightarrow \pi_0(\text{Diff}_{\partial C}(C)).$$

By [Gra73, Théorème 1], $\text{Diff}_{\partial C}(C)$ is contractible. We deduce that $(*)$ is a bijection. \square

Therefore \mathcal{LB}' and \mathcal{LB} are respectively called the *extended loop braid groupoid* and the (*non-extended*) *loop braid groupoid*. By Lemma 4.3, the groupoids \mathcal{LB}' and \mathcal{LB} inherit a symmetric monoidal structure from the semi-monoidal structure of \mathcal{D}_3 . Hence we define Quillen's bracket constructions \mathcal{ULB}' and \mathcal{ULB} . By Lemma 4.4, we see that $\pi_0(\mathcal{ULB}') \cong \mathcal{ULB}'$ and $\pi_0(\mathcal{ULB}) \cong \mathcal{ULB}$.

Properties of loop braid groups. There are alternative ways to introduce braid groups on surfaces which we recollect here for further use. We fix non-negative integers n and k . Analogously to surface braid groups of §4.5, the group $\pi_0(\text{Diff}_{\text{dec}}(\mathbb{D}^3, \underline{k}\mathbb{S}^1, \underline{n}\mathbb{S}^1))$ is called the *partitioned (k, n) -extended loop braid group* and denoted by $\mathbf{LB}'_{k,n}$ and the group $\pi_0(\text{Diff}_{\text{dec}}(\mathbb{D}^3, \underline{k}\mathbb{S}_+^1, \underline{n}\mathbb{S}_+^1))$ is called the *partitioned (k, n) -loop braid group* and denoted by $\mathbf{LB}_{k,n}$. The embedding spaces $\mathcal{E}(\underline{k}, \mathring{\mathbb{D}}^3 \setminus \underline{n}\mathbb{S}^1)$ and $\mathcal{E}(\underline{k}\mathbb{S}^1, \mathring{\mathbb{D}}^3 \setminus \underline{n}\mathbb{S}^1)$ are the *ordered* configuration space of k *points* and k *unlinks* in \mathbb{D}_v^3 respectively, and denoted by $F_k(\mathbb{D}_n^3)$ and $\hat{U}_k(\mathbb{D}_n^3)$ respectively. Also, the embedding spaces $\mathcal{E}(\underline{k}, \mathbb{D}^3 \setminus \underline{n}\mathbb{S}^1)/\mathfrak{S}_k$ and $\mathcal{E}(\underline{k}\mathbb{S}^1, \mathring{\mathbb{D}}^3 \setminus \underline{n}\mathbb{S}^1)/\mathfrak{S}_k$ are the *unordered* configuration spaces of k *points* and k *unlinks* in \mathbb{D}_n^3 , and denoted by $C_k(\mathbb{D}_n^3)$ and $U_k(\mathbb{D}_n^3)$ respectively. Let $\text{Diff}^+(\underline{k}\mathbb{S}^1)$ be the subgroup of $\text{Diff}(\underline{k}\mathbb{S}^1)$ of those diffeomorphisms that preserve the orientation of each circle. We denote the *ordered* configuration spaces $\mathcal{E}_{\text{Diff}^+(\underline{k}\mathbb{S}^1)}(\underline{k}\mathbb{S}^1, \mathring{\mathbb{D}}^3 \setminus \underline{n}\mathbb{S}^1)$ by $\hat{U}_k^+(\mathbb{D}_n^3)$. Also, we denote the *unordered* configuration spaces $\mathcal{E}_{\text{Diff}^+(\underline{k}\mathbb{S}^1)}(\underline{k}\mathbb{S}^1, \mathring{\mathbb{D}}^3 \setminus \underline{n}\mathbb{S}^1)/\mathfrak{S}_k$ by $U_k^+(\mathbb{D}_n^3)$.

Finally, we have explicit presentations of extended and non-extended loop braid groups by generators and relations. The loop braid group \mathbf{LB}_n admits a presentation given by generators $\{\sigma_i, \tau_i \mid 1 \leq i \leq n-1\}$, where $\{\sigma_1, \dots, \sigma_{n-1}\}$ satisfy the relations of the classical braid group

\mathbf{B}_n and $\{\tau_1, \dots, \tau_{n-1}\}$ those of the symmetric group \mathfrak{S}_n and we have three additional mixed relations; see [Dam17, Proposition 3.14]. Each τ_i corresponds to a loop in $U_n^+(\mathbb{D}^3)$ that interchanges two unknots without either of them passing through the other; each σ_i corresponds to a loop in $U_n^+(\mathbb{D}^3)$ that interchanges two unknots, while one passes through the other. The pure version \mathbf{LP}_n admits a presentation given by generators $\{\chi_{ij} \mid 1 \leq i \neq j \leq n\}$ and three relations; see [Dam17, Proposition 3.18]. Finally the extended loop braid group \mathbf{LB}'_n admits a presentation with generators and relations of \mathbf{LB}_n plus additional generators $\{\rho_1, \dots, \rho_{n-1}\}$ satisfying the relations of the abelian group $(\mathbb{Z}/2\mathbb{Z})^n$ and we have five extra mixed relations; see [Dam17, Proposition 3.16]. Each ρ_i corresponds to a loop in $U_n(\mathbb{D}^3)$ that rotates a single circle by 180 degrees.

5. Global homological representations

In this section, we apply the general construction of §2 to the categories of decorated manifolds \mathcal{UD}_d introduced and studied in §3.2 and §4. This is the more geometric or topological part of our construction of representations, whereas §2 is the more algebraic and formal part.

In more detail, in §5.1, we construct continuous functors $\mathcal{UD}_d \rightarrow \text{Cov}_\bullet$, which lead, via the general construction of §2, to “global homological representations” $\pi_0(\mathcal{UD}_d) \rightarrow \text{Mod}_\bullet$. In §5.2.2–§5.2.3, we study in more detail certain particular cases of these global homological representations, restricted to the subcategories of \mathcal{UD}_2 and \mathcal{UD}_3 that are relevant for *surface braid groups*, *mapping class groups of surfaces* and *loop braid groups*. The key point is that, in many cases, these restrictions of global homological representations have image contained in a subcategory $\text{Mod}_R \subset \text{Mod}_\bullet$, for some ring R . In §5.3, we vary this construction in such a way that it is better adapted to mapping class groups of manifolds (rather than motion groups), and we study corresponding particular cases of this construction in §5.4.

Although we do not discuss it in more detail in this section, one may equally well consider examples of these global homological representations restricted to subcategories of \mathcal{UD}_2 and \mathcal{UD}_3 (or \mathcal{UD}_d for higher d) relevant for automorphism groups of free groups, Torelli groups, handlebody mapping class groups, pure braid groups and mapping class groups of non-orientable surfaces, as well as higher-dimensional motion groups.

The automorphism groups of the category \mathcal{UD}_d contain all mapping class groups and all motion groups of d -dimensional (decorated) manifolds M with *non-empty* boundary. Representations of \mathcal{UD}_d therefore give a “global” way to describe representations of all of these groups simultaneously. Our construction below does not, on the other hand, apply directly to give representations of mapping class groups or motion groups of *closed* manifolds M . However, the idea of the construction may also be carried over to this setting, to produce representations of these groups too.

5.1. Global functors for motion groups

Fix an integer $d \geq 2$, a closed submanifold $Z \subset \mathbb{R}^d$ and an open subgroup $G \leq \text{Diff}(Z)$. Also fix another integer $\ell \geq 0$. From this data, we construct two functors

$$\mathring{F}_{(Z,G,\ell)} \text{ and } F_{(Z,G,\ell)}: \mathcal{UD}_d \longrightarrow \text{Cov}_\bullet. \quad (5.1)$$

The construction of these functors occupies §5.1.1–§5.1.3, and their key basic properties are established in §5.1.4–§5.1.5. In §5.1.6 we discuss possible variations and in §5.1.7 we explain how, fixing (Z, G) and allowing ℓ to vary, these functors fit together into a tower that may be thought of as a single “pro-nilpotent” representation of the category \mathcal{UD}_d ; see diagram (5.15).

Before we begin the construction of these functors, we make a few remarks about their place in our general construction. We recall that, roughly, our general construction consists in concatenating:

1. one of the functors (5.1),
2. a functor that passes from covering spaces to bundles of modules,
3. an optional fibrewise tensor product with a functor $V: \mathcal{UD}_d \rightarrow \bullet\text{Mod}_\bullet$,
4. twisted homology.

See §2.3–§2.5 for more details of steps (2)–(4). The resulting functor takes values in the category Mod_\bullet , which is a *discrete* category, and so it factors through $\pi_0(\mathfrak{U}\mathcal{D}_d)$, which is identified canonically with $\mathfrak{U}\pi_0(\mathcal{D}_d)$, by Lemma 3.8 and Proposition 3.24, yielding a functor:

$$L_i(F_{(Z,G,\ell)}; V): \mathfrak{U}\pi_0(\mathcal{D}_d) \longrightarrow \text{Mod}_\bullet. \quad (5.2)$$

See §2.6, and in particular Lemma 2.20, for the precise construction.

Semi-functors and functors. As discussed in §3.4, the left-hand side of (5.1) is only a (topological) *semi*-category, since it is constructed from the (topological) *semi*-monoidal groupoid \mathcal{D}_d . So (5.1) is, necessarily, just a (continuous) semi-functor. However, it induces a functor on π_0 ; see Lemma 5.10 below. All of the other steps (2)–(4) in our general construction are functors, so the final output (5.2) of the construction is a functor (not just a semi-functor). We will not emphasise this small subtlety in the rest of this section, however, and in particular we will henceforth write “functor” instead of “semi-functor”.

Borel-Moore homology and functoriality. We will show (*cf.* Lemma 5.11) that the functor $F_{(Z,G,\ell)}$ takes values in the subcategory $\text{Cov}_\bullet^{\text{pr}}$ of Cov_\bullet with the same objects and whose morphisms are those which are *proper* as maps between spaces. This functor therefore works equally well as an input for our general construction when step (4) is twisted *Borel-Moore* homology, which is functorial only with respect to proper maps of spaces.

On the other hand, the functor $\mathring{F}_{(Z,G,\ell)}$ does not take values in $\text{Cov}_\bullet^{\text{pr}}$, although its restriction to the underlying groupoid \mathcal{D}_d of $\mathfrak{U}\mathcal{D}_d$ does. Thus, if we wish to use Borel-Moore homology together with the functor $\mathring{F}_{(Z,G,\ell)}$ in our general construction, it will not be fully functorial, but only functorial for automorphisms: $\pi_0(\mathcal{D}_d) \rightarrow \text{Mod}_\bullet$. In other words, we just obtain representations of the individual groups in this case.

A natural homotopy equivalence. There is (*cf.* Lemma 5.13) a natural homotopy equivalence $\mathring{F}_{(Z,G,\ell)} \Rightarrow F_{(Z,G,\ell)}$. Thus, in the case where step (4) consists of *ordinary* twisted homology (or any other homotopy invariant flavour of twisted homology), it will not matter which of these two functors ($F_{(Z,G,\ell)}$ or $\mathring{F}_{(Z,G,\ell)}$) we use in step (1).

On the other hand, the natural homotopy equivalence $\mathring{F}_{(Z,G,\ell)} \Rightarrow F_{(Z,G,\ell)}$ is not a *proper* natural homotopy equivalence. Hence, in the case where step (4) consists of *Borel-Moore* twisted homology – which is invariant only under proper homotopy equivalences – the two choices $F_{(Z,G,\ell)}$ and $\mathring{F}_{(Z,G,\ell)}$ in step (1) will lead to *a priori* different homological representations. As noted above, the homological representations obtained using Borel-Moore homology in step (4) are defined on all of $\mathfrak{U}\mathcal{D}_d$ when using $F_{(Z,G,\ell)}$ in step (1), but only on \mathcal{D}_d when using $\mathring{F}_{(Z,G,\ell)}$ in step (1). Hence we can only compare them on $\mathcal{D}_d \subset \mathfrak{U}\mathcal{D}_d$.

In certain specific cases, such as the classical *Lawrence-Bigelow representations* of the braid groups in Borel-Moore homology, if the coefficients are “generic” – a certain condition on the fibrewise tensor product taken in step (3) – the two choices of $F_{(Z,G,\ell)}$ and $\mathring{F}_{(Z,G,\ell)}$ in step (1) do in fact lead to the same homological representations, due to the fact that they actually agree with the corresponding representations using ordinary homology instead of Borel-Moore homology. This is due to [Koh17, Theorem 3.1] when the ground ring is \mathbb{C} ; see also [AP20, Proposition D] for more general ground rings.

Cohomology. One can of course replace step (4) of the general construction with twisted cohomology or twisted compactly-supported cohomology, to obtain functors of the form $\mathfrak{U}\pi_0(\mathcal{D}_d)^{\text{op}} \rightarrow \text{Mod}_\bullet$. The same considerations apply to compactly-supported cohomology as for Borel-Moore homology, so choosing the version $\mathring{F}_{(Z,G,\ell)}$ in step (1) together with twisted compactly-supported cohomology in step (4) leads only to functors $\pi_0(\mathcal{D}_d)^{\text{op}} \rightarrow \text{Mod}_\bullet$.

5.1.1. The construction of the functor on objects.

In this section and the next, we describe the construction of the functor (5.1) (on objects and on morphisms respectively) in its $\check{F}_{(Z,G,\ell)}$ (“open”) variant. The modifications involved in defining the $F_{(Z,G,\ell)}$ (“closed”) variant of (5.1) are summarised in §5.1.3.

Let $(M, A) \in \mathcal{D}_d$ be a decorated d -dimensional manifold. We show how to associate to this

- (i) a based, path-connected space $X_{(Z,G,\ell)}(M, A)$ that admits a universal cover,
- (ii) a surjective homomorphism $\phi_{(Z,G,\ell)}(M, A): \pi_1(X_{(Z,G,\ell)}(M, A)) \rightarrow Q_{(Z,G,\ell)}(M, A)$.

Together, these data determine an object of Cov_\bullet . To simplify the notation, since the choice of (Z, G, ℓ) is fixed throughout this construction, we will drop the subscripts, denoting the space by $X(M, A)$ and the surjective homomorphism by $\phi(M, A): \pi_1(X(M, A)) \rightarrow Q(M, A)$.

The space. We denote by \check{M} the interior of $\mathbb{B}_1^d \natural M$, where \natural denotes the boundary connected sum along boundary-cylinder-germs (the semi-monoidal structure of \mathcal{D}_d). Note that the interior of \mathbb{B}_1^d may be identified canonically with \mathbb{R}^d , so there is a canonical embedding $\mathbb{R}^d \hookrightarrow \check{M}$. Its image is disjoint from M , hence in particular disjoint from $A \subset M$. Thus, restricting this to $Z \subset \mathbb{R}^d$, we obtain a canonical embedding $Z \hookrightarrow \check{M} \setminus A$, which determines a basepoint of the relative embedding space

$$\mathcal{E}_G(Z, \check{M} \setminus A) = \text{Emb}(Z, \check{M} \setminus A)/G.$$

Definition 5.1 We define $X(M, A)$ to be the path-component of this space containing the basepoint.

To complete step (i) of the construction, we will show that $\mathcal{E}_G(Z, \check{M} \setminus A)$ is locally path-connected and semi-locally simply-connected (it will then follow that $X(M, A)$ also has these properties, and hence admits a universal cover). There is a quotient map

$$\text{Emb}(Z, \check{M} \setminus A) \longrightarrow \mathcal{E}_G(Z, \check{M} \setminus A). \quad (5.3)$$

Embedding spaces, equipped with the Whitney topology, are locally path-connected. Hence it follows that $\mathcal{E}_G(Z, \check{M} \setminus A)$ is also locally path-connected (since this property is preserved under taking quotients).

For semi-local simply-connectedness we need to use a stronger property of (5.3) than just the fact that it is a quotient map. The space $\mathcal{E}_G(Z, \check{M} \setminus A)$ is locally retractile with respect to the action of $\text{Diff}_c(\check{M} \setminus A)$, by [Pal18a, Proposition 4.15] (here we use the assumption that G is an *open* subgroup of $\text{Diff}(Z)$). Now, the map (5.3) is equivariant with respect to the action of $\text{Diff}_c(\check{M} \setminus A)$, so by [Pal60b, Theorem A], the map (5.3) is a fibre bundle.

In general, we have the following point-set topological lemma:

Lemma 5.2 *Let $f: X \rightarrow Y$ be a surjective fibre bundle, and suppose that X is semi-locally simply-connected. Then Y is also semi-locally simply-connected.*

Proof. Let $y \in Y$ and let U be an open neighbourhood of y in Y . We need to find a smaller open neighbourhood $V \subseteq U$ of y such that any loop in V based at y is nullhomotopic in U . First, choose a smaller open neighbourhood $U' \subseteq U$ such that f is trivialisable over U' , and choose a trivialisation $\varphi: f^{-1}(U') \cong U' \times F$. Also choose a point $z \in F$ (this is possible since we have assumed that f is surjective). Since X is semi-locally simply-connected, we may find an open neighbourhood $W \subseteq f^{-1}(U')$ of $\tilde{y} = \varphi^{-1}(y, z)$ such that any loop in W based at \tilde{y} is nullhomotopic in $f^{-1}(U')$. By the definition of the product topology, we may then find open subsets $V \subseteq U'$ and $F' \subseteq F$ such that $y \in V$, $z \in F'$ and $\varphi^{-1}(V \times F') \subseteq W$. Now let γ be any loop in V based at y . Then $\tilde{\gamma} = \varphi^{-1} \circ (\gamma \times \{z\})$ is a loop in W based at \tilde{y} . By above, we may find a nullhomotopy of $\tilde{\gamma}$ in $f^{-1}(U')$. Composing this nullhomotopy with f , it becomes a nullhomotopy of γ in $U' \subseteq U$. \square

Corollary 5.3 *The space $X(M, A)$ admits a universal cover.*

Proof. As remarked above, it will suffice to show that the space $\mathcal{E}_G(Z, \check{M} \setminus A)$ is locally path-connected and semi-locally simply-connected, since $X(M, A)$ is one path-component of this space. We have already explained why $\mathcal{E}_G(Z, \check{M} \setminus A)$ is locally path-connected. For the second property, note that the embedding space $\text{Emb}(Z, \check{M} \setminus A)$ is locally contractible, thus in particular semi-locally simply-connected. The quotient map (5.3) is a fibre bundle, so Lemma 5.2 implies that its target $\mathcal{E}_G(Z, \check{M} \setminus A)$ is also semi-locally simply-connected. \square

The surjective homomorphism. To complete the definition of the functor (5.1) on objects, we need to choose a quotient $\phi(M, A)$ of $\pi_1(X(M, A)) = \pi_1(\mathcal{E}_G(Z, \check{M} \setminus A))$.

First note that there is a canonical isomorphism $\pi_1(\mathcal{E}_G(Z, \check{M} \setminus A)) \cong \pi_1(\mathcal{E}_G(Z, \mathring{M} \setminus A))$, where \mathring{M} denotes the interior of M , given by identifying $\mathbb{B}_1^d \natural M$ with M using one of the boundary-cylinders of the decorated manifold (M, A) ;¹ see the proof of Lemma 4.12. We therefore need to choose a quotient of $\pi_1(\mathcal{E}_G(Z, \mathring{M} \setminus A))$. To do this, we use the split homotopy fibration sequence (4.4) of Lemma 4.12, which induces the split short exact sequence (4.6); this is the top row of diagram (5.4) below. We then apply the functor $G \mapsto G/\Gamma_\ell(G)$ that quotients a group by the ℓ -th term in its lower central series to the middle and right-hand terms. Finally, we factor the composition of the top-left horizontal map and the middle vertical map in the unique possible way as a surjection followed by an injection, and use these to define the left-hand vertical map $\phi(M, A)$ and the bottom-left horizontal map in the 6-term diagram below.

$$\begin{array}{ccccccc}
& & \pi_1(X(M, A)) & & & & \\
& & \downarrow \cong & & & & \\
1 & \longrightarrow & \pi_1(\mathcal{E}_G(Z, \mathring{M} \setminus A)) & \longrightarrow & \pi_1(\mathcal{E}_{\text{Diff}(A) \times G}(A \sqcup Z, \mathring{M})) & \xrightarrow{\quad \leftarrow \text{dashed} \quad} & \pi_1(\mathcal{E}(A, \mathring{M})) \longrightarrow 1 \\
& & \downarrow \phi(M, A) & & \downarrow \gamma(M, A) & & \downarrow \bar{\gamma}(M, A) \\
1 & \longrightarrow & Q(M, A) & \longrightarrow & \frac{\pi_1(\mathcal{E}_{\text{Diff}(A) \times G}(A \sqcup Z, \mathring{M}))}{\Gamma_\ell(\pi_1(\mathcal{E}_{\text{Diff}(A) \times G}(A \sqcup Z, \mathring{M})))} & \xrightarrow{\quad \leftarrow \text{dashed} \quad} & \frac{\pi_1(\mathcal{E}(A, \mathring{M}))}{\Gamma_\ell(\pi_1(\mathcal{E}(A, \mathring{M})))} \longrightarrow 1
\end{array} \tag{5.4}$$

This completes the construction of $X(M, A)$ and $\phi(M, A)$, and hence of the functor (5.1) on objects.

An immediate key observation about this diagram is that the bottom row is again a split short exact sequence; the only property of the quotient $G/\Gamma_\ell(G)$ that this uses is that it is *functorial*. We recall that Grp denotes the category of groups.

Lemma 5.4 *Let $Q: \text{Grp} \rightarrow \text{Grp}$ be a functorial quotient of groups, i.e., it is equipped with a natural transformation $q: \text{id} \Rightarrow Q$ such that $q(G): G \rightarrow Q(G)$ is a quotient map for each group G . Then for any split short exact sequence*

$$1 \longrightarrow A \xrightarrow{f} B \xrightarrow{\quad \leftarrow \text{dashed} \quad} C \longrightarrow 1$$

we have $\text{im}(q(B) \circ f: A \rightarrow B \rightarrow Q(B)) = \ker(Q(g): Q(B) \rightarrow Q(C))$. Denoting this group by $Q^(A)$, this means that we have an induced 6-term diagram*

$$\begin{array}{ccccccc}
1 & \longrightarrow & A & \xrightarrow{f} & B & \xrightarrow{\quad \leftarrow \text{dashed} \quad} & C \longrightarrow 1 \\
& & \downarrow & & \downarrow & & \downarrow \\
1 & \longrightarrow & Q^*(A) & \longrightarrow & Q(B) & \longrightarrow & Q(C) \longrightarrow 1
\end{array}$$

¹ We recall that (M, A) comes equipped with an ordered pair of *germs* of boundary-cylinders. We choose the first of this ordered pair, and then choose a boundary cylinder representing this germ. The identification of $\mathbb{B}_1^d \natural M$ with M depends on this choice of representative, but the induced isomorphism $\pi_1(\mathcal{E}_G(Z, \mathring{M} \setminus A)) \cong \pi_1(\mathcal{E}_G(Z, \check{M} \setminus A))$ does not.

in which both rows are split short exact sequences and the middle and right-hand vertical maps are given by the natural transformation q .

Proof. The second statement is clear, given the first one: we define the right-hand square by applying the functor Q to the map g and its given section and by applying the natural transformation q to the groups B and C . We then fill in the left-hand square by factoring $q(B) \circ f$ uniquely as a surjection followed by an injection. The final thing to check is exactness in the middle of the bottom row, which is precisely the first statement of the lemma.

To prove the first statement, first note that the inclusion $\text{im}(q(B) \circ f) \subseteq \ker(Q(g))$ follows immediately from exactness of the top row. To prove the opposite inclusion, let $x \in Q(B)$ with $Q(g)(x) = 1$; we need to find a lift $y \in B$ of x such that $y \in f(A)$. To do this, first pick any lift $y' \in B$ of x and set $z = g(y') \in C$. Denoting the given section of g by s , note that $s(z)$ projects to $1 \in Q(B)$, since z projects to $1 \in Q(C)$. Thus $y = y' \cdot s(z)^{-1} \in B$ is another lift of x , and moreover $g(y) = g(y') \cdot z^{-1} = z \cdot z^{-1} = 1$, so $y \in f(A)$ by exactness of the top row. \square

Notation 5.5 Two notational points concerning the above diagram:

- To be fully rigorous, the vertical morphisms in (5.4) should have subscripts with the data (Z, G, ℓ) , which have been elided above to avoid cluttering the diagram. For example, the middle vertical morphism is called $\gamma_{(Z, G, \ell)}(M, A)$ in general.
- As convention, we take $\Gamma_0(G)$ to be the trivial subgroup of G (of course, when $\ell \geq 1$, $\Gamma_\ell(G)$ is the ℓ -th term in the lower central series of G).

Remark 5.6 Note that when $\ell = 1$ we have $Q(M, A) = \{\text{id}\}$, corresponding to the trivial cover of $X(M, A)$, and when $\ell = 0$ we have $\phi(M, A) = \text{id}$, corresponding to the universal cover of $X(M, A)$.

5.1.2. The construction of the functor on morphisms.

Given a morphism $\varphi: (M, A) \rightarrow (N, B)$ in \mathcal{UD}_d , we use the identification of Proposition 4.8, which describes it as an embedding of manifolds satisfying the three properties of Definition 3.22. From this description, we see that it induces a map of split homotopy fibration sequences of the form $(4.4)_{(M, A)} \rightarrow (4.4)_{(N, B)}$, in particular a map

$$f_\varphi: \mathcal{E}_G(Z, \overset{\circ}{M} \setminus A) \longrightarrow \mathcal{E}_G(Z, \overset{\circ}{N} \setminus B).$$

Notation of the form $(4.4)_{(N, B)}$ means the diagram (4.4) with each instance of (M, A) replaced by (N, B) . If we first use the semi-monoidal structure of \mathcal{UD}_d to multiply φ with the identity map of $(\mathbb{B}_1^d, \emptyset)$, then we also obtain a map

$$f_\varphi^\sim: \mathcal{E}_G(Z, \overset{\circ}{M} \setminus A) \longrightarrow \mathcal{E}_G(Z, \overset{\circ}{N} \setminus B),$$

which preserves basepoints and therefore restricts to a based map

$$f_\varphi^X: X(M, A) \longrightarrow X(N, B).$$

As explained in §5.1.1 above, $\pi_1(X(M, A))$ is naturally identified with $\pi_1(\mathcal{E}_G(Z, \overset{\circ}{M} \setminus A))$ (and similarly for (N, B)); under these identifications the homomorphisms $\pi_1(f_\varphi)$ and $\pi_1(f_\varphi^X)$ agree.

The map of split homotopy fibration sequences $(4.4)_{(M, A)} \rightarrow (4.4)_{(N, B)}$ induces a map of split short exact sequences $(4.6)_{(M, A)} \rightarrow (4.6)_{(N, B)}$, and hence (since the construction of (5.4) from (4.6) is functorial) a map of diagrams of the form $(5.4)_{(M, A)} \rightarrow (5.4)_{(N, B)}$, in particular a homomorphism

$$\theta_\varphi: Q(M, A) \longrightarrow Q(N, B).$$

That this is a map of diagrams says, in particular, that

$$\theta_\varphi \circ \phi(M, A) = \phi(N, B) \circ \pi_1(f_\varphi) = \phi(N, B) \circ \pi_1(f_\varphi^X),$$

and hence $\pi_1(f_\varphi^X)$ sends $\ker(\phi(M, A))$ into $\ker(\phi(N, B))$. Thus $(f_\varphi^X, \theta_\varphi)$ is a morphism in Cov from $(X(M, A), \phi(M, A))$ to $(X(N, B), \phi(N, B))$; cf. Definition 2.2. This completes the definition of the functor (5.1) on morphisms.

Two actions agree. The entire diagram (5.4) is functorial in the input (M, A) as an object of \mathcal{UD}_d . If we restrict to the automorphism group $\text{Diff}_{\text{dec}}(M, A)$ of a single object (M, A) and look just at the bottom-left group $Q(M, A)$ of (5.4), we obtain an action of $\text{Diff}_{\text{dec}}(M, A)$ on $Q(M, A)$. Since $Q(M, A)$ is discrete, this factors through an action of $\pi_0(\text{Diff}_{\text{dec}}(M, A))$ on $Q(M, A)$. If we restrict this further to the subgroup $\pi_0(\text{Diff}_{\text{dec}}^{\text{br}}(M, A)) \subseteq \pi_0(\text{Diff}_{\text{dec}}(M, A))$, we obtain an action of $\pi_0(\text{Diff}_{\text{dec}}^{\text{br}}(M, A))$ on $Q(M, A)$.

We also have an action of the top-right group of (5.4) on the bottom-left group $Q(M, A)$ of (5.4), given either by lifting elements of $Q(M, A)$ along the vertical map $\phi(M, A)$ and using the conjugation action of the semi-direct product on the top row, or equivalently by projecting along the vertical map $\bar{\gamma}(M, A)$ to the bottom-right group of (5.4) and then using the conjugation action of the semi-direct product on the bottom row. By Proposition 4.23, the top-right group of (5.4) is naturally identified with $\pi_0(\text{Diff}_{\text{dec}}^{\text{br}}(M, A))$, so this gives us another action of $\pi_0(\text{Diff}_{\text{dec}}^{\text{br}}(M, A))$ on $Q(M, A)$.

Proposition 5.7 *The two actions of $\pi_0(\text{Diff}_{\text{dec}}^{\text{br}}(M, A))$ on $Q(M, A)$ described above are equal.*

Proof. This follows immediately from the second part of Lemma 4.25. \square

5.1.3. A closed variant.

The construction of the “closed” variant $F_{(Z, \mathcal{G}, \ell)}$ of the functor (5.1) is a very slight modification of the constructions in §5.1.1 and §5.1.2, in which we constructed the “open” variant $\mathring{F}_{(Z, \mathcal{G}, \ell)}$.

To do this, we simply replace \mathring{M} with M (and similarly \mathring{M} with $\mathbb{B}_1^d \natural M$) everywhere, and we consider the embedding space $\mathcal{E}_G^{\ell}(Z, M \setminus A)$ of Definition 4.14 instead of $\mathcal{E}_G(Z, \mathring{M} \setminus A)$. In other words, we consider embeddings of Z whose image is either contained entirely in the interior or entirely in the boundary of M . This has the effect of replacing $X(M, A)$ with another space that is homotopy equivalent (cf. Lemma 4.15),² so it does not change $\pi_1(X(M, A))$.³ The construction of the quotient $\phi(M, A): \pi_1(X(M, A)) \rightarrow Q(M, A)$ is identical. On morphisms, the construction is exactly analogous, viewing the split short exact sequence (4.6) as (4.7).

5.1.4. Elementary properties.

By Lemma 3.7, if we have subgroupoids \mathcal{M} and \mathcal{G} of \mathcal{D}_d such that \mathcal{G} is closed under \natural and \mathcal{M} is closed under the action of \mathcal{G} via \natural , and if \mathcal{G} is moreover a *full* subgroupoid of \mathcal{D}_d , then there is a natural inclusion of categories

$$\langle \mathcal{G}, \mathcal{M} \rangle \hookrightarrow \mathcal{UD}_d. \quad (5.5)$$

On objects, this is just the inclusion of the objects of \mathcal{M} into the objects of \mathcal{D}_d . On morphisms, under the identification of morphism spaces of Proposition 4.8, this is an inclusion of embedding spaces (roughly – see Definition 4.7 for precise details – morphisms in \mathcal{UD}_d are given by embeddings whose complement is another decorated manifold, whereas morphisms in $\langle \mathcal{G}, \mathcal{M} \rangle$ are given by embeddings whose complement is a decorated manifold in the set $\text{ob}(\mathcal{G})$).

Summarising the constructions in §5.1.1–§5.1.3, we have:

Proposition 5.8 *For any integers $d \geq 2$ and $\ell \geq 0$, closed submanifold $Z \subset \mathbb{R}^d$ and open subgroup $G \leq \text{Diff}(Z)$, the recipe described above gives well-defined functors*

$$\mathring{F}_{(Z, \mathcal{G}, \ell)} \text{ and } F_{(Z, \mathcal{G}, \ell)}: \mathcal{UD}_d \longrightarrow \text{Cov}_{\bullet}, \quad (5.6)$$

and hence, by restriction, well-defined functors $\mathring{F}_{(Z, \mathcal{G}, \ell)}$ and $F_{(Z, \mathcal{G}, \ell)}: \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \text{Cov}_{\bullet}$, for any subcategory $\langle \mathcal{G}, \mathcal{M} \rangle \subseteq \mathcal{UD}_d$ as in (5.5).

² Although not *proper* homotopy equivalent, which is why the open and closed variants of (5.1) give rise to different homological representations when using Borel-Moore homology.

³ The only small technicality comes when checking that the closed version of $X(M, A)$ has good local properties, so it admits a universal cover (Corollary 5.3): the argument is exactly analogous, except that one has to use a variant of [Pal18a, Proposition 4.15] where the target manifold is allowed to have non-empty boundary. This may be proved by a small adaptation of the proof of Proposition 4.15 of [Pal18a], using ideas of [Cer61] (where all manifolds are allowed to have corners of any codimension).

Remark 5.9 Note that the construction, up to homotopy, depends only on the isotopy class (mod G) of the embedded submanifold $Z \subseteq \mathbb{R}^d$. More precisely, if we choose two embeddings of the closed manifold Z in \mathbb{R}^d such that the corresponding points of $\text{Emb}(Z, \mathbb{R}^d)/G$ lie in the same path-component, then the two resulting continuous functors $\mathfrak{UD}_d \rightarrow \text{Cov}_\bullet$ will be homotopic.

We now record a few immediate observations about the the topological functors (5.6). First, recall that these are in fact only semifunctors, since \mathfrak{UD}_d is only a semi-category (cf. §3.4). The first observation says that this technical issue does not matter once we pass to π_0 .

Lemma 5.10 *The topological semifunctors (5.6) induce functors on π_0 .*

Proof. Let (M, A) be an object of $\pi_0(\mathfrak{UD}_d)$ (i.e., an object of \mathcal{D}_d , a decorated manifold). We have to show that $\pi_0(5.6)$ sends $\text{id}_{(M,A)}$ to an identity morphism of $\pi_0(\text{Cov}_\bullet)$. To do this, we first have to find the identity morphism of (M, A) in $\pi_0(\mathfrak{UD}_d)$.

Let $e_1: \mathbb{D}^{d-1} \times [0, 1] = \mathbb{B}_1^d \hookrightarrow M$ denote one of the boundary cylinders that (M, A) is equipped with (more precisely, a representative of one of the *germs of boundary cylinders* that (M, A) is equipped with). The boundary connected sum $(\mathbb{B}_1^d, \emptyset) \natural (M, A)$ may be viewed as the union of $\mathbb{D}^{d-1} \times [-1, 1]$ with M along $\mathbb{D}^{d-1} \times [0, 1]$ via the embedding e_1 . Choose a diffeomorphism $[-1, 1] \rightarrow [0, 1]$ that is given by $t \mapsto t + 1$ on $[-1, -1 + \epsilon]$ and by $t \mapsto t$ on $[1 - \epsilon, 1]$ for some $\epsilon > 0$. Multiplying this by \mathbb{D}^{d-1} and extending by the identity over $M \setminus \text{im}(e_1)$, this determines an isomorphism of decorated manifolds

$$v_{(M,A)}: (\mathbb{B}_1^d, \emptyset) \natural (M, A) \longrightarrow (M, A). \quad (5.7)$$

Consider the endomorphism of (M, A) in \mathfrak{UD}_d given by

$$\Upsilon_{(M,A)} = ((\mathbb{B}_1^d, \emptyset), v_{(M,A)}). \quad (5.8)$$

One may check that, for any endomorphism φ of (M, A) in \mathfrak{UD}_d , the compositions $\Upsilon_{(M,A)} \circ \varphi$ and $\varphi \circ \Upsilon_{(M,A)}$ are both isotopic to φ . Hence $\Upsilon_{(M,A)}$ is the identity of (M, A) in the category $\pi_0(\mathfrak{UD}_d)$. Under the identification of Proposition 4.8, this corresponds to the self-embedding of (M, A) given by restricting the diffeomorphism (5.7) to the submanifold $(M, A) \subset (\mathbb{B}_1^d, \emptyset) \natural (M, A)$. Since this self-embedding is isotopic to the identity, the induced self-map of embedding spaces $X(M, A) \rightarrow X(M, A)$ is homotopic to the identity, and hence is an identity morphism in $\pi_0(\text{Cov}_\bullet)$. \square

Lemma 5.11 *The topological semifunctor $F_{(Z,G,\ell)}$ of (5.6) takes values in the subcategory $\text{Cov}_\bullet^{\text{pr}}$ of Cov_\bullet .*

Proof. We have to show that, for any morphism $\varphi: (M, A) \rightarrow (N, B)$ of decorated manifolds, the induced map of spaces $X(M, A) \rightarrow X(N, B)$ (in the *closed* variant of the construction in §5.1.1) is a proper map (preimages of compact subspaces are compact). We recall that this is (a restriction to particular path-components of) an inclusion of embedding spaces

$$\mathcal{E}'_G(Z, \mathbb{B}_1^d \natural M \setminus A) \longrightarrow \mathcal{E}'_G(Z, \mathbb{B}_1^d \natural N \setminus B)$$

induced by an embedding of pairs of manifolds $(M, A) \hookrightarrow (N, B)$ satisfying the three properties of Definition 3.22 (cf. Proposition 4.8). In particular, the third property implies that the inclusion $M \setminus A \hookrightarrow N \setminus B$ has closed image, so the inclusion of embedding spaces above also has closed image; any closed inclusion is a proper map. \square

Remark 5.12 On the other hand, the topological semifunctor $\mathring{F}_{(Z,G,\ell)}$ of (5.6) does *not* take values in the subcategory $\text{Cov}_\bullet^{\text{pr}}$ of Cov_\bullet . The above proof breaks down in this setting because the inclusion of the *interior* of $M \setminus A$ into the *interior* of $N \setminus B$ does not have closed image.

Lemma 5.13 *There is a natural homotopy equivalence $\mathring{F} \Rightarrow F$ between the two functors (5.6).*

Proof. The natural transformation may easily be constructed from the inclusions of spaces of embeddings into the interior of $M \setminus A$ into spaces of embeddings with image contained either in the interior or the boundary of $M \setminus A$. The fact that this is a natural *homotopy equivalence* (although not a natural *proper* homotopy equivalence) follows from Lemma 4.15. \square

5.1.5. The image of the functor.

Under certain conditions, the functor (5.6), restricted to a subcategory of the form $\langle \mathcal{G}, \mathcal{M} \rangle$ from (5.5), takes values in a subcategory of Cov_\bullet of the form Cov_Q or Cov_Q^{tw} for a fixed group Q .

Definition 5.14 (*Q-stability*.) Let \mathcal{M} and \mathcal{G} be subgroupoids as in (5.5). Suppose that there is a group Q and there are identifications $Q(M, A) \cong Q$ for each object (M, A) of $\langle \mathcal{G}, \mathcal{M} \rangle$ (recall that these are exactly the objects of \mathcal{M}), such that, for each object (M', A') of \mathcal{G} , the homomorphism of groups

$$Q(M, A) \longrightarrow Q((M', A') \natural (M, A)),$$

induced by the canonical morphism $((M', A'), \text{id})$ of $\langle \mathcal{G}, \mathcal{M} \rangle$, is equal to the identity under these identifications. In this case, we say that *the functor $F_{(Z, \mathcal{G}, \ell)}$ is Q-stable on $\langle \mathcal{G}, \mathcal{M} \rangle$* .

We recall that $\text{Cov}_Q^{\text{tw}} \subset \text{Cov}_\bullet$ is the full subcategory on objects $(X, \phi: \pi_1(X) \twoheadrightarrow Q')$ where $Q' = Q$. As an immediate observation, we note:

Lemma 5.15 *Suppose that $F_{(Z, \mathcal{G}, \ell)}$ is Q-stable on $\langle \mathcal{G}, \mathcal{M} \rangle$. Then $F_{(Z, \mathcal{G}, \ell)}$ is equivalent to a functor with image contained in $\text{Cov}_Q^{\text{tw}} \subset \text{Cov}_\bullet$.*

Proof. The hypothesis implies that the image of $F_{(Z, \mathcal{G}, \ell)}$ is contained in the slightly larger full subcategory $\text{Cov}_{\cong Q}^{\text{tw}} \subset \text{Cov}_\bullet$ on objects $(X, \phi: \pi_1(X) \twoheadrightarrow Q')$ where Q' is *isomorphic* to Q , and the inclusion $\text{Cov}_Q^{\text{tw}} \hookrightarrow \text{Cov}_{\cong Q}^{\text{tw}}$ is an equivalence of categories. \square

Under certain additional conditions, when $\ell = 2$, we may restrict the image further to Cov_Q , after passing to π_0 . Write $|\mathcal{G}|$ for the set of isomorphism classes of objects of $\pi_0(\mathcal{G})$, which is naturally a monoid, and similarly write $|\mathcal{M}|$ for the set of isomorphism classes of objects of $\pi_0(\mathcal{M})$, which is naturally a $|\mathcal{G}|$ -set.

Proposition 5.16 *Let \mathcal{M} and \mathcal{G} be subgroupoids of \mathcal{D}_d such that \mathcal{G} is closed under \natural and \mathcal{M} is closed under the action of \mathcal{G} via \natural . Set $\ell = 2$, suppose that $F_{(Z, \mathcal{G}, 2)}$ is Q-stable on $\langle \mathcal{G}, \mathcal{M} \rangle$, and*

- \mathcal{G} is full in \mathcal{D}_d and \mathcal{M} is 0-full in \mathcal{D}_d ,
- for each object (M, A) of \mathcal{M} , the subgroup $\text{Aut}_{\mathcal{M}}(M, A) \subseteq \text{Diff}_{\text{dec}}(M, A)$ lies in $\text{Diff}_{\text{dec}}^{\text{br}}(M, A)$,
- $|\mathcal{G}|$ is free as a monoid and $|\mathcal{M}|$ is free as a $|\mathcal{G}|$ -set.

Under these assumptions, the functor

$$\pi_0(F_{(Z, \mathcal{G}, 2)}|_{\langle \mathcal{G}, \mathcal{M} \rangle}): \pi_0(\langle \mathcal{G}, \mathcal{M} \rangle) \longrightarrow \pi_0(\text{Cov}_\bullet)$$

is equivalent to a functor with image contained in $\pi_0(\text{Cov}_Q) \subset \pi_0(\text{Cov}_\bullet)$.

Proof. To prove this, we will go step by step through the following diagram, where $F = F_{(Z, \mathcal{G}, 2)}|_{\langle \mathcal{G}, \mathcal{M} \rangle}$.

$$\begin{array}{ccc} \langle \mathcal{G}, \mathcal{M} \rangle & \xrightarrow{F \cong \hat{F}} & \text{Cov}_\bullet \\ \downarrow & & \downarrow \\ \pi_0(\langle \mathcal{G}, \mathcal{M} \rangle) & \cong & \langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle \xrightarrow{\pi_0(\hat{F})} \pi_0(\text{Cov}_\bullet) \\ & \uparrow & \uparrow \\ & \langle \mathcal{G}_0, \mathcal{M}_0 \rangle & \xrightarrow{\pi_0(\hat{F})} \pi_0(\text{Cov}_Q) \end{array} \quad (5.9)$$

The functor \hat{F} is defined just like F , except that, on objects, the surjection $\phi(M, A): \pi_1(X(M, A)) \twoheadrightarrow Q(M, A)$ is composed with the given identification $Q(M, A) \cong Q$ from the definition of Q -stability. It is easy to see that this is again a well-defined functor, and the isomorphisms $Q(M, A) \cong Q$ induce an isomorphism of functors $F \cong \hat{F}$. Clearly, \hat{F} has image contained in Cov_Q^{tw} : this is a slightly different alternative proof of Lemma 5.15, with the slightly stronger conclusion that $F_{(Z, \mathcal{G}, \ell)}$ is *isomorphic* to a functor with image contained in Cov_Q^{tw} . Moreover, the definition of Q -stability

immediately tells us that each canonical morphism $(M', \text{id}): M \rightarrow M' \natural M$ in $\langle \mathcal{G}, \mathcal{M} \rangle$ is sent by \hat{F} to the subcategory Cov_Q .

We now pass to the middle row of diagram (5.9). Using the first assumption of the proposition, Lemma 4.4 gives us a canonical isomorphism of categories $\pi_0(\langle \mathcal{G}, \mathcal{M} \rangle) \cong \langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle$, so we may view $\pi_0(\hat{F})$ as defined on the category $\langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle$. Since $\pi_0(\mathcal{G})$ has a monoidal unit $\mathbf{1}$, we may consider, for each automorphism φ of (M, A) in $\pi_0(\mathcal{M})$, the automorphism $(\mathbf{1}, \varphi)$ of (M, A) in $\langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle$. More precisely, since the unit $\mathbf{1} = (\mathbb{B}_1^d, \emptyset)$ of $\langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle$ is not *strict*, the corresponding automorphism of (M, A) in $\langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle$ is given by $(\mathbf{1}, \varphi \circ \lambda_{(M, A)})$, where λ is the left unitor for the monoidal structure. Under the functor $\pi_0(\hat{F})$, this is sent to an automorphism of the object $(X(M, A), \pi_1(X(M, A))) \rightarrow Q(M, A) \cong Q$, so in particular it induces an automorphism of Q . We now show that this is the *identity* automorphism; in other words, this says that $\pi_0(\hat{F})$ sends $(\mathbf{1}, \varphi)$ into the subcategory Cov_Q . Since the induced automorphism of Q is pulled back along the given identification $Q(M, A) \cong Q$ from the induced automorphism of $Q(M, A)$, it is equivalent to show that the induced automorphism of $Q(M, A)$ is the identity. To see this, note that, by the second assumption of the proposition and by Proposition 5.7, the action of $(\mathbf{1}, \varphi)$ on $Q(M, A)$ is given by viewing φ as an element of the top-right group of diagram (5.4) (via the identification of Proposition 4.23), projecting it to the bottom-right group and then letting it act on $Q(M, A)$ via the semi-direct product structure of the bottom row of (5.4). But we have assumed that $\ell = 2$, so all groups on the bottom row of (5.4) are abelian, so the semi-direct product is a *direct* product, and the action on $Q(M, A)$ is trivial.

Putting together the two paragraphs above, we see that the functor $\pi_0(\hat{F})$ has image contained in $\pi_0(\text{Cov}_Q^{\text{tw}})$, and moreover it takes all canonical morphisms (M', id) and all automorphisms of the form $(\mathbf{1}, \varphi)$ (for φ an automorphism of $\pi_0(\mathcal{M})$) into the subcategory $\pi_0(\text{Cov}_Q)$. However, we cannot yet conclude: although each morphism of $\langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle$ decomposes into a composition of a canonical morphism (M', id) followed by $(\mathbf{1}, \varphi)$ for an isomorphism φ of $\pi_0(\mathcal{M})$, it is not necessarily the case that φ is an *automorphism* of $\pi_0(\mathcal{M})$.

To deal with this, we now choose skeleta $\mathcal{G}_0 \subseteq \pi_0(\mathcal{G})$ and $\mathcal{M}_0 \subseteq \pi_0(\mathcal{M})$ such that:

- \mathcal{G}_0 is a *monoidal* subcategory of $\pi_0(\mathcal{G})$, and
- \mathcal{M}_0 is a \mathcal{G}_0 -*module* subcategory of $\pi_0(\mathcal{M})$.

It is possible to construct skeleta with these properties due to the third assumption of the proposition. Since the set $|\mathcal{G}|$ of isomorphism classes of objects of $\pi_0(\mathcal{G})$ is a *free* monoid, we may choose a free generating set for $|\mathcal{G}|$, then choose arbitrarily one object of $\pi_0(\mathcal{G})$ in each isomorphism class corresponding to a generator, and then use the monoidal structure of $\pi_0(\mathcal{G})$ to choose all other representatives of isomorphism classes of objects. This procedure is well-defined since we used a *free* generating set for the monoid $|\mathcal{G}|$. We then define \mathcal{G}_0 to be the full subcategory of $\pi_0(\mathcal{G})$ on the objects that we have chosen in this way. Exactly the same idea allows us to construct a skeleton \mathcal{M}_0 of $\pi_0(\mathcal{M})$ that is closed under the action of $\mathcal{G}_0 \subseteq \pi_0(\mathcal{G})$, using a free generating set for $|\mathcal{M}|$ as a $|\mathcal{G}|$ -set.

It is then routine to verify that there is a canonical functor $\langle \mathcal{G}_0, \mathcal{M}_0 \rangle \rightarrow \langle \pi_0(\mathcal{G}), \pi_0(\mathcal{M}) \rangle$ that is full, faithful and essentially surjective on objects. In other words, it is an inclusion of a subcategory that is also an equivalence of categories. It therefore suffices to show that $\pi_0(\hat{F})$, restricted to $\langle \mathcal{G}_0, \mathcal{M}_0 \rangle$, takes values in $\pi_0(\text{Cov}_Q)$. But this now follows from what we have already shown above. We recall that we have shown that $\pi_0(\hat{F})$ sends any morphism of the form (M', id) or $(\mathbf{1}, \varphi)$ for φ an automorphism of $\pi_0(\mathcal{M})$ into $\pi_0(\text{Cov}_Q)$. Any morphism in $\langle \mathcal{G}_0, \mathcal{M}_0 \rangle$ decomposes as $(\mathbf{1}, \varphi) \circ (M', \text{id})$ for an isomorphism φ of \mathcal{M}_0 . But \mathcal{M}_0 is a skeletal category, so φ must be an *automorphism*. \square

In several of our examples, Q -stability does not hold on $\langle \mathcal{G}, \mathcal{M} \rangle \subseteq \mathcal{UD}_d$ (for certain \mathcal{G} and \mathcal{M}), but *does* hold on the full subcategory $\langle \mathcal{G}, \mathcal{M}' \rangle \subseteq \langle \mathcal{G}, \mathcal{M} \rangle$ for a certain full sub- \mathcal{G} -module $\mathcal{M}' \subset \mathcal{M}$. Typically, the isomorphism classes of objects of \mathcal{G} will form the monoid \mathbb{N} , the isomorphism classes of objects of \mathcal{M} will form the \mathbb{N} -module \mathbb{N} (acting on itself by addition), and \mathcal{M}' will correspond to the sub- \mathbb{N} -module $\{n, n+1, n+2, \dots\}$ for some $n \geq 1$.

The functor $F_{(\mathcal{Z}, \mathcal{G}, 2)}$ restricted to $\pi_0(\langle \mathcal{G}, \mathcal{M}' \rangle)$ will therefore factor (up to equivalence) through $\pi_0(\text{Cov}_Q)$, and hence the general construction will result in a functor of the form $\pi_0(\langle \mathcal{G}, \mathcal{M}' \rangle) \rightarrow$

Mod_R for some ring R . It will sometimes be convenient to extend this functor “trivially” to the larger category $\pi_0(\langle \mathcal{G}, \mathcal{M} \rangle)$ with the help of the following observation.

Lemma 5.17 *Let \mathcal{D} be a category and $\mathcal{C} \subseteq \mathcal{D}$ the full subcategory on a subset $\text{ob}(\mathcal{C}) \subseteq \text{ob}(\mathcal{D})$. Assume that there are no morphisms in \mathcal{D} from $\text{ob}(\mathcal{C})$ to $\text{ob}(\mathcal{D}) \setminus \text{ob}(\mathcal{C})$. Then any functor $\mathcal{C} \rightarrow \mathcal{A}$ to a category \mathcal{A} with initial object I may be extended to \mathcal{D} by sending each object of $\text{ob}(\mathcal{D}) \setminus \text{ob}(\mathcal{C})$ to I .*

5.1.6. Choosing quotients.

The construction of quotients of $\pi_1(X(M, A))$ in diagram (5.4) admits several natural generalisations.

Functorial descending normal series. We recall that a *functorial descending normal series* is a sequence of endofunctors $s_i: \text{Grp} \rightarrow \text{Grp}$ with $s_1 = \text{id}$, such that $s_{i+1}(G) \subseteq s_i(G)$ for each i and these inclusions assemble to a natural transformation $s_{i+1} \Rightarrow s_i$, and each subgroup $s_i(G) \subseteq G$ is normal in G . Examples are the lower central series and derived series. In the construction of (5.1), we have used the lower central series to define certain quotients of fundamental groups. *However, exactly the same construction goes through if one replaces the lower central series by any other functorial descending normal series.* We have chosen to focus on constructions based on the lower central series due to the fact that there are many results in the literature about the lower central quotients of surface braid groups (and hence one may also deduce results about lower central quotients of *partitioned* surface braid groups, cf. [DPS21]) – see the later subsections of this section – which are useful for understanding specific examples of homological representations fitting into this framework.

Functorial quotients of groups. More generally, it would suffice to fix any functorial choice of normal subgroup (rather than an entire functorial descending normal series), in other words, a functor $s: \text{Grp} \rightarrow \text{Grp}$ with the property that $s(G) \triangleleft G$ for all groups G and $s(\phi) = \phi|_G$ for any homomorphism $\phi: G \rightarrow H$ (so that the inclusions assemble into a natural transformation $s \Rightarrow \text{id}$). We may then define $\gamma(M, A)$ and $\bar{\gamma}(M, A)$ in (5.4) by the natural transformation $G \mapsto G/s(G)$.

Transfinite series. The lower central series (and derived series) may be extended to *transfinite* series, indexed by arbitrary ordinals, and indeed one may take ℓ to be any (possibly infinite) ordinal in the construction. Infinite values of ℓ may be relevant for constructing homological representations of motion groups (for example loop-braid groups in 3-manifolds), since there exist 3-manifolds whose fundamental groups have lower central series that stop only at ordinals greater than ω [CO98].

Invariant non-functorial quotients. The diagram (5.4) is functorial in $\mathcal{M}\mathcal{D}_d$, so in particular there is an action of $\text{Aut}_{\mathcal{D}_d}(M, A)$ on $Q(M, A)$. If we focus on the single object (M, A) and its automorphisms, we may wish to modify the construction so that the quotient of $\pi_1(X(M, A))$ has the *trivial* action of $\text{Aut}_{\mathcal{D}_d}(M, A)$. There are two options:

- Restrict the automorphism group $\text{Aut}_{\mathcal{D}_d}(M, A)$ to the kernel of its action on $Q(M, A)$; see §5.4.1.2 for a specific example of this.
- Replace $Q(M, A)$ by a further quotient that is invariant under the action of $\text{Aut}_{\mathcal{D}_d}(M, A)$; see [PS21, §3] for a specific example of this.

5.1.7. Pro-nilpotent representations.

If we fix the inputs $Z \subseteq \mathbb{R}^d$ and $G \leq \text{Diff}(Z)$ in our construction and allow the integer ℓ to vary, one may package together the homological representations arising from the general construction for each value of ℓ into a kind of “pro-nilpotent” homological representation, which may be

truncated to recover the representation corresponding to each level ℓ . Namely, the resulting homological representations fit together into a tower, which one may think of as a single *pro-nilpotent representation*. In this section we explain how this may be done.

Remark 5.18 More generally, one may do this for any functorial descending normal series in place of the lower central series; in particular one may easily adapt this section to describe “pro-solvable” homological representations. In addition, one may also do this for any limit ordinal λ in place of ω , to obtain “pro- λ -nilpotent” homological representations, etc.

Definition 5.19 For an integer $\ell \geq 0$, let $\text{Cov}_\bullet^{(\ell)}$ be the full subcategory of Cov_\bullet on those objects $(X, \pi_1(X) \twoheadrightarrow Q)$ where the group Q has nilpotency class at most ℓ . If $\ell = 0$ this means Q must be the trivial group, corresponding to the trivial covering of X . If $\ell = 1$, this means that Q must be abelian, so $\text{Cov}_\bullet^{(1)}$ is the category of spaces equipped with abelian coverings.

Definition 5.20 Fix a unital ring \mathbb{A} and an integer $\ell \geq 0$, and let $\text{Mod}_\bullet^{(\ell)}$ be the full subcategory of Mod_\bullet on those objects (R, V) where $R = \mathbb{A}[Q]$ for a group Q of nilpotency class at most ℓ . Similarly, $\text{Top}_\bullet^{(\ell)}$ is the full subcategory of Top_\bullet on those objects (X, R, ξ) where $R = \mathbb{A}[Q]$ for a group Q of nilpotency class at most ℓ (and ξ is a bundle of right R -modules over the space X).

Construction 5.21 For $k \geq 0$, there is a functor

$$\varpi_k: \text{Cov}_\bullet \longrightarrow \text{Cov}_\bullet^{(k)},$$

which is a section for the inclusion $\text{Cov}_\bullet^{(k)} \subseteq \text{Cov}_\bullet$, defined as follows. On objects, it is given by

$$(X, \pi_1(X) \twoheadrightarrow Q) \longmapsto (X, \pi_1(X) \twoheadrightarrow Q \twoheadrightarrow Q/\Gamma_k(Q)).$$

We recall that a morphism $(X, \pi_1(X) \twoheadrightarrow Q) \rightarrow (X', \pi_1(X') \twoheadrightarrow Q')$ is a based map $f: X \rightarrow X'$ with the property that $\pi_1(f)$ descends to a map $Q \rightarrow Q'$. Clearly this implies that it also descends to a map $Q/\Gamma_k(Q) \rightarrow Q'/\Gamma_k(Q')$. Hence we may define the functor on morphisms simply by $f \mapsto f$.

Construction 5.22 Similarly, for $k \geq 0$ there are functors

$$\varpi_k: \text{Mod}_\bullet \longrightarrow \text{Mod}_\bullet^{(k)} \quad \text{and} \quad \varpi_k: \text{Top}_\bullet \longrightarrow \text{Top}_\bullet^{(k)}$$

which are sections for the inclusions $\text{Mod}_\bullet^{(k)} \subseteq \text{Mod}_\bullet$ and $\text{Top}_\bullet^{(k)} \subseteq \text{Top}_\bullet$, defined as follows. The first is defined on objects by

$$(\mathbb{A}[Q], V) \longmapsto (\mathbb{A}[Q/\Gamma_k(Q)], V \otimes_{\mathbb{A}[Q]} \mathbb{A}[Q/\Gamma_k(Q)])$$

and the second is defined on objects by

$$(X, \mathbb{A}[Q], \xi: \pi_{\leq 1}(X) \rightarrow \text{Mod}_{\mathbb{A}[Q]}) \longmapsto (X, \mathbb{A}[Q/\Gamma_k(Q)], \pi_{\leq 1}(X) \rightarrow \text{Mod}_{\mathbb{A}[Q]} \rightarrow \text{Mod}_{\mathbb{A}[Q/\Gamma_k(Q)]}),$$

where the functor $\text{Mod}_{\mathbb{A}[Q]} \rightarrow \text{Mod}_{\mathbb{A}[Q/\Gamma_k(Q)]}$ is given by $-\otimes_{\mathbb{A}[Q]} \mathbb{A}[Q/\Gamma_k(Q)]$. As in the construction above, it is easy to see how to extend these constructions to morphisms.

The functors $\text{Lift}: \text{Cov}_\bullet \rightarrow \text{Top}_\bullet$ and $H_i: \text{Top}_\bullet \rightarrow \text{Mod}_\bullet$ respect the filtrations of these categories given by nilpotency class. With respect to the functors ϖ_k , they fit together as follows:

$$\begin{array}{ccccc} \text{Cov}_\bullet & \xrightarrow{\text{Lift}} & \text{Top}_\bullet & \xrightarrow{H_i} & \text{Mod}_\bullet \\ \varpi_k \downarrow & \parallel & \varpi_k \downarrow & \not\parallel & \downarrow \varpi_k \\ \text{Cov}_\bullet^{(k)} & \xrightarrow{\text{Lift}} & \text{Top}_\bullet^{(k)} & \xrightarrow{H_i} & \text{Mod}_\bullet^{(k)} \end{array} \quad (5.10)$$

The left-hand square commutes on the nose, but the second one does not. However, for any space X , local system \mathcal{L} on X defined over a ring R and ring homomorphism $\theta: R \rightarrow S$, there is a natural homomorphism of S -modules

$$H_i(X; \mathcal{L}) \otimes_R S \longrightarrow H_i(X; \mathcal{L} \otimes_R S),$$

where S is viewed as an (R, S) -bimodule via θ . This homomorphism is the subject of the universal coefficient theorem (although we will not need more than its existence). This natural homomorphism provides the natural transformation filling the right-hand square as indicated.

Lemma 5.23 *The homological functor $F_{(Z,G,\ell)}: \mathfrak{UD}_d \rightarrow \text{Cov}_\bullet^{(\ell)}$ has image contained in $\text{Cov}_\bullet^{(\ell)}$.*

Proof. This follows from the fact that, in diagram (5.4), the middle group on the bottom row has nilpotency class at most ℓ by construction. The property of having nilpotency class at most ℓ passes to subgroups, so $Q(M, A)$ also has nilpotency class at most ℓ . \square

This means that we may consider the following triangle, where $\ell \geq k$.

$$\begin{array}{ccc} & & \text{Cov}_\bullet^{(\ell)} \\ & \nearrow^{F_{(Z,G,\ell)}} & \downarrow \varpi_k \\ \mathfrak{UD}_d & & \text{Cov}_\bullet^{(k)} \\ & \searrow_{F_{(Z,G,k)}} & \end{array} \quad (5.11)$$

Lemma 5.24 *There is a natural transformation $\tau: \varpi_k \circ F_{(Z,G,\ell)} \Rightarrow F_{(Z,G,k)}$ filling the triangle (5.11).*

Proof. The functor going clockwise around (5.11) sends the object (M, A) of \mathfrak{UD}_d to the space $X(M, A)$ together with the quotient

$$\varpi_k \circ \phi_\ell(M, A): \pi_1(X(M, A)) \longrightarrow Q_\ell(M, A) \longrightarrow Q_\ell(M, A)/\Gamma_k(Q_\ell(M, A)), \quad (5.12)$$

where $\phi_\ell(M, A)$ is defined in diagram (5.4). The functor going anticlockwise around (5.11) sends the object (M, A) to the same space $X(M, A)$ together with the quotient

$$\phi_k(M, A): \pi_1(X(M, A)) \longrightarrow Q_k(M, A). \quad (5.13)$$

We recall that a morphism in $\text{Cov}_\bullet^{(k)}$ is a based map of spaces f having the property that $\pi_1(f)$ descends to a homomorphism between the respective quotients. We define the natural transformation $\tau_{(M,A)}$ to be the identity map $X(M, A) \rightarrow X(M, A)$. To see that this really is a morphism, we have to check that the quotient (5.13) factors through the quotient (5.12). Naturality will then be clear. First, note that the quotient of $\pi_1(X(M, A))$ onto $Q_k(M, A)$ factors through its quotient onto $Q_\ell(M, A)$, by the construction of these quotients in the diagram (5.4), so we have a quotient

$$Q_\ell(M, A) \longrightarrow Q_k(M, A).$$

But its target is nilpotent of class at most k , so the subgroup $\Gamma_k(Q_\ell(M, A))$ must be sent to zero under this quotient. Hence it factors further through the quotient onto $Q_\ell(M, A)/\Gamma_k(Q_\ell(M, A))$, as required. \square

Putting together the diagrams (5.11) and (5.10) (with its top row restricted to the subcategories $(-)^{(\ell)}$), we obtain:

$$\begin{array}{ccccccc} & & \text{Cov}_\bullet^{(\ell)} & \xrightarrow{\text{Lift}} & \text{Top}_\bullet^{(\ell)} & \xrightarrow{H_i} & \text{Mod}_\bullet^{(\ell)} \\ & \nearrow^{F_{(Z,G,\ell)}} & \downarrow \varpi_k & \Downarrow & \downarrow \varpi_k & \Downarrow & \downarrow \varpi_k \\ \mathfrak{UD}_d & & \text{Cov}_\bullet^{(k)} & \xrightarrow{\text{Lift}} & \text{Top}_\bullet^{(k)} & \xrightarrow{H_i} & \text{Mod}_\bullet^{(k)} \\ & \searrow_{F_{(Z,G,k)}} & & & & & \end{array} \quad (5.14)$$

Composing these functors and pasting together the natural transformations, this gives us a tower of homological representations $L_i(F_{(Z,G,\ell)})$ of the category \mathfrak{UD}_d :

$$\begin{array}{ccc}
& & \downarrow \varpi_\ell \\
& & \text{Mod}_\bullet^{(\ell)} \\
& & \downarrow \varpi_{\ell-1} \\
& & \vdots \\
& & \downarrow \varpi_2 \\
& & \text{Mod}_\bullet^{(2)} \\
& & \downarrow \varpi_1 \\
& & \text{Mod}_\bullet^{(1)} \\
& & \downarrow \varpi_0 \\
\mathfrak{UD}_d & \begin{array}{l} \nearrow \\ \nearrow \\ \nearrow \\ \longrightarrow \end{array} & \text{Mod}_\bullet^{(0)}
\end{array} \tag{5.15}$$

We emphasise that this diagram does not commute. Instead, the natural transformations of (5.14) paste together to give a natural transformation going “downwards” filling each of the triangles in (5.15).

Definition 5.25 The tower (5.15) of homological representations of \mathfrak{UD}_d is called the *pro-nilpotent homological representation* of \mathfrak{UD}_d associated to the input data (Z, G) .

To justify this name, note that, if we restrict to a single object (M, A) of \mathfrak{UD}_d , we obtain a tower of (possibly twisted) representations of the mapping class group $\pi_0(\text{Diff}_{\text{dec}}(M, A))$ defined over the tower of groups $Q_\bullet(M, A)$. Since the group $Q_\ell(M, A)$ has nilpotency class at most ℓ by construction, the inverse limit of this tower of groups is a pro-nilpotent group.

Recall from §5.1.5 above that if the functor $F_{(Z, G, \ell)}$ is Q -stable on a subcategory $\langle \mathcal{G}, \mathcal{M} \rangle \subseteq \mathfrak{UD}_d$ (in the sense of Definition 5.14), then the homological representation $L_i(F_{(Z, G, \ell)})$ restricted to $\langle \mathcal{G}, \mathcal{M} \rangle$ takes values in a subcategory of Mod_\bullet of the form $\text{Mod}_{Q_\ell}^{\text{tw}}$ for some fixed group Q_ℓ . Hence also in this case we have a tower of (possibly twisted) representations over a tower of groups Q_\bullet whose inverse limit is a pro-nilpotent group.

In many specific examples, the tower of groups Q_\bullet actually stops at a finite stage, so we only have finitely many different representations in the corresponding tower. However, there are *also* many examples in which the tower of groups Q_\bullet does *not* stop, and hence we have a tower of representations that becomes richer at every stage. The question of whether and when the tower Q_\bullet stops is investigated in many important examples related to (surface) braid groups, mapping class groups and loop braid groups in the article [DPS21] by Darné and the authors.

Remark 5.26 In all of the discussion of this section, we may replace the ordinary twisted homology functor H_i with Borel-Moore twisted homology H_i^{BM} , and everything goes through in exactly the same way. In this case, of course, we also have to restrict the category Top_\bullet to its subcategory of proper maps.

5.2. Applications for motion groups

We apply the general construction of §5.1 to families of motion groups (*cf.* Definition 4.22) of type $\{\text{Mot}_{Y_n}(M) \mid n \in \mathbb{N}\}$ where M is a manifold of dimension $d = 2$ or $d = 3$ and Y_n is a closed submanifold of M . For that purpose, we consider the restriction of the continuous functor $\hat{F}_{(Z, G, \ell)}: \mathfrak{UD}_d \rightarrow \text{Cov}_\bullet$ of Proposition 5.8 to the appropriate full subcategory of the form $\langle \mathcal{G}, \mathcal{M} \rangle$ which automorphism groups correspond to the family of motion groups: these are given by Lemma

3.7 applied to the examples of subgroupoids \mathcal{G} and \mathcal{M} described in §4. Moreover we will make some particular choices for the type of submanifold Z and the group G :

- $Z = \underline{k}$ is either a finite set of points of size $k \geq 1$ in \mathbb{R}^d or Z is an unlink $\underline{k}\mathbb{S}^1$ with k components in \mathbb{R}^3 ;
- $G = \mathfrak{S}_k$ (when $Z = k$) or $G = \text{Diff}(\underline{k}\mathbb{S}^1)$ or $\text{Diff}^+(\underline{k}\mathbb{S}^1)$ (when $Z = \underline{k}\mathbb{S}^1$).

Furthermore, the possibilities on the parameter ℓ on the lower central series degree will generally be restricted to $\ell \leq 3$, because the lower central series of the considered motion groups generally stop at I_2 or I_3 . Using the framework and tools of §2, we take \mathbb{Z} as ground ring \mathbb{A} and V to be the trivial fibrewise tensor product functor $V_{\hat{F}(Z,G,\ell)} : \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \text{Mod}_\bullet$. Then, by Lemma 2.20, we define the homological functors for all $i \geq 1$:

$$L_i(\hat{F}_{(Z,G,\ell)}): \pi_0 \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \text{Mod}_\bullet. \quad (5.16)$$

For $\ell = 0$, we obtain the representations corresponding to the action on the homology groups of the universal covers of the spaces $X(M, Y_n)$, and we recover the natural action of the motion groups on the homology groups of the spaces $X(M, Y_n)$ for $\ell = 1$; cf. Remark 5.6. For $\ell = 2$ or $\ell = 3$, as a consequence of some abelianisation computations, we will typically prove that $Q_{(Z,G,\ell)}(X(M, A_n))$ does not depend on n for $n \geq \mu_{(Z,G,\ell)}$, and denote this quotient of by $Q_{(Z,G,\ell)}$. In this case we deduce that:

Proposition 5.27 *For all $i \geq 0$, trivialising the assignment for $n \leq \mu_{(Z,G,\ell)}$, the functor (5.16) defines homological functors*

$$L_i(\hat{F}_{(Z,G,2)}): \pi_0 \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \text{Mod}_{\mathbb{Z}[Q_{(Z,G,2)}]} \quad \text{and} \quad L_i(\hat{F}_{(Z,G,3)}): \pi_0 \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \text{Mod}_{\mathbb{Z}[Q_{(Z,G,3)}]}^{\text{tw}}.$$

Proof. Let $\langle \mathcal{G}, \mathcal{M} \rangle_{\geq \mu_{(Z,G,\ell)}}$ be the full subcategory of $\langle \mathcal{G}, \mathcal{M} \rangle$ on all objects *except* those indexed by non-negative integers strictly less than $\mu_{(Z,G,\ell)}$. By Proposition 5.16, the functor (5.16) induces functors $L_i(\hat{F}_{(Z,G,2)}): \pi_0 \langle \mathcal{G}, \mathcal{M} \rangle_{\geq \mu_{(Z,G,2)}} \rightarrow \text{Mod}_{\mathbb{Z}[Q_{(Z,G,2)}]}$ and $L_i(\hat{F}_{(Z,G,3)}): \pi_0 \langle \mathcal{G}, \mathcal{M} \rangle_{\geq \mu_{(Z,G,3)}} \rightarrow \text{Mod}_{\mathbb{Z}[Q_{(Z,G,3)}]}^{\text{tw}}$. By Lemma 5.17, we extend them to functors over $\langle \mathcal{G}, \mathcal{M} \rangle$ by sending the first objects to the trivial group. \square

General alternatives. For all the constructions presented in §§5.2.1–5.2.3, we could define interesting alternatives by taking $G = 0_{\text{Grp}}$ the trivial group and define $L_i(\hat{F}_{(Z,0_{\text{Grp}},2)}): \pi_0 \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \text{Mod}_\bullet$ and $L_i(\hat{F}_{(Z,0_{\text{Grp}},k)}): \pi_0 \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \text{Mod}_\bullet$ for each $i \geq 0$ and $k \geq 1$. This corresponds to taking *ordered* configuration spaces for the parameter $X(M, A)$ of §5.1.1 instead of *unordered* configuration spaces as done in §§5.2.1–5.2.3. More generally, interpolating the parameter G between 0_{Grp} and $\text{Diff}(Z)$, one may in particular take it such that we consider any *partitioned* configurations. The associated lower central series of the corresponding fundamental group is more complicated generally speaking and does not necessarily stop and therefore more sophisticated representations may arise from these alternatives.

5.2.1. Classical braid groups.

We consider the restriction of the continuous functor $\hat{F}_{(Z,G,\ell)}: \mathcal{UD}_2 \rightarrow \text{Cov}_\bullet$ of Proposition 5.8 to the full subcategory $\mathcal{UB}^{\mathbb{D}^2}$. In particular, we only have $M = \mathbb{D}^2$ the unit 2-disc and $Y_n = \underline{n}$ a set of $n \geq 0$ distinct points in the interior. Moreover, we restrict to $Z = \underline{k}$ a set of $k \geq 1$ distinct points seen as a closed submanifold of \mathbb{R}^2 and take $G = \mathfrak{S}_k$. Hence we define the homological functor

$$L_i(\hat{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}): \mathcal{UB} \rightarrow \text{Mod}_\bullet, \quad (5.17)$$

associated with non-negative integers $i \geq 1$, $k \geq 1$, $\ell \geq 0$ and with groups $Q_{(\underline{k}, \mathfrak{S}_k, \ell)}(\mathbb{D}_n)$. We recall from [DPS21, §4] that:

- $\Gamma_2(\mathbf{B}_{k,n}) = \Gamma_3(\mathbf{B}_{k,n})$ for $k \in \mathbb{N} \setminus \{2\}$ and $n \geq 3$;
- $\Gamma_i(\mathbf{B}_{2,n}) \neq \Gamma_{i+1}(\mathbf{B}_{2,n})$ and $\Gamma_i(\mathbf{B}_{2,n}) \neq \Gamma_{i+1}(\mathbf{B}_{2,n})$ for all $i \geq 1$.

Therefore, if $k \neq 2$, the construction produced for each $\ell > 2$ is equivalent to the one for $\ell = 2$, which is detailed in §5.2.1.1 below. In contrast, if $k = 2$ each $\ell \geq 2$ provides a brand new construction; cf. §5.2.1.2.

5.2.1.1. Standard situation of the abelianisation.

As a direct consequence of the computation for the abelianisations of partitioned braid groups of [DPS21, §4], we have that for all $n \geq 2$, $Q_{(\underline{k}, \mathfrak{S}_k, 2)}(\mathbb{D}_n) = \mathbb{Z}$ if $k = 1$ and \mathbb{Z}^2 if $k \geq 2$. We denote these quotients of by Q_k and deduce from Proposition 5.27:

Proposition 5.28 *For each $i \geq 0$ and $k \geq 1$, the functor (5.17) for $\ell = 2$ defines a homological functor $L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, 2)}): \mathfrak{UB} \rightarrow \text{Mod}_{\mathbb{Z}[Q_k]}$ trivialising the assignment for $n \leq 1$.*

Alternative topological interpretation. The representations encoded by the functor of Proposition 5.28 may alternatively be introduced with a more geometrical point of view. We fix a non-negative integer $n \geq 1$ and recall that γ_2 generically denotes the abelianisation map of a group. For $k = 1$, we consider the composite $\chi \circ \gamma_2: \mathbf{B}_1(\mathbb{D}_n) \rightarrow \mathbb{Z}^n \rightarrow \mathbb{Z}$, where χ denotes the sum map. For $k \geq 2$, first we consider the inclusion map $i: C_k(\mathbb{D}_n) \hookrightarrow C_k(\mathbb{D}^2)$ induced by forgetting the marked points of \mathbb{D}_n , and denote by i_* the induced surjective homomorphism on π_1 . Let \mathcal{T} be the composite $\gamma_2 \circ i_*: \mathbf{B}_k(\mathbb{D}_n) \rightarrow \mathbf{B}_k \rightarrow \mathbb{Z}$. We may have the following geometrical interpretation of this morphism: for $\lambda \in \pi_1(C_k(\mathbb{D}_n))$ and c_λ a simple closed curve in $C_k(\mathbb{D}_n)$ representative of λ , one can think of $\mathcal{T}(\lambda)$ as counting the total number of half-twists (i.e. half a Dehn twist in a tubular neighbourhood along the path c_λ) that occur in the path c_λ of the k configuration points in \mathbb{D}^2 . Furthermore, let j be the composite $C_k(\mathbb{D}_n) \hookrightarrow C_{k,n}(\mathbb{D}^2) \hookrightarrow C_{k+n}(\mathbb{D}^2)$ where the first map is the canonical fiber inclusion of (4.4) and the second map is induced by the inclusion of symmetric groups $\mathfrak{S}_k \times \mathfrak{S}_n \hookrightarrow \mathfrak{S}_{k+n}$, and denote by j_* the induced injective homomorphism on π_1 . For $\lambda \in \pi_1(C_k(\mathbb{D}_n))$ and c_λ as above, the composite $\gamma_2 \circ j_*(\lambda)$ can geometrically be interpreted as counting the total number of half-twists that occur in the path c_λ , and also between configuration points and the n marked points. Hence $\gamma_2 \circ j_*(\lambda) - \mathcal{T}(\lambda)$ is twice the total number of times that the configuration points wind around the marked points: it thus corresponds to twice the total winding number of c_λ . Hence we define $\mathcal{W}: \mathbf{B}_k(\mathbb{D}_n) \rightarrow \mathbb{Z}$ to be the surjective morphism defined by the total winding number, i.e. $\frac{1}{2}(\gamma_2 \circ j_* - \mathcal{T})$. These descriptions come from [Bud05, §2]. Finally, denoting by Δ the diagonal morphism, we consider the product $(\mathcal{W} \times \mathcal{T}) \circ \Delta: \mathbf{B}_k(\mathbb{D}_n) \rightarrow \mathbb{Z}^2$, defined by $\lambda \mapsto (\mathcal{W}(\lambda), \mathcal{T}(\lambda))$, for the choice of the quotient of the fundamental group.

It is straightforward computation to show that the morphisms $(\mathcal{W} \times \mathcal{T}) \circ \Delta$ and $\phi_{(\underline{k}, \mathfrak{S}_k, 2)}(\mathbb{D}_n): \mathbf{B}_k(\mathbb{D}_n) \rightarrow \mathbb{Z}^2$ (defining the functor $\mathring{F}_{(\underline{k}, \mathfrak{S}_k, 2)}: \mathfrak{UB}r_{\geq 1}^{\mathbb{D}^2} \rightarrow \text{Cov}_{Q_k}$) are equal. Since $C_k(\mathbb{D}_n)$ is a path-connected, locally path-connected and semi-locally simply-connected space, it follows from covering space theory (see, for example, [Hat02, Theorem 1.38]) that

Proposition 5.29 *The representation of \mathbf{B}_n encoded by the functor $L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, 2)})$ of Proposition 5.28 is equivalent to the action on the homology groups of associated with the quotient $\chi \circ \gamma_2$ if $k = 1$ and $(\mathcal{W} \times \mathcal{T}) \circ \Delta$ if $k \geq 2$.*

Bigelow's construction and Lawrence-Bigelow representations. Bigelow [Big04] introduced a general method to construct a representation of the braid group \mathbf{B}_n from a representation of the braid group \mathbf{B}_k for two integers k and n . It builds the well-known families of *Lawrence-Bigelow representations*, originally introduced by Lawrence [Law90] as representations of Hecke algebras. The most famous families among them are the Burau representations, originally introduced in [Bur35], and the *Lawrence-Krammer-Bigelow* representations that Bigelow [Big01] and Krammer [Kra02] independently proved to be faithful. We will prove that the Lawrence-Bigelow representations are actually recovered by the constructions of §5.2.1.1 (see Theorem 5.32).

We first reformulate Bigelow's construction with the framework and tools of §2. We fix an associative, unital ring R and $k \geq 1$ a non-negative integer. We consider the following ingredients.

- Let $\text{Big}: \mathfrak{UB}r^{\mathbb{D}^2} \rightarrow \bullet\text{Cov}\bullet$ be the functor that sends n to the k -th unordered configuration space $C_k(\mathbb{D}_n)$ together with the jointly surjective pair of quotients $\mathcal{W}: \mathbf{B}_k(\mathbb{D}_n) \rightarrow \mathbb{Z}$ the total winding number and $\mathcal{T}: \mathbf{B}_k(\mathbb{D}_n) \rightarrow \mathbf{B}_k$ the homomorphism that forgets the marked points. A morphism of $\mathfrak{B}r^{\mathbb{D}^2}$ is the embedding space $\mathfrak{UB}r^{\mathbb{D}^2}(\mathbb{D}_m, \mathbb{D}_n) \cong \text{Emb}_{\text{dec}}^{\text{Diff}}(\mathbb{D}_m, \mathbb{D}_n)$ for $m \leq n$ by Proposition 4.8, which induces a map of configuration spaces: one straightforwardly checks

from the definitions that this embedding commutes with the quotients of \mathcal{W} and \mathcal{T} , so it preserves their kernels, making F a valid functor into $\bullet\text{Cov}\bullet$.

- Let V be a left $R[\mathbf{B}_k]$ -module and $\mathbf{V}: \mathfrak{UBr}^{\mathbb{D}^2} \rightarrow \bullet\text{Mod}\bullet$ be the constant functor at the object $(R[\mathbf{B}_k], R, V)$.

Using Definition 2.21, we recover Bigelow's construction [Big04, §2]:

Definition 5.30 The functor $L_k(\text{Big}; \mathbf{V}): \mathfrak{UB} \rightarrow \bullet\text{Mod}\bullet$ encodes the construction of [Big04, §2]: for each $n \geq 1$, the n -th Bigelow construction is the representation of \mathbf{B}_n on the $R[\mathbb{Z}]$ -module $L_k(\text{Big}; \mathbf{V})(n)$.

Assigning $R = \mathbb{Z}$ and $V = \mathbb{Z}[\mathbb{Z}]$ the \mathbf{B}_k -representation where \mathbf{B}_k acts on \mathbb{Z} through the abelianisation and left-multiplication, the functor $L_k(\text{Big}; \mathbb{Z}[\mathbb{Z}])$ encodes the k -th Lawrence-Bigelow representations. In particular, the representations encoded in $L_1(\text{Big}; \mathbb{Z}[\mathbb{Z}])$ are the reduced Burau representations, and those encoded in $L_2(\text{Big}; \mathbb{Z}[\mathbb{Z}])$ are the Lawrence-Krammer-Bigelow representations; see [Big03; Big01]. In addition, by [PP02, Theorem 1.2], the tensor product with the field of fractions $L_2(\text{Big}; \mathbb{Z}[\mathbb{Z}]) \otimes_{\mathbb{Z}[\mathbb{Z}]} \mathbb{Q}(\mathbb{Z}^2)$ is isomorphic to the Lawrence-Krammer representation [Kra02] tensored by the field of fractions $\mathbb{Q}(\mathbb{Z}^2)$.

Remark 5.31 In Bigelow's setting [Big04, §2], the ground ring R is \mathbb{C} and a choice of non-zero complex number q is fixed. Then $\mathbb{C} \otimes_{\mathbb{C}[\mathbb{Z}]} L_m(\text{Big}; M)(n)$ is exactly the result of Bigelow's construction, with \mathbb{C} viewed as a right $\mathbb{C}[\mathbb{Z}]$ -module by letting $1 \in \mathbb{Z}$ act by multiplication by q .

Theorem 5.32 For each $k \geq 1$, the functor $L_k(\mathring{F}_{(k, \mathfrak{S}_k, 2)})$ of §5.2.1.1 is isomorphic to the functor $L_k(\text{Big}; \mathbb{Z}[\mathbb{Z}])$. Hence we denote this functor by $\mathfrak{LB}_k: \mathfrak{UB} \rightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Q}_k]}$ and call it the k -th Lawrence-Bigelow functor.

Proof. Recalling the result from Proposition 5.29, we then deduce from Shapiro's lemma an abelian group isomorphism for each n

$$H_k(C_k(\mathbb{D}_n)^{\mathcal{W}}; \mathbb{Z}[\mathbb{Z}]) \cong H_k(C_k(\mathbb{D}_n)^{\phi_{(k, \mathfrak{S}_k, 2)}(\mathbb{D}_n)}; \mathbb{Z}).$$

It follows from the definitions of both $L_k(\text{Big}; \mathbb{Z}[\mathbb{Z}])$ and $L_k(\mathring{F}_{(k, \mathfrak{S}_k, 2)})$ that the effect of a morphism σ of \mathfrak{UB} proceeds from the unique lift of an embedding of a marked disc into another marked disc for the associated covering space. The above isomorphism thus provides an isomorphism between the functors $L_k(\text{Big}; \mathbb{Z}[\mathbb{Z}])$ and $L_k(\mathring{F}_{(k, \mathfrak{S}_k, 2)})$. \square

5.2.1.2. Exceptional situation for $k = 2$.

Recall from [DPS21, §4] that $\Gamma_i(\mathbf{B}_{2,n}) \neq \Gamma_{i+1}(\mathbf{B}_{2,n})$ for all $i \geq 1$. Applying the procedure of §5.1 thus provides a specific global homological representation for each $\ell \geq 2$. In particular, additional properties may be deduced from the study of §5.1.7.

We fix some $\ell \geq 3$ and $n \geq 2$. We consider the variant of the global homological representation construction using Borel-Moore homology and we restrict to taking the second homology group. The adaptation of the diagram (5.14) using Borel-Moore homology (cf. Remark 5.26) defines a natural transformation $\eta_{\ell, n}: L_2^{BM}(F_{(2, \mathfrak{S}_2, \ell)})(n) \rightarrow L_2^{BM}(F_{(2, \mathfrak{S}_2, 2)})(n)$.

A general property of [Big04, Lemma 3.1] describes the Borel-Moore homology of a configuration space of points in the punctured disc and thus provides the isomorphisms

$$L_2^{BM}(F_{(2, \mathfrak{S}_2, \ell)})(n) \cong \mathbb{Z}[Q_{\ell, n}]^{\oplus n(n-1)/2} \quad \text{and} \quad L_2^{BM}(F_{(2, \mathfrak{S}_2, 2)})(n) \cong \mathbb{Z}[\mathbb{Z}^2]^{\oplus n(n-1)/2}.$$

Therefore, the natural transformation $\eta_{\ell, n}$ is thus a \mathbf{B}_n -equivariant surjection $L_2^{BM}(F_{(2, \mathfrak{S}_2, \ell)})(n) \rightarrow L_2^{BM}(F_{(2, \mathfrak{S}_2, 2)})(n)$ induced by the canonical group ring surjection $\mathbb{Z}[Q_{\ell, n}] \rightarrow \mathbb{Z}[\mathbb{Z}^2]$.

Faithfulness results. Note the following general result on the quotient of a representation.

Lemma 5.33 Let R be a ring and V be a R -module. We consider (ρ, V) a representation of a group G . If any quotient (μ, W) of (ρ, V) is a faithful representation of G , then (ρ, V) is also faithful.

Proof. We denote by $\alpha : (\rho, V) \rightarrow (\mu, W)$ the quotient G -equivariant map. Let g be an element of G such that $\rho(g) = \text{id}_V$. Then $\mu(g) \circ \alpha = \alpha$ and therefore $\mu(g) = \text{id}_W$ since α is an epimorphism. We deduce from the faithfulness of (μ, W) that $g = \text{id}_G$, which ends the proof. \square

Recall from [Big02, §4] that for each n the \mathbf{B}_n -representation $L_2^{BM}(F_{(2, \mathfrak{S}_{2,2})})(n)$ is faithful.

Corollary 5.34 *For each $\ell \geq 3$, the functor $L_2^{BM}(F_{(2, \mathfrak{S}_{2,\ell})})$ encodes a faithful linear representation of \mathbf{B}_n for each n .*

5.2.2. Surface braid groups.

Let S be a compact, connected, smooth surface with boundary different from the 2-disc: it is therefore isomorphic either to $\Sigma_{g,1}$ or else to $\mathcal{N}_{h,1}$ for some fixed non-negative integers $g \geq 1$ and $h \geq 1$. We consider the restriction of the continuous functor $\mathring{F}_{(Z,G,\ell)} : \mathfrak{UD}_2 \rightarrow \text{Cov}_\bullet$ of Proposition 5.8 to the full subcategory $\langle \mathcal{B}r^{\mathbb{D}^2}, \mathcal{B}r^S \rangle$. In particular, we only have $M = S$ and $Y_n = \underline{n}$ a set of n distinct points in the interior. Moreover, we take $Z = \underline{k}$ a set of $k \geq 1$ distinct points seen as a closed submanifold of \mathbb{R}^2 and set $G = \mathfrak{S}_k$. Hence we consider an homological functor

$$L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}): \langle \beta, \beta^S \rangle \longrightarrow \text{Mod}_\bullet \quad (5.18)$$

associated with non-negative integers $i \geq 1$, $\ell \geq 0$ and $k \geq 1$, and with groups $Q_{(\underline{k}, \mathfrak{S}_k, \ell)}(S^{(n)})$. We recall from [DPS21, §6] that for any $g \geq 1$, $h \geq 1$ and $n \geq 3$, for $S = \Sigma_{g,1}$ or $\mathcal{N}_{h,1}$:

- $\Gamma_2(\mathbf{B}_{k,n}(S)) \neq \Gamma_3(\mathbf{B}_{k,n}(S)) = \Gamma_4(\mathbf{B}_{k,n}(S))$ if $k \geq 3$;
- $\Gamma_\ell(\mathbf{B}_{1,n}(S)) \neq \Gamma_{\ell+1}(\mathbf{B}_{1,n}(S))$ and $\Gamma_\ell(\mathbf{B}_{2,n}(S)) \neq \Gamma_{\ell+1}(\mathbf{B}_{2,n}(S))$ for all $\ell \geq 1$.

Hence we study the construction for $\ell = 2$ in §5.2.2.1 in all the situations and then we consider the further situation of $\ell = 3$ for orientable surfaces in §5.2.2.2. As in §5.2.1.2, the procedure for the particular parameter $k = 1$ and 2 of §5.1 provides a specific representation for each $\ell \geq 3$, but will not be addressed in details here.

5.2.2.1. Standard situation of the abelianisation.

We deduce from the computation for the abelianisations of partitioned surface braid groups of [DPS21, §6] that for all $k \geq 1$, if $n \geq 2$ then $Q_{(\underline{k}, \mathfrak{S}_k, 2)}(S^{(n)}) \cong \mathbb{Z}^{p_S} \oplus (\mathbb{Z}/2\mathbb{Z})^{d_k}$ where

$$p_S = \begin{cases} 2g & \text{if } S = \Sigma_{g,1}, \\ h & \text{if } S = \mathcal{N}_{h,1}, \end{cases} \text{ and } d_k = \begin{cases} 0 & \text{if } k = 1, \\ 1 & \text{if } k \geq 2. \end{cases}$$

We denote this quotient by $Q_{(k,2)}(S)$ and deduce from Proposition 5.27 that:

Proposition 5.35 *For each $i \geq 0$ and $k \geq 1$, the functor (5.18) for $\ell = 2$ defines a homological functor $L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, 2)}): \langle \beta, \beta^S \rangle \rightarrow \text{Mod}_{\mathbb{Z}[Q_{(k,2)}(S)]}$ trivialising the assignment for $n \leq 1$.*

5.2.2.2. Situation of $\ell = 3$.

First of all, we note the following result for the orientable surfaces $\Sigma_{g,1}$ for $g \geq 1$.

Lemma 5.36 *For all $k \geq 1$, if $n \geq 3$ then $Q_{(\underline{k}, \mathfrak{S}_k, 3)}(\Sigma_{g,1}^{(n)}) = Q_{(\underline{k}, \mathfrak{S}_k, 3)}(\Sigma_{g,1}^{(n+1)})$. For each k , we denote the common quotient by $Q_{(k,3)}(\Sigma_{g,1})$.*

Proof. We fix $k \geq 0$ and $n \geq 3$. The full presentation of the group $\mathbf{B}_{k,n}(\Sigma_{g,1}) \cong \mathbf{B}_k(\Sigma_{g,1}^{(n)}) \rtimes \mathbf{B}_n(\Sigma_{g,1})$ is detailed in [BGG17, Proposition 3.2] and is used for this proof. Note that the elements that depend on n in its presentations are the set of braid generators $\{\sigma_1, \dots, \sigma_{n-1}\}$ of $\mathbf{B}_n(\Sigma_{g,1})$ and the set of the generators $\{\xi_1, \dots, \xi_n\}$ of $\mathbf{B}_k(\Sigma_{g,1}^{(n)})$. Because the commutativity of (5.4), we abuse the notation γ_3 to denote the projections on the metabelian quotients.

First, since $\gamma_3([\sigma_i, [\sigma_{i+1}, \sigma_i]]) = 1$, the braid relation $\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$ implies that $\gamma_3(\sigma_i) = \gamma_3(\sigma_{i+1})$ for all $i \in \{1, \dots, n-2\}$. Moreover, from the relations in the presentation of [BGG17, Proposition 3.2, relations (c.3)], we have the following relations: $\sigma_i \xi_i \sigma_i^{-1} = \xi_i^{-1} \xi_{i+1} \xi_i$, $\sigma_i \xi_{i+1} \sigma_i^{-1} = \xi_i$ and $\sigma_i \xi_j \sigma_i^{-1} = \xi_j$ if $j \neq i, i+1$. Then, since $\gamma_3(\sigma_i) = \gamma_3(\sigma_{i+1}) = \gamma_3(\sigma_{i+2})$, we deduce that $\gamma_3(\xi_{i+1}) = \gamma_3(\sigma_i^{-1} \xi_i \sigma_i) = \gamma_3(\sigma_{i+2}^{-1} \xi_i \sigma_{i+2}) = \gamma_3(\xi_i)$ for all $i \in \{1, \dots, n-2\}$. Hence the presentation of $\mathbf{B}_{k,n}(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_{k,n}(\Sigma_{g,1}))$ is independent of n . In addition, the above proof for $k=0$ shows that $\mathbf{B}_n(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_n(\Sigma_{g,1}))$ is also independent of n . Hence, for each $k \geq 1$, the canonical morphism $\mathbf{B}_{k,n}(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_{k,n}(\Sigma_{g,1})) \twoheadrightarrow \mathbf{B}_n(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_n(\Sigma_{g,1}))$ is independent of n , hence so is the kernel $Q_{(\underline{k}, \mathfrak{S}_k)}(\Sigma_{g,1}^{(n)})$. \square

For $k, n \geq 3$, explicit calculations of $\mathbf{B}_{k,n}(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_{k,n}(\Sigma_{g,1}))$ and $\mathbf{B}_n(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_n(\Sigma_{g,1}))$ are provided by [BGG17, Corollaries 3.9(i) and 3.14(i)]. In particular we deduce that $Q_{(k,3)}(\Sigma_{g,1})$ is equal to $\mathbb{Z} \times (\mathbb{Z}^{g+1} \rtimes \mathbb{Z}^g)$ for any $k \geq 3$. Finally, it follows from Proposition 5.27 that:

Proposition 5.37 *For each $i \geq 0$ and $k \geq 1$, the functor (5.18) for $\ell = 3$ provides the homological functors (trivialising the assignment for $n \leq 2$)*

- $L_i(\hat{F}_{(\underline{k}, \mathfrak{S}_k, 3)}): \langle \beta, \beta^{\Sigma_{g,1}} \rangle \rightarrow \text{Mod}_{\mathbb{Z}[Q_{(k,3)}(\Sigma_{g,1})]}^{\text{tw}}$ for each $g \geq 1$.
- $L_i(\hat{F}_{(\underline{k}, \mathfrak{S}_k, 3)}): \langle \beta, \beta^{\mathcal{N}_{h,1}} \rangle \rightarrow \text{Mod}_{\bullet}$ for each $h \geq 1$.

Conjecture 5.38 For each $g \geq 1$, we conjecture that $\mathbf{B}_{1,n}(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_{1,n}(\Sigma_{g,1})) \cong (\mathbb{Z}^2 \times \mathbb{Z}^{2g}) \rtimes \mathbb{Z}^{2g}$ and $\mathbf{B}_{2,n}(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_{2,n}(\Sigma_{g,1})) \cong (\mathbb{Z}^3 \times \mathbb{Z}^{2g}) \rtimes \mathbb{Z}^{2g}$ for all $n \geq 3$. Therefore $Q_{(1,3)}(\Sigma_{g,1})$ would be equal to \mathbb{Z}^{2g+1} and $Q_{(2,3)}(\Sigma_{g,1})$ would be equal to $\mathbb{Z} \times (\mathbb{Z}^{g+1} \rtimes \mathbb{Z}^g)$.

For each $h \geq 1$, we conjecture that $\mathbf{B}_{k,n}(\mathcal{N}_{h,1})/\Gamma_3(\mathbf{B}_{k,n}(\mathcal{N}_{h,1})) \cong \mathbb{Z}/2\mathbb{Z} \times (\mathbb{Z} \times \mathbb{Z}^h) \rtimes \mathbb{Z}^h$ if $k=1$ and $(\mathbb{Z}/2\mathbb{Z})^2 \times (\mathbb{Z} \times \mathbb{Z}^h) \rtimes \mathbb{Z}^h$ if $k \geq 2$ for all $n \geq 3$. Then the analogue of Lemma 5.36 would hold and therefore the target category of the associated homological functor would be $\text{Mod}_{\mathbb{Z}[Q_{(k,3)}(\mathcal{N}_{h,1})]}^{\text{tw}}$ where $Q_{(k,3)}(\mathcal{N}_{h,1}) \cong \mathbb{Z}^{1+h}$ if $k=1$ and $\mathbb{Z}^{1+h} \oplus \mathbb{Z}/2\mathbb{Z}$ if $k \geq 2$.

Remark 5.39 In the orientable case for $k \geq 3$ (and conjecturally in both cases for all $k \geq 1$), there is a \mathbb{Z} summand in the quotient group $Q_{(k,3)}(S)$, generated by (the image of) the surface braid ξ where one of the first k strands winds once around one of the last n strands. In order to obtain *untwisted* representations, in other words a homological functor taking values in a category of the form $\text{Mod}_{\mathbb{Z}[Q]}$ rather than $\text{Mod}_{\mathbb{Z}[Q]}^{\text{tw}}$, we need to take a quotient of $Q_{(k,3)}(S)$ on which the natural action of each $B_n(S)$ (for $n \geq 3$) is trivial. For $n \geq 3$, the coinvariants of the action of $B_n(S)$ on $Q_{(k,3)}(S)$ are given by killing the \mathbb{Z} summand generated by ξ . Thus, if we define $Q'_{(k,3)}(S)$ to be this quotient of $Q_{(k,3)}(S)$, we obtain homological functors

- $\langle \beta, \beta^S \rangle \longrightarrow \text{Mod}_{\mathbb{Z}[Q'_{(k,3)}(S)]}$ for $S = \Sigma_{g,1}$ or $S = \mathcal{N}_{h,1}$ for each $g, h \geq 1$,

where

- $Q'_{(k,3)}(\Sigma_{g,1})$ is \mathbb{Z}^{2g} for $k=1$ and $\mathbb{Z}^{g+1} \rtimes \mathbb{Z}^g$ for $k \geq 2$,
- $Q'_{(k,3)}(\mathcal{N}_{h,1})$ is \mathbb{Z}^h for $k=1$ and $\mathbb{Z}^h \oplus \mathbb{Z}/2\mathbb{Z}$ for $k \geq 2$.

In the orientable case with $k \geq 3$, this statement is unqualified; in the other cases it depends on Conjecture 5.38. These quotients and the corresponding untwisted homological functors will be explained and studied in more detail in the sequel paper [PS21].

The An-Ko representations. The procedure described here is in a sense a reinterpretation following [BGG17] of the work [AK10] to extend some homological representations from the classical braid groups to the surface braid groups. Namely, the functor $L_k(\hat{F}_{(\underline{k}, \mathfrak{S}_k, 3)})$ induces representation of $\mathbf{B}_n(\Sigma_{g,1})$ for any $k \geq 1$ and $n \geq 3$. Furthermore, let $\mathbf{B}_{k,n}(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_{k,n}(\Sigma_{g,1})) \twoheadrightarrow \mathcal{Q}$ be an epimorphism: it automatically induces a $\mathbf{B}_n(\Sigma_{g,1})$ - $Q_{(\underline{k}, \mathfrak{S}_k, 3)}$ -bimodule structure on \mathcal{Q} . We denote by $\psi_n^{\mathcal{Q}}$ the induced action of $\mathbf{B}_n(\Sigma_{g,1})$ on $\mathbb{Z}[\mathcal{Q}]$.

The k -th An-Ko representation of $\mathbf{B}_n(\Sigma_{g,1})$ [AK10, Theorem 3.2] is the tensor product $\psi_n^{\mathcal{Q}} \otimes_{\mathbb{Z}[Q_{(\underline{k}, \mathfrak{S}_k, 3)}]} L_k(\cdot)$ for $\mathcal{Q} = \mathbf{B}_{k,n}(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_{k,n}(\Sigma_{g,1}))$. The groups $Q_{(\underline{k}, \mathfrak{S}_k, 3)}$ and \mathcal{Q} are abstractly defined in [AK10] in terms of group presentation to satisfy certain technical homological constraints: [BGG17, §4] gives all the connections to the third lower central quotient. The general method applied in

§5.2.2 underlines the mainspring of these groups, proving moreover that the use of the third lower central quotient is a key tool to define the homological representations and giving an alternative to the technical result [AK10, Lemma 3.1]. Finally, we refer the reader to [AK10, §3.B, pp. 273-274] for explicit computations for $k = 1, 2, 3$ of the matrices of the representation of $\mathbf{B}_n(\Sigma_{1,1})$.

5.2.3. Loop braid groups.

To apply the construction of §5.1 to extended loop braid groups, we consider restrictions of the continuous functor $\mathring{F}_{(Z,G,\ell)}: \mathcal{UD}_3 \rightarrow \text{Cov}_\bullet$ of Proposition 5.8 to the full subcategory \mathcal{ULB}' of \mathcal{UD}_3 (cf. §4.6), thus assigning $M = \mathbb{D}^3$ and $A = Y_n = \underline{n}\mathbb{S}^1$ a set of n disjoint, unlinked circles in the interior. For non-extended loop braid groups, we instead consider restrictions to $\mathcal{ULB} \subseteq \mathcal{UD}_3^+$ of the analogous continuous functor

$$\mathring{F}_{(Z,G,\ell)}: \mathcal{UD}_3^+ \longrightarrow \text{Cov}_\bullet$$

given by the analogue of Proposition 5.8 for \mathcal{D}_d^+ instead of \mathcal{D}_d . This general construction is exactly parallel to the construction of the non-oriented version, the only difference being that, in diagram (5.4), embeddings of the manifold A are always considered modulo $\text{Diff}^+(A)$, rather than the full diffeomorphism group $\text{Diff}(A)$. The space $X(M, A)$ is therefore the same in this construction, but the quotient $Q(M, A)$ of its fundamental group is generally different (in particular, it will be in our setting of loop braid groups).

Two choices for the submanifold Z naturally arise as relevant parameters to construct homological representations: either we consider a set of points (studied in §5.2.3.1 below), or we take Z to be an unlink (detailed in §5.2.3.2).

5.2.3.1. Using configurations of points.

We take $Z = \underline{k}$ a set of $k \geq 1$ distinct points seen as a closed submanifold of \mathbb{R}^3 and take $G = \mathfrak{S}_k$. We therefore consider the functors $\mathring{F}'_{(\underline{k}, \mathfrak{S}_k, \ell)}: \mathcal{ULB}' \rightarrow \text{Cov}_\bullet$ and $\mathring{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}: \mathcal{ULB} \rightarrow \text{Cov}_\bullet$ associated with a pair of non-negative integers (k, ℓ) where $k \geq 1$ and $\ell \geq 0$. By Lemma 2.20, we have the following homological functors for all $i \geq 1$:

$$L_i(\mathring{F}'_{(\underline{k}, \mathfrak{S}_k, \ell)}): \mathcal{ULB}' \longrightarrow \text{Mod}_\bullet \quad \text{and} \quad L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}): \mathcal{ULB} \longrightarrow \text{Mod}_\bullet. \quad (5.19)$$

We denote by $Q_{(\underline{k}, \mathfrak{S}_k, \ell)}(\mathbb{D}_n^3)$ the quotient group $Q(M, A)$ in diagram (5.4) for $(Z, G, \ell) = (\underline{k}, \mathfrak{S}_k, \ell)$ and recall from [DPS21, §5] that for any $n \geq 4$,

- $\Gamma_2(\pi_1(C_k(\mathbb{D}_n^3)) \rtimes \mathbf{LB}_n) = \Gamma_3(\pi_1(C_k(\mathbb{D}_n^3)) \rtimes \mathbf{LB}_n)$ if $k = 1$ or $k \geq 3$;
- for all $\ell \geq 1$, $\Gamma_\ell(\pi_1(C_2(\mathbb{D}_n^3)) \rtimes \mathbf{LB}_n) \neq \Gamma_{\ell+1}(\pi_1(C_2(\mathbb{D}_n^3)) \rtimes \mathbf{LB}_n)$ and $\Gamma_\ell(\pi_1(C_2(\mathbb{D}_n^3)) \rtimes \mathbf{LB}_n) \neq \Gamma_{\ell+1}(\pi_1(C_2(\mathbb{D}_n^3)) \rtimes \mathbf{LB}'_n)$.

Hence we just focus on the construction for $\ell = 2$ here. Again, the procedure for the particular parameters $k = 2$ of §5.1 provides a specific representation for each $\ell \geq 2$, but will not be addressed in details here.

We deduce from the computation for the abelianisations of partitioned loop braid groups of [DPS21, §5] that for all $n \geq 2$, $Q_{(\underline{k}, \mathfrak{S}_k, 2)}(\mathbb{D}_n^3) = \mathbb{Z}$ if $k = 1$ and $\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ if $k \geq 2$, whereas $Q'_{(\underline{k}, \mathfrak{S}_k, 2)}(\mathbb{D}_n^3) = \mathbb{Z}/2\mathbb{Z}$ if $k = 1$ and $(\mathbb{Z}/2\mathbb{Z})^2$ if $k \geq 2$. We deduce from Proposition 5.27 that:

Proposition 5.40 *For each $i \geq 0$ and $k \geq 1$, trivialising the assignment for $n \leq 1$, the functors (5.19) determine homological representations:*

$$\begin{aligned} \text{(for } k = 1) \quad \mathcal{ULB}' &\longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}/2\mathbb{Z}]} & \text{and} & \quad \mathcal{ULB} \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}]}; \\ \text{(for } k \geq 2) \quad \mathcal{ULB}' &\longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}]} & \text{and} & \quad \mathcal{ULB} \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}]} \end{aligned}$$

Notation 5.41 We denote these homological representations by $L_i(\mathcal{LB}'_{k,2})$ and $L_i(\mathcal{LB}_{k,2})$ respectively. When $i = k$ we will also write them as $\mathfrak{L}_k(k, \mathcal{LB}')$ and $\mathfrak{L}_k(k, \mathcal{LB})$ respectively.

Among the families of representations constructed by Proposition 5.40, the easiest to understand are those corresponding to $i = k = 1$. For instance, we explicitly compute that the matrices of the representations $L_1(\mathcal{LB}_{1,2})(n): \mathbf{LB}_n \rightarrow \text{Aut}_{\mathbb{Z}[\mathbb{Z}]}(\mathbb{Z}[\mathbb{Z}]^{\oplus n-1})$ are those of:

- the reduced Burau representation of the braid group \mathbf{B}_n for the generators $\{\sigma_1, \dots, \sigma_{n-1}\}$;
- the standard representation of the symmetric group \mathfrak{S}_n for the generators $\{\tau_1, \dots, \tau_{n-1}\}$.

We also explicitly compute (see §7) matrices for the representations $L_1(\mathcal{LB}'_{1,2})(n)$ of the extended loop-braid groups \mathbf{LB}'_n over $R = \mathbb{Z}[\mathbb{Z}/2\mathbb{Z}] = \mathbb{Z}[t^{\pm 1}]/(t^2 - 1)$. These are more subtle, since the underlying R -module of the representation is not free: it is $R^{n-1} \oplus R/(t-1)$. On the other hand, the variants of $L_1(\mathcal{LB}_{1,2})(n)$ and $L_1(\mathcal{LB}'_{1,2})(n)$ using *reduced* homology are representations of \mathbf{LB}_n on $\mathbb{Z}[\mathbb{Z}]^n$ (respectively of \mathbf{LB}'_n on R^n), which extend the *unreduced* Burau representations of the classical braid groups. In the case of non-extended loop-braid groups, this extension has already been introduced in [Ver01, §4] and [BS20, §1.3.1]. See §7 for full details.

5.2.3.2. Using configurations of unlinks.

We now consider $Z = \underline{k}\mathbb{S}^1$ a set of $k \geq 1$ distinct unlinks seen as a closed submanifold of \mathbb{R}^3 . However, we essentially have two choices for the groups G : fixing an orientation on each unlink, we can assume that the considered maps preserve or not these orientations.

5.2.3.3. For configurations of oriented unlinks. We take $G = \text{Diff}^+(\underline{k}\mathbb{S}^1)$. We consider the homological functors

$$L_i(\mathring{F}'_{(\underline{k}\mathbb{S}^1, \text{Diff}^+(\underline{k}\mathbb{S}^1), \ell)}): \mathcal{ULB}' \longrightarrow \text{Mod}_{\bullet} \quad \text{and} \quad L_i(\mathring{F}_{(\underline{k}\mathbb{S}^1, \text{Diff}^+(\underline{k}\mathbb{S}^1), \ell)}): \mathcal{ULB} \longrightarrow \text{Mod}_{\bullet} \quad (5.20)$$

associated with non-negative integers $i \geq 1$, $\ell \geq 0$ and $k \geq 1$. We recall from [DPS21, §5] that for any $n \geq 4$,

- for $k \in \mathbb{N} \setminus \{2, 3\}$: $\Gamma_2(\mathbf{LB}_{k,n}) = \Gamma_3(\mathbf{LB}_{k,n})$ and $\Gamma_2(\pi_1(U_k^+(\mathbb{D}_n^3)) \rtimes \mathbf{LB}'_n) = \Gamma_3(\pi_1(U_k^+(\mathbb{D}_n^3)) \rtimes \mathbf{LB}'_n)$;
- for $k \in \{2, 3\}$: $\Gamma_\ell(\mathbf{LB}_{k,n}) \neq \Gamma_{\ell+1}(\mathbf{LB}_{k,n})$ and $\Gamma_\ell(\pi_1(U_k^+(\mathbb{D}_n^3)) \rtimes \mathbf{LB}'_n) \neq \Gamma_{\ell+1}(\pi_1(U_k^+(\mathbb{D}_n^3)) \rtimes \mathbf{LB}'_n)$ for all $\ell \geq 1$.

$\Gamma_2(\mathbf{LB}_{k,n}) = \Gamma_3(\mathbf{LB}_{k,n})$ and $\Gamma_2(\pi_1(U_k^+(\mathbb{D}_n^3)) \rtimes \mathbf{LB}'_n) = \Gamma_3(\pi_1(U_k^+(\mathbb{D}_n^3)) \rtimes \mathbf{LB}'_n)$. Therefore, we detail below the construction produced for $\ell = 2$. Similarly to §5.2.1.2, the procedure for the particular parameters $k = 2$ or 3 of §5.1 may provide a specific representation for each $\ell \geq 2$, but will not be addressed in details here.

It follows from the abelianisations computations of [DPS21, §5] that for $n \geq 2$:

- $Q'_{(\underline{k}\mathbb{S}^1, \text{Diff}^+(\underline{k}\mathbb{S}^1), 2)}(\mathbb{D}_n^3) = \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$ if $k = 1$ and $\mathbb{Z}^2 \oplus (\mathbb{Z}/2\mathbb{Z})^2$ if $k \geq 2$;
- $Q_{(\underline{k}\mathbb{S}^1, \text{Diff}^+(\underline{k}\mathbb{S}^1), 2)}(\mathbb{D}_n^3) = \mathbb{Z}^2$ if $k = 1$ and $\mathbb{Z}^3 \oplus \mathbb{Z}/2\mathbb{Z}$ if $k \geq 2$.

Then it follows from Proposition 5.27 that:

Proposition 5.42 *For each $i \geq 0$ and $k \geq 1$, trivialising the assignment for $n \leq 1$, the functors (5.20) for $\ell = 2$ define homological functors:*

$$\begin{aligned} (\text{for } k = 1) \quad & \mathcal{ULB}' \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}]} \quad \text{and} \quad \mathcal{ULB} \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}^2]}; \\ (\text{for } k \geq 2) \quad & \mathcal{ULB}' \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}^2 \oplus (\mathbb{Z}/2\mathbb{Z})^2]} \quad \text{and} \quad \mathcal{ULB} \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}^3 \oplus \mathbb{Z}/2\mathbb{Z}]}. \end{aligned}$$

For configurations of unoriented unlinks. We take $G = \text{Diff}(\underline{k}\mathbb{S}^1)$. We consider the homological functor

$$L_i(\mathring{F}'_{(\underline{k}\mathbb{S}^1, \text{Diff}(\underline{k}\mathbb{S}^1), \ell)}): \mathcal{ULB}' \longrightarrow \text{Mod}_{\bullet} \quad \text{and} \quad L_i(\mathring{F}_{(\underline{k}\mathbb{S}^1, \text{Diff}(\underline{k}\mathbb{S}^1), \ell)}): \mathcal{ULB} \longrightarrow \text{Mod}_{\bullet} \quad (5.21)$$

$$L_i(\mathring{F}'_{(\underline{k}\mathbb{S}^1, \text{Diff}(\underline{k}\mathbb{S}^1), \ell)}): \mathcal{ULB}' \longrightarrow \text{Mod}_{\bullet}.$$

associated with non-negative integers $i \geq 1$, $\ell \geq 0$ and $k \geq 1$ and groups $Q_{(\underline{k}\mathbb{S}^1, \text{Diff}(\underline{k}\mathbb{S}^1), \ell)}(\mathbb{D}_n^3)$. We recall from [DPS21, §5] that for any $n \geq 4$,

- for $k \in \mathbb{N} \setminus \{2, 3\}$: $\Gamma_2(\mathbf{LB}'_{k,n}) = \Gamma_3(\mathbf{LB}'_{k,n})$ and $\Gamma_2(\pi_1(U_k(\mathbb{D}_n^3))) \rtimes \mathbf{LB}_n = \Gamma_3(\pi_1(U_k(\mathbb{D}_n^3))) \rtimes \mathbf{LB}_n$;
- for $k \in \{2, 3\}$: $\Gamma_\ell(\mathbf{LB}'_{k,n}) \neq \Gamma_{\ell+1}(\mathbf{LB}'_{k,n})$ and $\Gamma_\ell(\pi_1(U_k(\mathbb{D}_n^3))) \rtimes \mathbf{LB}_n \neq \Gamma_{\ell+1}(\pi_1(U_k(\mathbb{D}_n^3))) \rtimes \mathbf{LB}_n$ for all $\ell \geq 1$.

Therefore, the construction produced for $\ell = 2$ is detailed below. Again, the procedure for the particular parameters $k = 2$ or 3 provides a specific representation for each $\ell \geq 2$, but will not be addressed in details here.

It follows from the computations of [DPS21, §5] that for $n \geq 2$:

- $Q'_{(k\mathbb{S}^1, \text{Diff}(k\mathbb{S}^1), 2)}(\mathbb{D}_n^3) = (\mathbb{Z}/2\mathbb{Z})^3$ if $k = 1$ and $(\mathbb{Z}/2\mathbb{Z})^5$ if $k \geq 2$;
- $Q_{(k\mathbb{S}^1, \text{Diff}(k\mathbb{S}^1), 2)}(\mathbb{D}_n^3) = \mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z})^2$ if $k = 1$ and $\mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z})^4$ if $k \geq 2$.

We deduce from Proposition 5.27 that:

Proposition 5.43 *For each $i \geq 0$ and $k \geq 1$, trivialising the assignment for $n \leq 1$, the functor (5.21) for $\ell = 2$ define homological functors:*

$$\begin{aligned} (\text{for } k = 1) \quad \mathfrak{ULB}' &\longrightarrow \text{Mod}_{\mathbb{Z}[(\mathbb{Z}/2\mathbb{Z})^3]} \quad \text{and} \quad \mathfrak{ULB} \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z})^2]}; \\ (\text{for } k \geq 2) \quad \mathfrak{ULB}' &\longrightarrow \text{Mod}_{\mathbb{Z}[(\mathbb{Z}/2\mathbb{Z})^5]} \quad \text{and} \quad \mathfrak{ULB} \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z} \oplus (\mathbb{Z}/2\mathbb{Z})^4]}. \end{aligned}$$

5.3. Global functors for mapping class groups

The construction in §5.1 of functors $\mathfrak{UD}_d \rightarrow \text{Mod}_\bullet$ is well-adapted for motion groups (a.k.a. *braided mapping class groups*, cf. Proposition 4.23). This is reflected in the fact that, *if* one restricts to a subcategory of \mathfrak{UD}_d of the form $\langle \mathcal{G}, \mathcal{M} \rangle$, where the automorphism groups of \mathcal{M} are contained in the corresponding *braided* diffeomorphism groups, then Proposition 5.16 implies (under additional conditions) that the functor takes values in $\text{Cov}_Q \subset \text{Cov}_\bullet$ (after passing to π_0), for a fixed group Q . Hence, applying the general construction of §2, we obtain a functor into Mod_R for some fixed ring R , rather than just Mod_\bullet . Thus the construction of §5.1 is adequate for constructing families of *untwisted* representations of motion groups (braided mapping class groups). However, for (full) mapping class groups, this construction typically only gives twisted representations, and one must restrict to smaller subgroups in order to obtain genuine, untwisted representations.

In this subsection, we describe a variant of the construction of §5.1, based on the split short exact sequence (4.9) instead of (4.6), which has good properties for the *full* mapping class groups. More precisely, this variant satisfies an analogue of Proposition 5.16 where one does *not* have to assume that the automorphism groups of \mathcal{M} are contained in the corresponding braided diffeomorphism groups (see Proposition 5.48 below).

Since the construction is very similar, we just sketch an outline and point out the differences from the construction of §5.1. The following is the analogue of Proposition 5.8.

Proposition 5.44 *For any integers $d \geq 2$ and $\ell \geq 0$, closed submanifold $Z \subset \mathbb{R}^d$ and open subgroup $G \leq \text{Diff}(Z)$, there are well-defined functors*

$$\mathring{F}_{(Z,G,\ell)} \quad \text{and} \quad F_{(Z,G,\ell)} : \mathfrak{UD}_d \longrightarrow \text{Cov}_\bullet, \quad (5.22)$$

constructed as described below.

The construction of the functor $\mathring{F}_{(Z,G,\ell)}$ on objects is similar to that of §5.1. The space $X(M, A)$ is defined in exactly the same way, but the quotient $\phi(M, A) : \pi_1(X(M, A)) \rightarrow Q(M, A)$ is defined using the following 6-term diagram instead of (5.4).

$$\begin{array}{ccccccc} & & \pi_1(X(M, A)) & & & & \\ & & \downarrow \mathbb{R} & & & & \\ 1 & \longrightarrow & \pi_1(\mathcal{E}_G(Z, \dot{M} \setminus A)) & \longrightarrow & \pi_0(\text{Diff}_{\text{dec}}(M, A, Z|G)) & \overset{\leftarrow}{\dashrightarrow} & \pi_0(\text{Diff}_{\text{dec}}(M, A)) \longrightarrow 1 \\ & & \downarrow \phi(M, A) & & \downarrow \gamma(M, A) & & \downarrow \bar{\gamma}(M, A) \\ 1 & \longrightarrow & Q(M, A) & \longrightarrow & \frac{\pi_0(\text{Diff}_{\text{dec}}(M, A, Z|G))}{\Gamma_\ell(\pi_0(\text{Diff}_{\text{dec}}(M, A, Z|G)))} & \overset{\leftarrow}{\dashrightarrow} & \frac{\pi_0(\text{Diff}_{\text{dec}}(M, A))}{\Gamma_\ell(\pi_0(\text{Diff}_{\text{dec}}(M, A)))} \longrightarrow 1 \end{array} \quad (5.23)$$

The top row is the split short exact sequence (4.9) of Corollary 4.19 when $G = \text{Diff}(Z)$; more generally it is the middle row of diagram (4.12) for open subgroups $G \leq \text{Diff}(Z)$. The rest of the diagram is constructed from this just as in §5.1, using Lemma 5.4. The construction of the functor on morphisms is then exactly as in §5.1.2, using the fact that the diagram (5.23) is functorial in the object (M, A) .

The closed variant $F_{(Z,G,\ell)}$ of the functor (5.22) is constructed similarly, using the embedding space $\mathcal{E}'_G(Z, M \setminus A)$ instead of $\mathcal{E}_G(Z, \dot{M} \setminus A)$ (cf. §5.1.3).

The observations in §5.1.4 hold also for the functors (5.22): they induce functors (not just semi-functors) on π_0 , the closed version $F_{(Z,G,\ell)}$ takes values in $\text{Cov}_{\bullet}^{\text{pr}}$ and there is a natural homotopy equivalence $\hat{F}_{(Z,G,\ell)} \Rightarrow F_{(Z,G,\ell)}$ between the two versions of (5.22).

The possible variants of the construction mentioned in §5.1.6, replacing the functorial quotient $G \mapsto G/\Gamma_\ell(G)$ by other quotients, apply similarly to the functors (5.22). Moreover, the constructions of §5.1.7 also carry over to these functors, and show that if we fix (Z, G) and allow ℓ to vary, the functors (5.22) form a “pro-nilpotent” tower of representations of \mathcal{UD}_d .

Remark 5.45 There is a natural morphism of diagrams (5.4) \rightarrow (5.23) induced by the map of split short exact sequences from the top row to the middle row of diagram (4.12). This induces a natural transformation of functors

$$(5.6) \Rightarrow (5.22). \quad (5.24)$$

Two actions agree. The diagram (5.23) is functorial in (M, A) as an object of \mathcal{UD}_d , so there is an action of $\text{Diff}_{\text{dec}}(M, A) = \text{Aut}_{\mathcal{D}_d}((M, A))$ on the bottom-left group $Q(M, A)$. Since $Q(M, A)$ is discrete, this factors through an action of $\pi_0(\text{Diff}_{\text{dec}}(M, A))$ on $Q(M, A)$.

On the other hand, $\pi_0(\text{Diff}_{\text{dec}}(M, A))$ is also the top-left group of (5.23), so it has another action on $Q(M, A)$ given either by lifting elements of $Q(M, A)$ along $\phi(M, A)$ and using the conjugation action of the semi-direct product on the top row, or equivalently by projecting along $\bar{\gamma}(M, A)$ to the bottom-right group of (5.23) and then using the conjugation action of the semi-direct product on the bottom row. The following is the analogue of Proposition 5.7.

Proposition 5.46 *The two actions of $\pi_0(\text{Diff}_{\text{dec}}(M, A))$ on $Q(M, A)$ described above are equal.*

Proof. This follows immediately from the first part of Lemma 4.25. \square

The definition of Q -stability in the setting of this section is exactly analogous to Definition 5.14.

Definition 5.47 Let \mathcal{M} and \mathcal{G} be subgroupoids as in (5.5). Suppose that there is a group Q and there are identifications $Q(M, A) \cong Q$ for each object (M, A) of $\langle \mathcal{G}, \mathcal{M} \rangle$, such that, for each object (M', A') of \mathcal{G} , the homomorphism of groups

$$Q(M, A) \longrightarrow Q((M', A')\natural(M, A)),$$

induced by the canonical morphism $((M', A'), \text{id})$ of $\langle \mathcal{G}, \mathcal{M} \rangle$, is equal to the identity under these identifications. In this case, we say that *the functor $F_{(Z,G,\ell)}$ is Q -stable on $\langle \mathcal{G}, \mathcal{M} \rangle$.*

The only difference between Definition 5.14 and Definition 5.47 is that, in the former, the group $Q(M, A)$ and the induced homomorphism $Q(M, A) \rightarrow Q((M', A')\natural(M, A))$ are those obtained from the 6-term diagram (5.4) (and its functoriality), whereas in the latter, they are obtained from the 6-term diagram (5.23) (and its functoriality).

The analogue of Lemma 5.15 is again immediate: Q -stability implies that $F_{(Z,G,\ell)}$ is equivalent to a functor taking values in $\text{Cov}_Q^{\text{tw}} \subset \text{Cov}_{\bullet}$. Moreover, we have an analogue of Proposition 5.16, except that we drop the second assumption of Proposition 5.16 in the following.

Proposition 5.48 *Let \mathcal{M} and \mathcal{G} be subgroupoids of \mathcal{D}_d such that \mathcal{G} is closed under \natural and \mathcal{M} is closed under the action of \mathcal{G} via \natural . Set $\ell = 2$, suppose that $F_{(Z,G,2)}$ is Q -stable on $\langle \mathcal{G}, \mathcal{M} \rangle$, and*

- \mathcal{G} is full in \mathcal{D}_d and \mathcal{M} is 0-full in \mathcal{D}_d ,

- $|\mathcal{G}|$ is free as a monoid and $|\mathcal{M}|$ is free as a $|\mathcal{G}|$ -set.

Under these assumptions, the functor

$$\pi_0(F_{(Z,G,2)}|_{\langle \mathcal{G}, \mathcal{M} \rangle}) : \pi_0(\langle \mathcal{G}, \mathcal{M} \rangle) \longrightarrow \pi_0(\text{Cov}_\bullet)$$

is equivalent to a functor with image contained in $\pi_0(\text{Cov}_Q) \subset \pi_0(\text{Cov}_\bullet)$.

Proof. The proof is exactly analogous to the proof of Proposition 5.16, except that we use Proposition 5.46, instead of Proposition 5.7, to understand the action of automorphisms of \mathcal{M} on Q . Since Proposition 5.46 applies to the whole mapping class group $\pi_0(\text{Diff}_{\text{dec}}(M, A))$ (whereas Proposition 5.7 applies only to the subgroup $\pi_0(\text{Diff}_{\text{dec}}^{\text{br}}(M, A))$, the braided mapping class group), we do not need to assume that the automorphisms of \mathcal{M} are contained in the braided diffeomorphism groups $\text{Diff}_{\text{dec}}^{\text{br}}(M, A)$, as we had to for Proposition 5.16. \square

5.4. Applications for mapping class groups of surfaces

Although it is best adapted to motion groups, the general construction of §5.1 may also be used to construct representations of mapping class groups, and recovers several classical constructions. This is detailed in §5.4.1. We then apply in §5.4.2 the adapted method for mapping class groups of §5.3 to define other representations of these groups.

We take this opportunity to recall and introduce some useful categories to handle mapping class groups of surfaces. We recall from §4.4 that $\mathcal{M}_2^t = \mathcal{D}_2^0$. Let $\mathcal{M}_2^{t,+}$ (respectively $\mathcal{M}_2^{t,-}$) be the full subcategories of \mathcal{M}_2^t with objects the orientable (respectively non-orientable) surfaces such that $\partial S = \partial_0 S$. We denote by \mathcal{M}_2^+ (respectively \mathcal{M}_2^-) the connected component $\pi_0 \mathcal{M}_2^{t,+}$ (respectively $\pi_0 \mathcal{M}_2^{t,-}$).

5.4.1. Homological representations from motion group functors.

A first idea to construct representations of the mapping class group of a d -manifold M is to use its action on covering spaces constructed in the spirit of §5.1. Let \mathcal{UD}_d^0 be the full category of the category \mathcal{UD}_d on the objects (M, \emptyset) . We consider the restriction of the continuous functor $\hat{F}_{(Z,G,\ell)} : \mathcal{UD}_d \rightarrow \text{Cov}_\bullet$ of Proposition 5.8 to \mathcal{UD}_d^0 . Note that this has the effect that the short exact sequences of the diagram (5.4) degenerates in the sense that its right-hand side is the trivial group.

We restrict our further study to the representations for mapping class groups of surfaces. Let $Z = \underline{k}$ a set of $k \geq 1$ distinct points seen as a closed submanifold of \mathbb{R}^2 . Under these restrictions, we thus consider the functors

$$\hat{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}^+ : \mathcal{UM}_2^+ \longrightarrow \text{Cov}_\bullet \quad \text{and} \quad \hat{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}^- : \mathcal{UM}_2^- \longrightarrow \text{Cov}_\bullet, \quad (5.25)$$

associated with a pair of non-negative integers (k, ℓ) where $\ell \geq 0$ and $k \geq 1$, with the lower central quotients of motion groups $\pi_1(\mathcal{E}_G(\underline{k}, \hat{S})) / \Gamma_\ell \pi_1(\mathcal{E}_G(\underline{k}, \hat{S}))$ (where S respectively belongs to $\mathcal{M}_2^{t,-}$ and $\mathcal{M}_2^{t,-}$) and with an open subgroup $G \leq \text{Diff}(\underline{k})$. Two choices for the group G naturally arise as relevant parameters to construct homological representations: either we consider *unordered* configurations spaces (see §5.4.1.1 below) or we take *ordered* configurations spaces; cf. §5.4.1.2.

5.4.1.1. Using unordered configuration spaces.

Similarly to the constructions of representations for motion groups in §5.2, we take $G = \mathfrak{S}_k$. Using the framework and tools of §2, we take \mathbb{Z} as ground ring and V to be the appropriate trivial fibrewise tensor product functor $V_{\hat{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}^+} : \mathcal{UM}_2^+ \rightarrow \text{Mod}_\bullet$ and $V_{\hat{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}^-} : \mathcal{UM}_2^- \rightarrow \text{Mod}_\bullet$. By Lemma 2.20, we define the homological functors for all $i \geq 1$:

$$L_i(\hat{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}^+) : \mathcal{UM}_2^+ \longrightarrow \text{Mod}_\bullet \quad \text{and} \quad L_i(\hat{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}^-) : \mathcal{UM}_2^- \longrightarrow \text{Mod}_\bullet. \quad (5.26)$$

For $\ell = 0$, we obtain the representations corresponding to the action on the homology groups of the universal covers of the spaces $X(M, \emptyset)$, and we recover the natural action of the motion groups

on the homology groups of the spaces $X(M, \emptyset)$ for $\ell = 1$; cf. Remark 5.6. We give more details of some representations encoded in the functor $L_i(\mathring{F}_{(k, \mathfrak{S}_k, \ell)}^+)$ depending on k .

If $k \leq 2$: For $k = 1$, we recall from [MKS04] that $\Gamma_\ell(\pi_1(\Sigma_{g,1})) \neq \Gamma_{\ell+1}(\pi_1(\Sigma_{g+1,1}))$ for all $\ell \geq 1$ and $g \geq 1$: the procedure of §5.1 thus provides a specific representation for each $\ell \geq 0$.

For $\ell = 0$, since a universal cover is simply connected, $L_1(\mathring{F}_{(\underline{1}, \mathfrak{S}_1, 0)}^+)$ is the trivial functor. However, considering the variant using *reduced* homology, $L_1^r(\mathring{F}_{(\underline{1}, \mathfrak{S}_1, 0)}^+)$ encodes the (*universal*) *Magnus representations* of the mapping class groups. These were originally defined using the Fox free differential calculus (see [Bir74; Sak12] for further details) and Suzuki [Suz05] introduced the equivalent topological definition.

For $\ell = 1$, the functor $L_1(\mathring{F}_{(\underline{1}, \mathfrak{S}_1, 1)}^+)$ encodes the natural actions on the first homology groups of the surfaces, that will be denoted for each g by $\mathfrak{a}_g : \Gamma_{g,1} \rightarrow \text{Aut}_{\mathbb{Z}}(H_1(\Sigma_{g,1}, \mathbb{Z}))$ for convenience.

For $\ell \geq 2$, considering the variant with *reduced* homology, $L_1^r(\mathring{F}_{(\underline{1}, \mathfrak{S}_1, \ell)}^+)$ defines the *level- ℓ Magnus representations* of the mapping class groups (see [Suz05, §3]). Note that the hypothesis (*Q-stability*) of Proposition 5.16 is satisfied if and only if we restrict to the kernels of the induced representations $\Gamma_{g,1} \rightarrow \text{Aut}_{\mathbb{Z}}(\pi_1(\Sigma_{g,1}, \mathbb{Z})/\Gamma_\ell \pi_1(\Sigma_{g,1}, \mathbb{Z}))$ (for instance, it is the Torelli group for $\ell = 2$). Then, it simply follows from the 4-term exact sequence of the homology of a free group that the representations encoded by $L_1(\mathring{F}_{(\underline{1}, \mathfrak{S}_1, \ell)}^+)$ are proper submodules of the level- ℓ Magnus representations.

The case of $k = 2$ represents a more difficult situation. In particular, for the torus with one boundary component $\Sigma_{1,1}$, [BGG08, §4] proves that $\Gamma_\ell(\mathbf{B}_2(\Sigma_{1,1})) \neq \Gamma_{\ell+1}(\mathbf{B}_2(\Sigma_{1,1}))$ for all $\ell \geq 1$. As far as the authors know, the representations defined by the functors $L_i(\mathring{F}_{(2, \mathfrak{S}_2, \ell)}^+)$ have not been addressed in the literature yet.

If $k \geq 3$: We recall from [BGG08, Theorem 2] that for any $g \geq 1$ and $k \geq 3$, $\Gamma_2(\mathbf{B}_k(\Sigma_{g,1})) \neq \Gamma_3(\mathbf{B}_k(\Sigma_{g,1})) = \Gamma_4(\mathbf{B}_k(\Sigma_{g,1}))$. Hence, for each $i \geq 0$, the homological functors $L_i(\mathring{F}_{(k, \mathfrak{S}_k, 2)}^+)$ and $L_i(\mathring{F}_{(k, \mathfrak{S}_k, 3)}^+)$ are the only relevant ones to study. We determine the best subgroup of the mapping class groups which acts on the representation spaces as $\mathbf{B}_k(\Sigma_{g,1})/\Gamma_\ell(\mathbf{B}_k(\Sigma_{g,1}))$ -modules.

We fix a symplectic basis $\{A_1, B_1, \dots, A_g, B_g\}$ for the first homology group of the surface $H_g := H_1(\Sigma_{g,1}; \mathbb{Z})$ with respect to the algebraic intersection form $\omega_g : H_g \times H_g \rightarrow \mathbb{Z}$. Moreover, the operation $(z, c) \cdot (z', c) = (z + z' + \omega_g(\lambda, \lambda'), \lambda + \lambda')$ for all $z, z' \in \mathbb{Z}$ and $\lambda, \lambda' \in H_g$ defines a central extension denoted by $\mathbb{Z} \times_{\omega_g} H_g$. Then:

Lemma 5.49 *The abelianisation $\mathbf{B}_k(\Sigma_{g,1})/\Gamma_2(\mathbf{B}_k(\Sigma_{g,1}))$ is isomorphic to the product $\mathbb{Z}/2\mathbb{Z} \times H_g$. The third lower central quotient $\mathbf{B}_k(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_k(\Sigma_{g,1}))$ is isomorphic to the central extension $\mathbb{Z} \times_{\omega_g} H_g$. In both situations, the action of the mapping class group $\Gamma_{g,1}$ on the quotient H_g is the natural action \mathfrak{a}_g .*

Proof. For $\ell = 2$, the result directly follows from the presentation of $\mathbf{B}_k(\Sigma_{g,1})$ of [Bel04, Theorem 1]: the factor $\mathbb{Z}/2\mathbb{Z}$ is the image of the braid generators $\{\sigma_1, \dots, \sigma_{k-1}\}$ and the factor H_g is the image of the generators $\{a_1, b_1, \dots, a_g, b_g\}$. For $\ell = 3$, by [BGG17, Corollary 3.14] the third lower central quotient $\mathbf{B}_k(\Sigma_{g,1})/\Gamma_3(\mathbf{B}_k(\Sigma_{g,1}))$ is isomorphic to the semi-direct product $(\mathbb{Z} \times \mathbb{Z}^g) \rtimes \mathbb{Z}^g$, the first factor \mathbb{Z} is central and is generated by $\sigma := \gamma_3(\sigma_i)$ for all $i \in \{1, \dots, k-1\}$, the second factor \mathbb{Z}^g is generated by $\{a_i := \gamma_3(a_i) \mid i \in \{1, \dots, g\}\}$, and the third factor \mathbb{Z}^g is generated by $\{b_i := \gamma_3(b_i) \mid i \in \{1, \dots, g\}\}$; for all $j \in \{1, \dots, g\}$, the generator b_j acts trivially on a_i for $i \in \{1, \dots, g\} \setminus \{j\}$ and $a_j b_j = \sigma^2 b_j a_j$. The isomorphism is given by sending σ to the generator of \mathbb{Z} in the central extension, a_i to A_i and b_i to B_i for all $i \in \{1, \dots, g\}$. The relation $a_j b_j = \sigma^2 b_j a_j$ is preserved through this morphism by the definition of the intersection form. \square

By Lemma 5.49, we must restrict to a subgroup of the Torelli group to obtain a trivial actions both on the abelianisation and on the third lower central quotient. In particular the $\mathbb{Z}/2\mathbb{Z}$ summand is generated by the image σ of the braid generators of $\mathbf{B}_n(\Sigma_{g,1})$. We recall from Corollary 4.19 that the splitting $\Gamma_{g,1} \hookrightarrow \Gamma_{g,1}^n$ is induced by the embedding of surfaces $\Sigma_{g,1} \hookrightarrow \Sigma_{0,1}^n \natural \Sigma_{g,1} \cong \Sigma_{g,1}^n$: this implies that $\Gamma_{g,1}$ acts trivially on σ , since σ is supported in the subsurface $\Sigma_{0,1}^n$. Also the action of $\Gamma_{g,1}$ on H_g is induced by \mathfrak{a}_g . Hence the result of Lemma 5.49 on $\mathbf{B}_k(\Sigma_{g,1})/\Gamma_2(\mathbf{B}_k(\Sigma_{g,1}))$

is a $\Gamma_{g,1}$ -module isomorphism, not just of abelian groups. Therefore, for $\ell = 2$, the homological representations defined by $L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_{k,2})}^+)$ acts as $(\mathbb{Z}/2\mathbb{Z} \times H_g)$ -modules if we restrict to the Torelli groups.

For $\ell = 3$, we deduce that there exists a map $\kappa : \Gamma_{g,1} \rightarrow \text{Hom}(H_g, \mathbb{Z})$ so that the action of the mapping class group on the central extension $\mathbb{Z} \times_{\omega_g} H_g$ is described by the matrix

$$\begin{bmatrix} Id_{\mathbb{Z}} & \kappa \\ 0 & \mathfrak{a}_g \end{bmatrix}.$$

We denote by *Chill* the *Chillingworth homomorphism* [Chi72] which describes the action of $\mathcal{I}_{g,1}$ on the winding numbers of the curves of $\Sigma_{g,1}$. Its kernel is called the *Chillingworth subgroup* and denoted by $\mathcal{C}_{g,1}$. By [Chi12, Corollary 4.8], it is the subgroup generated by the union of the *simply intersecting pair maps* and the *Johnson subgroup* [Joh83], which is the kernel of the natural map $\Gamma_{g,1} \rightarrow \text{Aut}_{\mathbb{Z}}(\pi_1(\Sigma_{g,1}, p_0)/\Gamma_3(\pi_1(\Sigma_{g,1}, p_0)))$. Then:

Lemma 5.50 *The map κ is a crossed homomorphism and its kernel coincides with $\mathcal{C}_{g,1}$.*

Proof. Since the action of the mapping class group on the central extension is a morphism, we deduce that $\kappa(\varphi \circ \psi) = \kappa(\psi) + \kappa(\varphi)\mathfrak{a}_g(\psi)$ for all $\varphi, \psi \in \Gamma_{g,1}$: this proves that κ is a crossed homomorphism. Moreover [Mor89] proves that $H^1(\Gamma_{g,1}, H_g) \cong \mathbb{Z}$. Hence $\kappa = \lambda \cdot \text{Chill} + c$ where $\lambda \in \mathbb{Z}$ and c is a principal crossed homomorphism: restricting to the Torelli group, we deduce that $\ker(\kappa) = \ker(\text{Chill})$. \square

Hence the homological representations defined by $L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_{k,3})}^+)$ acts as $(\mathbb{Z} \times_{\omega_g} H_g)$ -modules if we restrict to the Chillingworth subgroups.

5.4.1.2. Using ordered configuration spaces.

Alternatively, we consider ordered configuration spaces of points on the surfaces: we take the same assignments as in §5.4.1.1 except that $G = 0_{\text{Grp}}$ the trivial group. Hence we define the homological functors $L_i(\mathring{F}_{(\underline{k}, 0_{\text{Grp}}, \ell)}^+): \mathfrak{M}_2^+ \rightarrow \text{Mod}_{\bullet}$ and $L_i(\mathring{F}_{(\underline{k}, 0_{\text{Grp}}, \ell)}^-): \mathfrak{M}_2^- \rightarrow \text{Mod}_{\bullet}$ for all $i \geq 1$.

An interesting modification of this construction consists in removing the basepoint p_0 from the configuration space and allowing the configuration points to be in the boundary of the surface. Namely, we use the closed variant F of §5.1.3 instead of the functor \mathring{F} and consider the configurations in the surface $S \setminus \{p_0\}$ for each object S . By Lemma 5.11, Borel-Moore homology may be applied for this alternative and has the advantage to be endowed with a natural free generating set; cf. [Big04, Lemma 3.1], [AK10, Lemma 3.3], [AP20, Theorem 6.6] or [PS21, §5].

The Moriyama functors. For each $k \geq 1$ and $g \geq 0$, Moriyama [Mor07] studies the natural action of the mapping class group $\Gamma_{g,1}$ on the relative homology group $H_k(F_k(\Sigma_{g,1}), \mathcal{Y}_k(\Sigma_{g,1}, p_0); \mathbb{Z})$ where $\mathcal{Y}_k(\Sigma_{g,1}, p_0)$ denotes the set $\{(x_1, \dots, x_k) \in \Sigma_{g,1}^{\times k} \mid x_i = p_0 \text{ for some } i\}$ for each $k \geq 0$. Its kernel coincides with the kernel of the natural action on the k -th lower central quotient group of the fundamental group of $\Sigma_{g,1}$.

Theorem 5.51 *For each $k \geq 1$, the representations encoded by the functor $L_k^{BM}(F_{(\underline{k}, 0_{\text{Grp}}, 1)}^+): \mathfrak{M}_2^+ \rightarrow \text{Mod}_{\bullet}$ are equivalent to those studied in [Mor07]. We thus denote this functor by \mathfrak{Mor}_k and call it the k -th Moriyama functor.*

Proof. For each $g \geq 0$, it follows from the basic properties of Borel-Moore homology (see [Bre97, Corollary V.5.10] for instance) that we have the following abelian group isomorphism between $H_k^{BM}(F_k(\Sigma_{g,1} \setminus \{p_0\}); \mathbb{Z})$ and $H_k(F_k(\Sigma_{g,1}), \mathcal{Y}_k(\Sigma_{g,1}, p_0); \mathbb{Z})$. It defines a $\Gamma_{g,1}$ -module isomorphism since this structure is induced by the diagonal action in both situations. \square

5.4.2. Homological representations from mapping class group functors.

The quotient groups Q defining the representations constructed in §5.4.1 for the mapping class groups of the surfaces are typically *not* independent of the surface (i.e. we do not have Q -stability in the sense of Definition 5.14). The advantage of the method of §5.3 applied here is that it is much more adequate for this property to be satisfied. We consider the restriction of the continuous functor $\mathring{F}_{(Z,G,\ell)}: \mathcal{MD}_2 \rightarrow \text{Cov}_\bullet$ of Proposition 5.44 to the appropriate full subcategory of the form \mathcal{UG} given by Lemma 3.7 applied to the examples of subgroupoids \mathcal{G} of §4.4 (whose automorphism groups correspond to the family of mapping class groups). Moreover we take $Z = \underline{k}$ a set of $k \geq 1$ distinct points seen as a closed submanifold of \mathbb{R}^2 and set $G = \mathfrak{S}_k$.

Using the framework and tools of §2, we take \mathbb{Z} as ground ring \mathbb{A} and V to be the trivial fibrewise tensor product functor $V_{\mathring{F}_{(Z,G,\ell)}}: \mathcal{UG} \rightarrow \text{Mod}_\bullet$. Then, by Lemma 2.20, we define the homological functors for all $i \geq 1$:

$$L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}): \pi_0 \mathcal{UG} \longrightarrow \text{Mod}_\bullet. \quad (5.27)$$

associated with non-negative integers $i \geq 1$, $\ell \geq 0$ and $k \geq 1$. For $\ell = 0$, we obtain the representations corresponding to the action on the homology groups of the universal cover of the configuration spaces of points in the surface, and we recover the natural action of the motion groups on the homology groups on these configuration spaces for $\ell = 1$; cf. Remark 5.6. For $\ell = 2$, we will typically prove that the Q -stability property (cf. Definition 5.14) is satisfied. In this case, denoting by $Q_{(\underline{k}, \mathfrak{S}_k, 2)}$ the associated stable quotient, we deduce that:

Proposition 5.52 *We define a homological functor $L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, 2)}): \pi_0(\mathcal{UG}) \rightarrow \text{Mod}_{\mathbb{Z}[Q_{(\underline{k}, \mathfrak{S}_k, 2)}]}$ from the functor (5.27) for all $i \geq 0$.*

Proof. Let $\mathcal{UG}_{\text{stab}}$ be the full subcategory of \mathcal{UG} on all objects *except* those for which the obtained quotient is not isomorphic to $Q_{(\underline{k}, \mathfrak{S}_k, 2)}$. By Proposition 5.16, the functor (5.27) induces functors $L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, 2)}): \mathcal{UG}_{\text{stab}} \rightarrow \text{Mod}_{\mathbb{Z}[Q_{(\underline{k}, \mathfrak{S}_k, 2)}]}$. By Lemma 5.17, we extend them to functors over \mathcal{UG} by sending the other objects to the trivial group. \square

As far as the authors know, the representations encoded by the functors based on the above procedure and described in the following Propositions 5.54 and 5.55 do not appear in the literature and therefore appear to be new.

General alternatives. For all the constructions presented here, we could define interesting alternatives by taking the parameter G between the trivial group 0_{Grp} and $\text{Diff}(Z)$: this amounts to considering *partitioned* configurations. The associated lower central series being more complicated generally speaking (see [DPS21]), the representations that arise from these alternatives may be more sophisticated and would deserve more study.

5.4.2.1. For orientable surfaces.

We consider the restriction of the continuous functor $\mathring{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}: \mathcal{MD}_2 \rightarrow \text{Cov}_\bullet$ of Proposition 5.8 to the full subcategory $\mathcal{MM}_2^{t,+}$. Hence we obtain the homological functor

$$L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}): \mathcal{MM}_2^+ \longrightarrow \text{Mod}_\bullet, \quad (5.28)$$

associated with non-negative integers $i \geq 1$, $k \geq 1$, $\ell \geq 0$. To obtain some restrictions on the possibilities for the parameter ℓ , we study the lower central series of the mapping class group $\Gamma_{g,1}^k$. We compute its abelianisation:

Lemma 5.53 *For each $g \geq 3$, then $(\Gamma_{g,1}^k)^{\text{ab}}$ is trivial if $k \leq 1$ and isomorphic to $\mathbb{Z}/2\mathbb{Z}$ if $k \geq 2$.*

Proof. We recall from Corollary 4.19 that $\Gamma_{g,1}^k$ is the semi-direct product $\mathbf{B}_k(\Sigma_{g,1}) \rtimes \Gamma_{g,1}$. By abelianisation of a semi-direct product (see [GG09, Proposition 3.3] for instance), we deduce that

$(\Gamma_{g,1}^k)^{\text{ab}}$ is isomorphic to $(\mathbf{B}_k(\Sigma_{g,1})^{\text{ab}})_{\Gamma_{g,1}} \oplus (\Gamma_{g,1})^{\text{ab}}$, where $(\mathbf{B}_k(\Sigma_{g,1})^{\text{ab}})_{\Gamma_{g,1}}$ denotes the coinvariants of $\mathbf{B}_k(\Sigma_{g,1})^{\text{ab}}$ under the natural action of $\Gamma_{g,1}$ on $\Sigma_{g,1}$. We recall from [Kor02, Theorem 5.1] that the abelianisation of $(\Gamma_{g,1})^{\text{ab}}$ is trivial for any $g \geq 3$. It therefore just remains to show that the left-hand summand is isomorphic to $\mathbb{Z}/2\mathbb{Z}$. By [BGG17, Proposition 3.4], we know that

$$\mathbf{B}_k(\Sigma_{g,1})^{\text{ab}} \cong H_1(\Sigma_{g,1}) \oplus \mathbb{Z}/2\mathbb{Z}, \quad (5.29)$$

where the generator σ of the $\mathbb{Z}/2\mathbb{Z}$ summand is the image of all the braid generators of $\mathbf{B}_k(\Sigma_{g,1})$. The splitting $\Gamma_{g,1} \hookrightarrow \Gamma_{g,1}^k$ of the exact sequence is induced by the embedding of surfaces $\Sigma_{g,1} \hookrightarrow \Sigma_{0,b}^k \natural \Sigma_{g,1} \cong \Sigma_{g,1}^k$, so $\Gamma_{g,1}$ acts trivially on σ , since σ is supported in the subsurface $\Sigma_{0,b}^k$.

The action of $\Gamma_{g,1}$ on the left-hand summand of (5.29) is induced by the natural action of $\Gamma_{g,1}$ on the homology of the surface. Together with the fact that its action on σ is trivial, this in particular implies that (5.29) is a splitting of representations of $\Gamma_{g,1}$, not just of abelian groups. The action of the Dehn twists $\tau_{A_i}, \tau_{B_i} \in \Gamma_{g,1}$ on $H_1(\Sigma_{g,1}) = \langle A_1, \dots, A_g, B_1, \dots, B_g \rangle$ satisfies $\tau_{A_i}(B_i) - B_i = A_i$ and $\tau_{B_i}(A_i) - A_i = B_i$, so we have $H_1(\Sigma_{g,1})_{\Gamma_{g,1}} = 0$. Putting this all together, we end the proof. \square

Hence, since $(\Gamma_{g,1}^k)^{\text{ab}}$ is a cyclic group, it follows from [DPS21, Corollary 2.2] that $\Gamma_2(\Gamma_{g,1}^k) = \Gamma_3(\Gamma_{g,1}^k)$ for all $k \geq 1$ and $g \geq 3$: hence we can restrict our study to the case of $\ell = 2$. Moreover, we deduce from the computation of Lemma 5.53 that for all $k \geq 1$, if $g \geq 2$ then

$$Q_{(\underline{k}, \mathfrak{S}_k, 2)}(\Sigma_{g,1}) = Q_{(\underline{k}, \mathfrak{S}_k, 2)}(\Sigma_{g+1,1}) \cong \begin{cases} 0 & \text{if } k = 1, \\ \mathbb{Z}/2\mathbb{Z} & \text{if } k \geq 2. \end{cases}$$

We denote this quotient by $Q_k(\Sigma)$ and deduce from Proposition 5.52 that:

Proposition 5.54 *For each $i \geq 0$ and $k \geq 1$, the functor (5.28) for $\ell = 2$ defines a homological functor $L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, 2)}): \mathfrak{UM}_2^+ \rightarrow \text{Mod}_{\mathbb{Z}[Q_k(\Sigma)]}$.*

5.4.2.2. For non-orientable surfaces.

We consider the restriction of the continuous functor $\mathring{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}: \mathfrak{UD}_2 \rightarrow \text{Cov}_\bullet$ of Proposition 5.8 to the full subcategory \mathfrak{UM}_2^{\pm} . Hence we define the homological functor

$$L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, \ell)}): \mathfrak{UM}_2^- \longrightarrow \text{Mod}_\bullet, \quad (5.30)$$

associated with non-negative integers $i \geq 1$, $k \geq 1$, $\ell \geq 0$. Again, we study the lower central series of the mapping class group $\mathcal{N}_{h,1}^k$ to obtain some restrictions on the possibilities for the parameter ℓ . We consider the following elements of $\mathcal{N}_{h,1}^k$, for $k \geq 3$ and $h \geq 7$ introduced in [Stu10, §2]

- A braid-type element σ that interchanges two marked points.
- A marked point slide c that sends a marked point through the core of a cross-cap.
- A cross-cap slide y supported in a genus-2 subsurface of the surface.

We recall from [Stu10, Theorems 6.21 and 6.22] that for all $h \geq 7$ and $k \geq 3$, $(\mathcal{N}_{h,1}^k)^{\text{ab}} \cong (\mathbb{Z}/2\mathbb{Z})^3$ – generated by the elements σ , c and y – and that $(\mathcal{N}_{h,1})^{\text{ab}} \cong \mathbb{Z}/2\mathbb{Z}$ – generated by the element y . Using the same type of disjoint support arguments as [DPS21], we deduce that $\Gamma_2(\mathcal{N}_{h,1}^k) = \Gamma_3(\mathcal{N}_{h,1}^k)$ for $k \geq 1$ and h an **even** number such that $h \geq 7$. We thus restrict our study to the case of $\ell = 2$. We also deduce from the above abelianisation results that for all $k \geq 3$, if $h \geq 7$ then

$$Q_{(\underline{k}, \mathfrak{S}_k, 2)}(\mathcal{N}_{h,1}) = Q_{(\underline{k}, \mathfrak{S}_k, 2)}(\mathcal{N}_{h+1,1}) \cong (\mathbb{Z}/2\mathbb{Z})^2.$$

We deduce from Proposition 5.52 that:

Proposition 5.55 *For each $i \geq 0$ and $k \geq 3$, the functor (5.30) for $\ell = 2$ defines a homological functor $L_i(\mathring{F}_{(\underline{k}, \mathfrak{S}_k, 2)}): \mathfrak{UM}_2^- \rightarrow \text{Mod}_{\mathbb{Z}[(\mathbb{Z}/2\mathbb{Z})^2]}$.*

6. Iterative constructions

We illustrate in this section the applications of the iterative construction of §2.7. We fix an associative unital ring \mathbb{A} and write $\mathbb{A}\text{-Alg}$ for the category of associative, unital \mathbb{A} -algebras. Throughout this section, we fix a monoidal groupoid $(\mathcal{G}, \natural, 0)$ and a left-module (\mathcal{M}, \natural) over \mathcal{G} . For sake of simplicity, we directly take the categories \mathcal{G} and \mathcal{M} to be *discrete* (i.e. $\pi_0(\mathcal{G}) = \mathcal{G}$ and $\pi_0(\mathcal{M}) = \mathcal{M}$): typically we consider the skeletons of the path-components of the categories and modules introduced in §4. Moreover, forgetting all morphisms and viewing \mathcal{G} as a monoid acting on the set \mathcal{M} , we assume that these are both isomorphic to the non-negative integers \mathbb{N} and the action is equivalent to \mathbb{N} acting on itself by addition. Therefore, there exist two objects $1 \in \text{ob}(\mathcal{G})$ and $\hat{0} \in \text{ob}(\mathcal{G})$ so that any object X of \mathcal{M} is isomorphic to the monoidal product $\underline{n} := 1^{\natural n} \natural \hat{0}$ for some $n \in \mathbb{N}$. The monoidal structure thus defines an endofunctor $k\natural - : \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \langle \mathcal{G}, \mathcal{M} \rangle$ for each $k \geq 1$. We denote by G_n the isomorphism group $\text{Iso}_{\langle \mathcal{G}, \mathcal{M} \rangle}(\underline{n})$.

Let $F : \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \pi_0(\text{Cov}_\bullet)$ be a functor that sends each $\underline{n} \in \text{ob}(\langle \mathcal{G}, \mathcal{M} \rangle)$ to a path-connected, locally path-connected and semi-locally simply-connected based space $F(\underline{n})$, together with the quotient $\phi = \text{id}_{\pi_1(F(\underline{n}))} : \pi_1(F(\underline{n})) \rightarrow \pi_1(F(\underline{n}))$. In particular, the functor F thus encodes the universal cover of the space $F(\underline{n})$. Let $\langle \mathcal{G}, \mathcal{M} \rangle_F$ be the category with the objects of \mathcal{M} and the product of morphisms $\text{Im}(r \circ F([n' - n, \phi])) \times [n' - n, \phi]$ for all $[n' - n, \phi] \in \langle \mathcal{G}, \mathcal{M} \rangle(n, n')$, where $\text{Im}(r \circ F([n' - n, \phi]))$ is the induced group morphism $r \circ F(n) \rightarrow r \circ F(n')$.

In the notation of Definition 2.21, one may easily check that the assignment $W \mapsto L_i^r(F; W)$ defines a functor on the functor categories

$$L_i^r(F; -) : \mathbf{Fct}(\langle \mathcal{G}, \mathcal{M} \rangle_F, \text{Mod}_\bullet) \longrightarrow \mathbf{Fct}(\langle \mathcal{G}, \mathcal{M} \rangle, \text{Mod}_\bullet). \quad (6.1)$$

We recall that Grp denotes the category of groups and from Definition 2.28 that the functor $r \circ F : \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \text{Grp}$ defines a category $(r \circ F) \times \langle \mathcal{G}, \mathcal{M} \rangle$ by using the Grothendieck construction. The key to make the construction (6.1) iterative is to consider a functor $\chi_{r \circ F} : (r \circ F) \times \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \langle \mathcal{G}, \mathcal{M} \rangle$ such that the composite $\langle \mathcal{G}, \mathcal{M} \rangle \xrightarrow{s_{r \circ F}} (r \circ F) \times \langle \mathcal{G}, \mathcal{M} \rangle \xrightarrow{\chi_{r \circ F}} \langle \mathcal{G}, \mathcal{M} \rangle$ is equal to the endofunctor $k\natural -$ for some $k \geq 1$. The interpretation of this condition at the level of automorphisms is that the functor $\chi_{r \circ F}$ induces an homomorphism $(r \circ F)(\underline{n}) \rtimes G_n \rightarrow G_{k+n}$. Using *reduced* homology, Proposition 2.31 then defines the iterative construction

$$\Lambda_i^r(F; \chi_{r \circ F}) : \mathbf{Fct}(\langle \mathcal{G}, \mathcal{M} \rangle, \text{Mod}_\bullet) \longrightarrow \mathbf{Fct}(\langle \mathcal{G}, \mathcal{M} \rangle, \text{Mod}_\bullet). \quad (6.2)$$

6.1. Connection to the Long-Moody functors

Long and Moody [Lon94] introduced a recipe to construct representations of braid groups. This method and its variants have been studied with a functorial point of view in [Sou19b] for braid groups and then generalised in [Sou19a] for general families of groups. We give their connections to the above iterative construction of homological functors.

First of all, we consider a functor $\mathcal{A} : \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \text{Grp}$. We recall that the group rings and the augmentation ideals respectively define the group algebra functor $\mathbb{A}[-] : \text{Grp} \rightarrow \mathbb{A}\text{-Alg}$ and the augmentation ideal functor $\mathcal{I}_{\mathbb{A}[-]} : \text{Grp} \rightarrow \mathbb{A}\text{-Alg}$ respectively. We respectively denote by $\mathbb{A}[\mathcal{A}]$ and $\mathcal{I}_{\mathcal{A}}$ the composite functors $\mathbb{A}[-] \circ \mathcal{A}$ and $\mathcal{I}_{\mathbb{A}[-]} \circ \mathcal{A}$. Then $\mathbb{A}[\mathcal{A}]$ is a monoid object in $\mathbf{Fct}(\langle \mathcal{G}, \mathcal{M} \rangle, \text{Mod}_\bullet)$. Hence pointwise tensor product of functors induces the *tensor product functor over $\mathbb{A}[\mathcal{A}]$* (cf. [Sou19a, Definition 2.1])

$$- \otimes_{\mathbb{A}[\mathcal{A}]} - : \text{Mod}_{\mathbb{A}[\mathcal{A}]} \times_{\mathbb{A}[\mathcal{A}]} \text{Mod} \rightarrow \mathbf{Fct}(\langle \mathcal{G}, \mathcal{M} \rangle, \text{Mod}_\bullet)$$

where ${}_{\mathbb{A}[\mathcal{A}]} \text{Mod}$ and $\text{Mod}_{\mathbb{A}[\mathcal{A}]}$ respectively denote the categories of left and right modules over the monoid object $\mathbb{A}[\mathcal{A}]$. Moreover $\mathcal{I}_{\mathcal{A}}$ is a right $\mathbb{A}[\mathcal{A}]$ -module and thus defines a functor $\mathcal{I}_{\mathcal{A}} \otimes_{\mathbb{A}[\mathcal{A}]} - : {}_{\mathbb{A}[\mathcal{A}]} \text{Mod} \rightarrow \mathbf{Fct}(\langle \mathcal{G}, \mathcal{M} \rangle, \text{Mod}_\bullet)$.

Finally we consider a functor $\chi_{\mathcal{A}} : \mathcal{A} \times \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \langle \mathcal{G}, \mathcal{M} \rangle$ such that $\chi_{\mathcal{A}} \circ s_{\mathcal{A}} = 1\natural -$. The appropriate data \mathcal{A} and $\chi_{\mathcal{A}}$ naturally arise for many families of groups in connection with topology. In particular, we refer the reader to [Sou19a, §3] for the introduction of non-trivial and natural such functors for the families surface braid groups and mapping class groups of surfaces.

Definition 6.1 The Long-Moody functor $\mathbf{LM}_{\mathcal{A}, \chi_{\mathcal{A}}}$ associated with the functors \mathcal{A} and $\chi_{\mathcal{A}}$ is the composite

$$\mathbf{Fct}(\langle \mathcal{G}, \mathcal{M} \rangle, \mathbf{Mod}_{\bullet}) \xrightarrow{s_{\mathcal{A}}^* \circ \chi_{\mathcal{A}}^*} {}_{\mathbb{A}[\mathcal{A}]} \mathbf{Mod} \xrightarrow{\mathcal{I}_{\mathcal{A}} \otimes_{\mathbb{A}[\mathcal{A}]} -} \mathbf{Fct}(\langle \mathcal{G}, \mathcal{M} \rangle, \mathbf{Mod}_{\bullet}),$$

where $s_{\mathcal{A}}^*$ and $\chi_{\mathcal{A}}^*$ respectively denote the precomposition functors $s_{\mathcal{A}}$ and $\chi_{\mathcal{A}}$.

Let \mathbf{FGrp} be the full subcategory of \mathbf{Grp} on the finitely-generated free groups.

Theorem 6.2 *If the functor \mathcal{A} factors through the category of \mathbf{FGrp} , then there exists a functor $F: \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \pi_0(\mathbf{Cov}_{\bullet})$ such that $r \circ F = \mathcal{A}$ and $\mathbf{LM}_{\mathcal{A}, \chi_{\mathcal{A}}} \cong \Lambda_1^r(F; \chi_{r \circ F})$.*

Proof. The forgetful functor $U: \mathbf{Cov}_{\bullet} \rightarrow \mathbf{Top}_{*}$ admits a section given by sending a based space X to the pair (X, \tilde{X}) , where \tilde{X} denotes the universal covering of X . Also, the fundamental group functor $\pi_1: \pi_0(\mathbf{Top}_{*}) \rightarrow \mathbf{Grp}$ admits a section defined by sending a group G to an Eilenberg–MacLane space of type $K(G, 1)$, which is unique up to homotopy (see [Hat02, Proposition 1.B.8] for instance). Hence the functor $\mathcal{A}: \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \mathbf{Grp}$ automatically admits a lift $F: \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \pi_0(\mathbf{Cov}_{\bullet})$ such that $\mathcal{A} = r \circ F$.

We fix $\underline{n} \in \mathbf{ob}(\langle \mathcal{G}, \mathcal{M} \rangle)$ and $V \in \mathbf{ob}(\mathbf{Fct}(\langle \mathcal{G}, \mathcal{M} \rangle, \mathbf{Mod}_{\bullet}))$. Note that $\pi_1(F(\underline{n})) = \mathcal{A}(\underline{n})$ is a finitely generated free group and therefore $F(\underline{n})$ is homotopy equivalent to a bouquet of circles of the rank of $F(\underline{n})$ as a free group. From the cell structure, we deduce the following isomorphism for the reduced homology group with local coefficient:

$$H_1(F(\underline{n}), p_0; V(\underline{n+1})) \cong \mathcal{I}_{\mathcal{A}(\underline{n})} \otimes_{\mathbb{A}[\mathcal{A}(\underline{n})]} V(\underline{n+1}).$$

It directly follows from the definitions that this abelian group isomorphism defines an isomorphism $\mathbf{LM}_{\mathcal{A}, \chi_{\mathcal{A}}}(V) \cong \Lambda_1^r(F; \chi_{r \circ F})(V)$ of $\mathbf{Fct}(\langle \mathcal{G}, \mathcal{M} \rangle, \mathbf{Mod}_{\bullet})$, which is easily proved to be natural in V . \square

Remark 6.3 In most examples of [Sou19a], the various functors \mathcal{A} factor across \mathbf{FGrp} and are actually induced by geometrical constructions: the associated Long-Moody functors are thus particular cases of the iterative construction (6.2). However, since the first homology of a group is not necessarily related to its augmentation ideal generally speaking, the general framework of [Sou19a] which considers a functor $\mathcal{A}: \langle \mathcal{G}, \mathcal{M} \rangle \rightarrow \mathbf{Grp}$ cannot be fully recovered by the iterative homological constructions. On the other hand, the construction (6.2) offers others possibilities, in particular on the degree of the homology groups that are considered.

6.2. Recovering the Tong-Yang-Ma representations

Tong, Yang and Ma [TYM96] introduce a family of irreducible representations of braid groups, of same dimension as the unreduced Burau representations but not equivalent to them, and which can be extended to the loop braid groups (see [BS20]). Namely, the *Tong-Yang-Ma* representation $\mathbf{TYM}_n: \mathbf{B}_n \rightarrow GL_n(\mathbb{Z}[t^{\pm 1}])$ is defined by:

$$\sigma_i \mapsto Id_{i-1} \oplus \begin{bmatrix} 0 & t \\ 1 & 0 \end{bmatrix} \oplus Id_{n-i-1}$$

for all non-negative integers $i \in \{1, \dots, n-1\}$. We prove below that it is possible to reconstruct \mathbf{TYM}_n using the slight generalisation of the iterative homological construction (6.1).

First of all, fixing $\mathbf{F}_n = \langle x_1, \dots, x_n \rangle$, we consider the group morphism $a_{n,2}: \mathbf{B}_n \rightarrow \mathbf{Aut}(\mathbf{F}_n)$ defined for all $i \in \{1, \dots, n-1\}$ by $a_{n,2}(\sigma_i)(x_i) = x_{i+1}$, $a_{n,2}(\sigma_i)(x_{i+1}) = x_i^{-1}$, $a_{n,2}(\sigma_i)(x_j) = x_j$ for $j \notin \{i, i+1\}$. It belongs to the family of local representations of \mathbf{B}_n in $\mathbf{Aut}(\mathbf{F}_n)$ classified in [Wad92; Ito13]: more precisely, it corresponds to the Type 2 *Wada representation*.

For simplicity, for each non-negative integer n , we denote by \mathbb{D}_n the n -th punctured 2-disc (i.e. we remove the marked points of of the decorated surface of Notation 4.28) with the basepoint

$p_0 \in \partial \mathbb{D}^2$. Note that \mathbb{D}_n is an Eilenberg–MacLane space of type $K(\pi_1(\mathbb{D}_n), 1)$. Let $\mathbf{hAut}(\mathbb{D}_n)$ be the grouplike topological monoid of based homotopy self-equivalences of \mathbb{D}_n . In particular, $\mathbf{Aut}(\pi_1(\mathbb{D}_n))$ is naturally isomorphic to $\pi_0(\mathbf{hAut}(\mathbb{D}_n))$: $a_{n,2}$ thus defines a morphism $\mathbf{B}_n \rightarrow \pi_0(\mathbf{hAut}(\mathbb{D}_n))$. Hence we define a functor $A_2: \beta \rightarrow \pi_0(\mathbf{Cov}_\bullet)$ that sends each $n \in \text{ob}(\beta)$ to \mathbb{D}_n together with its universal cover $\tilde{\mathbb{D}}_n$ for objects, and by $a_{n,2}$ for the automorphisms of β .

Using the generalisation (6.1) of the iterative construction, we consider the functor $L_1^r(A_2; -)$. Let $\rho: \beta_{A_2} \rightarrow \text{Mod}_\bullet$ be the functor defined by sending each n to the group ring $\mathbb{Z}[t^{\pm 1}]$, the multiplication by $-t^{-1}$ for each generator of $\pi_1(\mathbb{D}_n)$ and the trivial action of the braid group \mathbf{B}_n . Then a straightforward computation shows that:

Proposition 6.4 *The functor $L_1^r(A_2; \rho)$ encodes the Tong–Yang–Ma representations.*

Remark 6.5 Taking the tensor product $\mathbb{C} \otimes \mathbb{Z}[t^{\pm 1}] \cong \mathbb{C}[t^{\pm 1}]$ and specializing t to complex values in the Burau and Tong–Yang–Ma representations, we obtain complex representations of \mathbf{B}_n . The classification of the irreducible complex representations of \mathbf{B}_n is well researched for various degrees. For $n \geq 7$, [For96] proves that the irreducible complex representations of \mathbf{B}_n of degree $\leq n - 1$ are either one-dimensional representation or a tensor product of a one-dimensional representation and a composition factor of the specialization of the Burau representation, and [Sys01] shows that those of degree n are equivalent to a tensor product of one-dimensional representation and specialization of the Tong–Yang–Ma representation if $n \geq 9$. Finally, it is shown in [Sys20] that there are no irreducible complex representations of degree $n + 1$ for $n \geq 10$. Hence all the low-dimensional complex irreducible representations of braid groups are homological.

7. Reduced Burau representations for loop braid groups

We consider the homological representations of Proposition 5.40 for $i = k = 1$, namely

$$\mathfrak{L}_1(1, \mathcal{L}\beta): \mathfrak{UL}\beta \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}]} \quad \text{and} \quad \mathfrak{L}_1(1, \mathcal{L}\beta'): \mathfrak{UL}\beta' \longrightarrow \text{Mod}_{\mathbb{Z}[\mathbb{Z}/2\mathbb{Z}]}.$$

Our goal is to give explicit formulas for the matrices of these representations, restricted to the non-extended loop braid groups \mathbf{LB}_n and the extended loop braid groups \mathbf{LB}'_n respectively. In the case of \mathbf{LB}_n , the representation is a *free* $\mathbb{Z}[\mathbb{Z}]$ -module, and we will give formulas with respect to a particular basis of this free module. In the case of \mathbf{LB}'_n , the representation is *not* a free $\mathbb{Z}[\mathbb{Z}/2\mathbb{Z}]$ -module, although it is not far from being free: it is the direct sum of a free module with the trivial $\mathbb{Z}[\mathbb{Z}/2\mathbb{Z}]$ -module \mathbb{Z} . In this case, we will give formulas with respect to a generating set given by a basis for the free summand together with a chosen generator of the (non-free) trivial \mathbb{Z} summand.

We first give a more concrete description of the homological representations. Fix once and for all an n -component unlink in the interior of the closed 3-disc and write \mathbb{D}_n^3 for its complement. The extended loop braid group \mathbf{LB}'_n (and hence also its subgroup \mathbf{LB}_n) acts up to homotopy on \mathbb{D}_n^3 by homeomorphisms. The homological representations of \mathbf{LB}_n and \mathbf{LB}'_n that we are considering are their induced actions on the first homology of \mathbb{D}_n^3 with coefficients in a certain local system over the ring $\mathbb{Z}[\mathbb{Z}]$ respectively $\mathbb{Z}[\mathbb{Z}/2\mathbb{Z}]$. These local systems arise from regular covering spaces of \mathbb{D}_n^3 with deck transformation group \mathbb{Z} respectively $\mathbb{Z}/2\mathbb{Z}$. By Shapiro’s lemma, we may therefore equivalently view the homological representations as the induced action on the first (untwisted, integral) homology of the respective covering spaces.

Let us make these two covering spaces of \mathbb{D}_n^3 more precise. First, recall that the fundamental group of \mathbb{D}_n^3 is the free group F_n on n generators. This is easy to see: the unlink-complement $\mathbb{D}_n^3 \subseteq \mathbb{D}^3$ deformation retracts onto a wedge of n circles and n copies of the 2-sphere. The n circles a_1, \dots, a_n are shown in Figure 7.1. Now let

$$\phi: \pi_1(\mathbb{D}_n^3) \longrightarrow \mathbb{Z}$$

be the surjective homomorphism defined by $\phi(a_i) = 1$ for all $i = 1, \dots, n$. Let

$$\phi': \pi_1(\mathbb{D}_n^3) \longrightarrow \mathbb{Z}/2\mathbb{Z}$$




Figure 7.1 The unlink-complement \mathbb{D}_n^3 with generators a_1, \dots, a_n for $\pi_1(\mathbb{D}_n^3) \cong F_n$.

be the composition of ϕ with the unique surjection $\mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z}$. We denote by $\widetilde{\mathbb{D}}_n^3$ the regular covering space corresponding to $\ker(\phi)$ and denote by $\widehat{\mathbb{D}}_n^3$ the regular covering space corresponding to $\ker(\phi')$. We therefore have coverings

$$\widetilde{\mathbb{D}}_n^3 \longrightarrow \widehat{\mathbb{D}}_n^3 \longrightarrow \mathbb{D}_n^3$$

and we are considering the action of \mathbf{LB}_n on $H_1(\widetilde{\mathbb{D}}_n^3)$ and the action of \mathbf{LB}'_n on $H_1(\widehat{\mathbb{D}}_n^3)$.

We can make these covering spaces even more precise by building explicit models for each of them. We embed n pairwise disjoint closed 3-discs into the interior of the unit 3-disc \mathbb{D}^3 as pictured in Figure 7.2, so that each little 3-disc looks like a “lens shape” and the union of their equators is precisely the n -component unlink that we fixed earlier. Let $\mathring{\mathbb{D}}_n^3$ denote \mathbb{D}_n^3 minus the interiors of these n little 3-discs, equivalently, \mathbb{D}^3 minus the interiors and equators of the n little 3-discs. Also, write N_i for the open northern hemisphere of the boundary of the i -th little 3-disc, and write S_i for the open southern hemisphere of the boundary of the i -th little 3-disc. Now consider

$$\mathbb{Z} \times \mathring{\mathbb{D}}_n^3$$

and glue $\{j\} \times N_i$ to $\{j-1\} \times S_i$ via the homeomorphism $N_i \cong S_i$ given by reflection in the plane passing through the equator. This is an explicit model for $\widetilde{\mathbb{D}}_n^3$. Similarly, we may consider

$$\mathbb{Z}/2\mathbb{Z} \times \mathring{\mathbb{D}}_n^3$$

and glue as before, where j is now considered mod 2. This is an explicit model for $\widehat{\mathbb{D}}_n^3$.

Matrices for non-extended loop braid groups. We first consider the action of the non-extended loop braid group \mathbf{LB}_n on $H_1(\widetilde{\mathbb{D}}_n^3)$. As noted above, the unlink-complement \mathbb{D}_n^3 deformation retracts onto a wedge of n circles and n copies of the 2-sphere. This deformation retraction lifts to a deformation retraction of the covering space $\widetilde{\mathbb{D}}_n^3$ onto the space pictured in Figure 7.3. This is an infinite 2-dimensional cell complex with vertices indexed by \mathbb{Z} , with exactly n edges between consecutive vertices (and none between non-consecutive vertices) and with exactly n copies of the 2-sphere wedged onto each vertex. Its fundamental group is freely generated by $t^k \cdot (a_2 \bar{a}_1), \dots, t^k \cdot (a_n \bar{a}_{n-1})$ for all $k \in \mathbb{Z}$, where \bar{a} denotes the reverse of a path a . Abelianising and writing $\mathbb{Z}[\mathbb{Z}] = \mathbb{Z}[t^{\pm 1}]$, we see that its first homology is freely generated, as a $\mathbb{Z}[t^{\pm 1}]$ -module, by $x_1 := a_2 \bar{a}_1, \dots, x_{n-1} := a_n \bar{a}_{n-1}$. This choice of free generating set determines an isomorphism of modules $H_1(\widetilde{\mathbb{D}}_n^3) \cong \mathbb{Z}[t^{\pm 1}]^{\oplus n-1}$, so we may write the action of \mathbf{LB}_n on $H_1(\widetilde{\mathbb{D}}_n^3)$ as

$$\mathbf{LB}_n \longrightarrow GL_{n-1}(\mathbb{Z}[t^{\pm 1}]).$$

Figure 7.4 describes, geometrically, the action of the generators σ_i and τ_i of \mathbf{LB}_n on the elements x_1, \dots, x_{n-1} , projected to the base space \mathbb{D}_n^3 . Ignore for the moment the bottom row corresponding




Figure 7.2 The complement $\hat{\mathbb{D}}_n^3$ of the interiors and equators of n closed little 3-discs (“lens shapes”) in the interior of the closed unit 3-disc \mathbb{D}^3 . The boundary of $\hat{\mathbb{D}}_n^3$ decomposes as the disjoint union of $2n + 1$ components: $\partial\hat{\mathbb{D}}_n^3 = \partial\mathbb{D}^3 \sqcup N_1 \sqcup \dots \sqcup N_n \sqcup S_1 \sqcup \dots \sqcup S_n$.




Figure 7.3 A deformation retract of the \mathbb{Z} -covering $\hat{\mathbb{D}}_n^3$. The edges are all oriented from left to right.

to ρ_i . Writing the loops on the right-hand side in terms of the generators x_1, \dots, x_{n-1} and the action of t , we see that σ_i acts on the generators (x_{i-1}, x_i, x_{i+1}) by the matrix

$$\begin{bmatrix} 1 & 0 & 0 \\ 1 & -1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

and trivially on all other generators, and τ_i acts on the generators (x_{i-1}, x_i, x_{i+1}) by the matrix

$$\begin{bmatrix} 1 & 0 & 0 \\ t & -t & 1 \\ 0 & 0 & 1 \end{bmatrix}$$

and trivially on all other generators. When $i = 1$ or $i = n - 1$ we pass to the appropriate 2×2 submatrices. These matrices give a purely algebraic description of the homological representation $\mathcal{L}_1(1, \mathcal{L}\beta)(n)$ of \mathbf{LB}_n .

Matrices for extended loop braid groups. Similarly to above, the deformation retraction of the unlink-complement \mathbb{D}_n^3 onto a wedge of circles and 2-spheres lifts to a deformation retraction of its covering $\hat{\mathbb{D}}_n^3$ onto the space pictured in Figure 7.5. This is a finite 2-dimensional cell complex with two vertices $\{v, tv\}$, exactly $2n$ edges between them and with exactly n copies of the 2-sphere wedged onto each vertex. Its fundamental group is freely generated by the $2n - 1$ loops

$$a_2\bar{a}_1, \dots, a_n\bar{a}_{n-1}, t.(a_2\bar{a}_1), \dots, t.(a_n\bar{a}_{n-1}), a_n(t.a_n),$$

where \bar{a} denotes the reverse of a path a . Hence its first homology $H_1(\hat{\mathbb{D}}_n^3)$ is generated, as an abelian group, by the same $2n - 1$ loops, viewed as homology classes. The first $2n - 2$ of these classes generate a summand isomorphic to $\mathbb{Z}[\mathbb{Z}/2\mathbb{Z}]^{\oplus n-1}$ as a $\mathbb{Z}[\mathbb{Z}/2\mathbb{Z}]$ -module. Here we are identifying $\mathbb{Z}[\mathbb{Z}/2\mathbb{Z}]$ with $R := \mathbb{Z}[t^{\pm 1}]/(t^2 - 1)$. The last element generates a summand isomorphic to \mathbb{Z} viewed as a *trivial* $\mathbb{Z}[\mathbb{Z}/2\mathbb{Z}]$ -module. Note that this is *torsion* as an R -module (although it is of




Figure 7.4 The action of \mathbf{LB}'_n on the loops x_1, \dots, x_{n-1} .




Figure 7.5 A deformation retract of the double covering $\widehat{\mathbb{D}}_n^3$. The top n edges are oriented from left to right; the bottom n edges are oriented from right to left.

course free as an abelian group), since it is isomorphic to $R/(t-1)$. Let us write $x_i := a_{i+1}\bar{a}_i$ for $i = 1, \dots, n-1$ and $y := a_n(t.a_n)$. These elements determine an isomorphism of modules $H_1(\widehat{\mathbb{D}}_n^3) \cong R^{\oplus n-1} \oplus R/(t-1)$, so we may write the action of \mathbf{LB}'_n on $H_1(\widehat{\mathbb{D}}_n^3)$ as

$$\mathbf{LB}'_n \longrightarrow \text{Aut}_R(R^{\oplus n-1} \oplus R/(t-1)).$$

To give a purely algebraic description of the homological representation $\mathfrak{L}_1(1, \mathcal{L}\beta')(n)$, we therefore just have to calculate the $n \times n$ matrices that describe the action of each generator σ_i , τ_i and ρ_i of \mathbf{LB}'_n on the generators x_1, \dots, x_{n-1}, y .

In order to do this, we first consider a variant of $\mathfrak{L}_1(1, \mathcal{L}\beta')(n)$, which contains it as a subrepresentation and is easier to calculate. The action of \mathbf{LB}'_n on the unlink-complement \mathbb{D}_n^3 fixes its boundary $\partial\mathbb{D}^3$ pointwise, so in particular it fixes any given basepoint in the boundary. We may arrange that the embedded wedge of circles and 2-spheres $(S^1 \vee S^2)^{\vee n} \hookrightarrow \mathbb{D}_n^3$ onto which \mathbb{D}_n^3 deformation retracts has its unique vertex v on the boundary. The lifted action of \mathbf{LB}'_n on the corresponding deformation retract (see Figure 7.5) of the covering $\widehat{\mathbb{D}}_n^3$ therefore fixes the set of vertices $\{v, tv\}$ setwise, so we may consider its induced action on *relative* first homology $H_1(\widehat{\mathbb{D}}_n^3, \{v, tv\})$. Note

matrix for ρ_1 is

$$\begin{bmatrix} -t & 0 & \cdots & 0 \\ -1-t & & & \\ \vdots & & \mathbf{I}_{n-1} & \\ -1-t & & & \\ 1 & & & \end{bmatrix},$$

the matrix for ρ_i with $2 \leq i \leq n-1$ is

$$\mathbf{I}_{i-2} \oplus \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ 1+t & -t & 0 & \cdots & 0 \\ 1+t & -1-t & & & \\ \vdots & \vdots & & \mathbf{I}_{n-i} & \\ 1+t & -1-t & & & \\ -1 & 1 & & & \end{bmatrix}$$

and the matrix for ρ_n is $\mathbf{I}_{n-2} \oplus \begin{bmatrix} 1 & 0 \\ -1 & -1 \end{bmatrix}$. Note that, since these matrices describe automorphisms of $R^{\oplus n-1} \oplus R/(t-1)$, each entry above the bottom row should be considered as an element of R , whereas each element of the bottom row should be considered as an element of $R/(t-1) \cong \mathbb{Z}$.

How these are related to each other. The action of \mathbf{LB}_n on $H_1(\widehat{\mathbb{D}}_n^3) \cong \mathbb{Z}[t^{\pm 1}]^{\oplus n-1}$ that we described earlier has been defined before, via assigning explicit matrices to the generators σ_i and τ_i , by Vershinin [Ver01, §4]. More precisely, it is the reduced version of the representation described there via explicit matrices. In particular, its restriction to the classical braid group, which embeds into \mathbf{LB}_n as the subgroup generated by the τ_i generators, is exactly the classical reduced Burau representation. We may therefore call the representation

$$\mathbf{LB}_n \longrightarrow \text{Aut}_{\mathbb{Z}[t^{\pm 1}]}(H_1(\widehat{\mathbb{D}}_n^3)) = GL_{n-1}(\mathbb{Z}[t^{\pm 1}])$$

the *reduced Burau representation of non-extended loop braid groups*. The map of covering spaces $\widehat{\mathbb{D}}_n^3 \longrightarrow \widehat{\mathbb{D}}_n^3$ induces a map of \mathbf{LB}_n -representations

$$H_1(\widehat{\mathbb{D}}_n^3) = \mathbb{Z}[t^{\pm 1}]^{\oplus n-1} \longrightarrow R^{\oplus n-1} \oplus R/(t-1) = H_1(\widehat{\mathbb{D}}_n^3),$$

where $R = \mathbb{Z}[t^{\pm 1}]/(t^2 - 1)$. Checking generators, we see that it is the direct sum of $n-1$ copies of the obvious projection $\mathbb{Z}[t^{\pm 1}] \rightarrow R$ followed by the inclusion of the left-hand summand $R^{\oplus n-1}$ of the codomain. This means that if we consider the reduced Burau representation of $\mathbf{LB}_n \bmod t^2$ (in other words, we tensor it over $\mathbb{Z}[t^{\pm 1}]$ with R), it is isomorphic to the direct summand $R^{\oplus n-1}$ of the representation $H_1(\widehat{\mathbb{D}}_n^3)$ of \mathbf{LB}'_n . Another, more concrete way of verifying this statement is to notice that, if we forget the generator y from the matrices above for σ_i and τ_i acting on $H_1(\widehat{\mathbb{D}}_n^3)$, they agree with the ones that we found for $H_1(\mathbb{D}_n^3)$ earlier. Note, however, that the action of ρ_i does *not* send the summand $R^{\oplus n-1}$ to itself, so this is only a sub- \mathbf{LB}_n -representation, not a sub- \mathbf{LB}'_n -representation. To see that ρ_i does not send $R^{\oplus n-1}$ to itself, one may either read this off from the explicit matrices that we obtained for ρ_i above, or one may see this geometrically from Figure 7.4, where we note that the loops $\rho_j(x_i)$ *cannot in general be expressed purely in terms of the generators* x_1, \dots, x_{n-1} ; the generator y is also needed.

We conclude from this that, in order to extend the reduced Burau representation from \mathbf{LB}_n to \mathbf{LB}'_n , we have to first consider it mod t^2 (tensor with R), then enlarge it from $R^{\oplus n-1}$ to $R^{\oplus n-1} \oplus R/(t-1)$ (extending the matrices for σ_i and τ_i as shown above) and finally define the action of ρ_i using the matrices pictured above. Notice that it is not trivial to extend the action of the σ_i and τ_i generators to the new $R/(t-1)$ summand: σ_{n-1} sends the new generator y to $y + (1+t).x_{n-1}$ and τ_{n-1} sends y to $y - (1+t).x_{n-1}$. We call the representation

$$\mathbf{LB}'_n \longrightarrow \text{Aut}_R(H_1(\widehat{\mathbb{D}}_n^3)) = \text{Aut}_R(R^{\oplus n-1} \oplus R/(t-1))$$

the *reduced Burau representation of extended loop-braid groups*.

Summary. The above representations may be summarised as follows:

$$\begin{array}{ccccccc}
x_1, \dots, x_{n-1} & & x_1, \dots, x_{n-1} & & x_1, \dots, x_{n-1}, y & & a_1, \dots, a_n \\
\mathbb{Z}[\mathbb{Z}]^{\oplus n-1} & \xrightarrow{\quad} & R^{\oplus n-1} & \hookrightarrow & R^{\oplus n-1} \oplus R/(t-1) & \hookrightarrow & R^{\oplus n} \\
\parallel & & \parallel & & \parallel & & \parallel \\
H_1(\widehat{\mathbb{D}}_n^3) & & H_1(\widetilde{\mathbb{D}}_n^3) \otimes_{\mathbb{Z}[\mathbb{Z}]} R & & H_1(\widehat{\mathbb{D}}_n^3) & & H_1(\widehat{\mathbb{D}}_n^3, \{v, tv\})
\end{array} \tag{7.2}$$

where:

- a generating set corresponding to the given direct sum decomposition is written in blue above each module,
- the left two modules are \mathbf{LB}_n -representations and the right two modules are \mathbf{LB}'_n -representations,
- all maps are \mathbf{LB}_n -equivariant and the right-hand map is moreover \mathbf{LB}'_n -equivariant,
- the ground ring on the left is $\mathbb{Z}[\mathbb{Z}] = \mathbb{Z}[t^{\pm 1}]$ and the ground ring for the other three modules is $R = \mathbb{Z}[\mathbb{Z}/2\mathbb{Z}] = \mathbb{Z}[t^{\pm 1}]/(t^2 - 1)$.

These representations are named, respectively (from left to right):

- the reduced Burau representation of \mathbf{LB}_n ,
- the reduction mod t^2 of the above,
- the reduced Burau representation of \mathbf{LB}'_n ,
- the unreduced Burau representation of \mathbf{LB}'_n .

References

- [AK10] B. H. An and K. H. Ko. *A family of representations of braid groups on surfaces*. *Pacific J. Math.* 247.2 (2010), pp. 257–282 (↑ 6, 60, 61, 67).
- [AP20] C. A.-M. Anghel and M. Palmer. *Lawrence-Bigelow representations, bases and duality*. ArXiv:2011.02388. 2020 (↑ 16, 44, 67).
- [BB05] J. S. Birman and T. E. Brendle. *Braids: a survey*. *Handbook of knot theory* (2005), pp. 19–103 (↑ 1).
- [BC18] F. Ben Aribi and A. Conway. *L^2 -Burau maps and L^2 -Alexander torsions*. *Osaka J. Math.* 55.3 (2018), pp. 529–545 (↑ 14).
- [Bel04] P. Bellingeri. *On presentations of surface braid groups*. *J. Algebra* 274.2 (2004), pp. 543–563 (↑ 66).
- [BGG08] P. Bellingeri, S. Gervais and J. Guaschi. *Lower central series of Artin-Tits and surface braid groups*. *J. Algebra* 319.4 (2008), pp. 1409–1427 (↑ 66).
- [BGG11] P. Bellingeri, E. Godelle and J. Guaschi. *Exact sequences, lower central series and representations of surface braid groups*. 2011 (↑ 7).
- [BGG17] P. Bellingeri, E. Godelle and J. Guaschi. *Abelian and metabelian quotient groups of surface braid groups*. *Glasg. Math. J.* 59.1 (2017), pp. 119–142 (↑ 6, 59, 60, 66, 69).
- [Big01] S. J. Bigelow. *Braid groups are linear*. *J. Amer. Math. Soc.* 14.2 (2001), pp. 471–486 (↑ 2, 8, 57, 58).
- [Big02] S. Bigelow. *Representations of braid groups*. *Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002)*. Higher Ed. Press, Beijing, 2002, pp. 37–45 (↑ 59).
- [Big03] S. Bigelow. *The Lawrence-Krammer representation*. *Topology and geometry of manifolds (Athens, GA, 2001)*. Vol. 71. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 2003, pp. 51–68 (↑ 58).
- [Big04] S. Bigelow. *Homological representations of the Iwahori-Hecke algebra*. *Proceedings of the Casson Fest*. Vol. 7. Geom. Topol. Monogr. Geom. Topol. Publ., Coventry, 2004, pp. 493–507 (↑ 5, 57, 58, 67).
- [Bir74] J. S. Birman. *Braids, links, and mapping class groups*. Annals of Mathematics Studies, No. 82. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1974, pp. ix+228 (↑ 66).
- [Bre97] G. E. Bredon. *Sheaf theory*. Second. Vol. 170. Graduate Texts in Mathematics. Springer-Verlag, New York, 1997, pp. xii+502 (↑ 67).
- [BS20] P. Bellingeri and A. Soulié. *A note on representations of welded braid groups*. preprint arXiv:2001.04272, submitted (2020) (↑ 62, 71).
- [Bud05] R. D. Budney. *On the image of the Lawrence-Krammer representation*. *J. Knot Theory Ramifications* 14.6 (2005), pp. 773–789 (↑ 57).

- [Bur35] W. Burau. *Über Zopfgruppen und gleichsinnig verdrillte Verkettungen. Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg*. Vol. 11. 1. Springer. 1935, pp. 179–186 (↑ [2](#), [7](#), [57](#)).
- [Cer61] J. Cerf. *Topologie de certains espaces de plongements*. *Bull. Soc. Math. France* 89 (1961), pp. 227–380 (↑ [29](#), [31](#), [41](#), [42](#), [48](#)).
- [Chi12] L. R. Childers. *Simply intersecting pair maps in the mapping class group*. *J. Knot Theory Ramifications* 21.11 (2012), pp. 1250107, 21 (↑ [67](#)).
- [Chi72] D. R. J. Chillingworth. *Winding numbers on surfaces. I*. *Math. Ann.* 196 (1972), pp. 218–249 (↑ [67](#)).
- [CO98] T. D. Cochran and K. E. Orr. *Stability of lower central series of compact 3-manifold groups*. *Topology* 37.3 (1998), pp. 497–526 (↑ [52](#)).
- [Con18] A. Conway. *Braid maps and twisted Alexander polynomials*. *Proc. Edinb. Math. Soc. (2)* 61.2 (2018), pp. 479–497 (↑ [14](#)).
- [Dam17] C. Damiani. *A journey through loop braid groups*. *Expo. Math.* 35.3 (2017), pp. 252–285 (↑ [41](#), [43](#)).
- [DK01] J. F. Davis and P. Kirk. *Lecture notes in algebraic topology*. Vol. 35. Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2001, pp. xvi+367 (↑ [16](#)).
- [DPS21] J. Darné, M. Palmer and A. Soulié. *When the lower central series stops. Preprint in progress, to be posted on the arXiv soon*. (2021) (↑ [3](#), [5](#), [6](#), [52](#), [55–59](#), [61–63](#), [68](#), [69](#)).
- [EN02] K. Erdmann and D. K. Nakano. *Representation type of Hecke algebras of type A*. *Trans. Amer. Math. Soc.* 354.1 (2002), pp. 275–285 (↑ [1](#), [8](#)).
- [For96] E. Formanek. *Braid group representations of low degree*. *Proc. London Math. Soc. (3)* 73.2 (1996), pp. 279–322 (↑ [72](#)).
- [FP92] E. Formanek and C. Procesi. *The automorphism group of a free group is not linear*. *J. Algebra* 149.2 (1992), pp. 494–499 (↑ [8](#)).
- [GG09] D. L. Gonçalves and J. Guaschi. *The lower central and derived series of the braid groups of the sphere*. *Trans. Amer. Math. Soc.* 361.7 (2009), pp. 3375–3399 (↑ [68](#)).
- [Gra73] A. Gramain. *Le type d’homotopie du groupe des difféomorphismes d’une surface compacte*. *Annales scientifiques de l’École Normale Supérieure*. Vol. 6. 1. Elsevier. 1973, pp. 53–66 (↑ [42](#)).
- [Gra76] D. Grayson. *Higher algebraic K-theory. II (after Daniel Quillen)* (1976), 217–240. Lecture Notes in Math., Vol. 551 (↑ [22](#)).
- [Hat02] A. Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002, pp. xii+544 (↑ [57](#), [71](#)).
- [Hir76] M. W. Hirsch. *Differential topology*. Graduate Texts in Mathematics, No. 33. New York: Springer-Verlag, 1976, pp. x+221 (↑ [26](#)).
- [Ito13] T. Ito. *The classification of Wada-type representations of braid groups*. *J. Pure Appl. Algebra* 217.9 (2013), pp. 1754–1763 (↑ [71](#)).
- [Joh83] D. Johnson. *The structure of the Torelli group. I. A finite set of generators for \mathcal{I}* . *Ann. of Math. (2)* 118.3 (1983), pp. 423–442 (↑ [67](#)).
- [Koh17] T. Kohno. *Quantum representations of braid groups and holonomy Lie algebras. Geometry, dynamics, and foliations 2013*. Vol. 72. Adv. Stud. Pure Math. Math. Soc. Japan, Tokyo, 2017, pp. 117–144 (↑ [44](#)).
- [Kor02] M. Korkmaz. *Low-dimensional homology groups of mapping class groups: a survey*. *Turkish J. Math.* 26.1 (2002), pp. 101–114 (↑ [69](#)).
- [Kra02] D. Kramer. *Braid groups are linear*. *Ann. of Math. (2)* 155.1 (2002), pp. 131–156 (↑ [2](#), [8](#), [57](#), [58](#)).
- [Kra19] M. Krannich. *Homological stability of topological moduli spaces*. *Geom. Topol.* 23.5 (2019), pp. 2397–2474 (↑ [22](#)).
- [KS80] S. A. Krugljak and J. S. Samoilenko. *Unitary equivalence of sets of selfadjoint operators*. *Funktsional. Anal. i Prilozhen.* 14.1 (1980), pp. 60–62 (↑ [1](#)).
- [Law90] R. J. Lawrence. *Homological representations of the Hecke algebra*. *Comm. Math. Phys.* 135.1 (1990), pp. 141–191 (↑ [2](#), [5](#), [57](#)).
- [Lim63] E. L. Lima. *On the local triviality of the restriction map for embeddings*. *Comment. Math. Helv.* 38 (1963), pp. 163–164 (↑ [29](#), [41](#)).
- [Lon94] D. D. Long. *Constructing representations of braid groups*. *Comm. Anal. Geom.* 2.2 (1994), pp. 217–238 (↑ [7](#), [70](#)).
- [Mac98] S. Mac Lane. *Categories for the working mathematician*. Second. Vol. 5. Graduate Texts in Mathematics. Springer-Verlag, New York, 1998, pp. xii+314 (↑ [21](#), [41](#)).

- [Mar19] D. Margalit. *Problems, questions, and conjectures about mapping class groups. Breadth in contemporary topology*. Vol. 102. Proc. Sympos. Pure Math. Amer. Math. Soc., Providence, RI, 2019, pp. 157–186 (↑ 1).
- [MKS04] W. Magnus, A. Karrass and D. Solitar. *Combinatorial group theory*. second. Presentations of groups in terms of generators and relations. Dover Publications, Inc., Mineola, NY, 2004, pp. xii+444 (↑ 66).
- [MMSS01] M. A. Mandell, J. P. May, S. Schwede and B. Shipley. *Model categories of diagram spectra*. *Proc. London Math. Soc. (3)* 82.2 (2001), pp. 441–512 (↑ 30).
- [Mor07] T. Moriyama. *The mapping class group action on the homology of the configuration spaces of surfaces*. *J. Lond. Math. Soc. (2)* 76.2 (2007), pp. 451–466 (↑ 7, 67).
- [Mor89] S. Morita. *Families of Jacobian manifolds and characteristic classes of surface bundles. I*. *Ann. Inst. Fourier (Grenoble)* 39.3 (1989), pp. 777–810 (↑ 67).
- [Pal18a] M. Palmer. *Homological stability for moduli spaces of disconnected submanifolds, I*. ArXiv:1805.03917. 2018. To appear in Algebraic & Geometric Topology (↑ 36, 37, 45, 48).
- [Pal18b] M. Palmer. *Twisted homological stability for configuration spaces*. *Homology, Homotopy and Applications* 20.2 (2018), pp. 145–178. {arxiv:1308.4397} (↑ 16).
- [Pal60a] R. S. Palais. *Extending diffeomorphisms*. *Proc. Amer. Math. Soc.* 11 (1960), pp. 274–277 (↑ 26).
- [Pal60b] R. S. Palais. *Local triviality of the restriction map for embeddings*. *Comment. Math. Helv.* 34 (1960), pp. 305–312 (↑ 29, 31, 36, 37, 41, 45).
- [PP02] L. Paoluzzi and L. Paris. *A note on the Lawrence-Krammer-Bigelow representation*. *Algebr. Geom. Topol.* 2 (2002), pp. 499–518 (↑ 58).
- [PS21] M. Palmer and A. Soulié. *Irreducibility and polynomiality of motion group and mapping class group representations*. Preprint in progress, to be posted on the arXiv soon. (2021) (↑ 2, 8, 52, 60, 67).
- [RW17] O. Randal-Williams and N. Wahl. *Homological stability for automorphism groups*. *Adv. Math.* 318 (2017), pp. 534–626. {arxiv:1409.3541} (↑ 22, 34, 40).
- [Sak12] T. Sakasai. *A survey of Magnus representations for mapping class groups and homology cobordisms of surfaces*. *Handbook of Teichmüller theory. Volume III*. Vol. 17. IRMA Lect. Math. Theor. Phys. Eur. Math. Soc., Zürich, 2012, pp. 531–594 (↑ 66).
- [Sma59] S. Smale. *Diffeomorphisms of the 2-sphere*. *Proc. Amer. Math. Soc.* 10 (1959), pp. 621–626 (↑ 41).
- [Sou19a] A. Soulié. *Generalized Long-Moody functors*. preprint arXiv:1709.04278, submitted (2019) (↑ 7, 8, 70, 71).
- [Sou19b] A. Soulié. *The Long-Moody construction and polynomial functors*. *Ann. Inst. Fourier (Grenoble)* 69.3 (2019), pp. 1799–1856 (↑ 7, 70).
- [Ste51] N. Steenrod. *The Topology of Fibre Bundles*. Princeton Mathematical Series, vol. 14. Princeton University Press, Princeton, N. J., 1951, pp. viii+224 (↑ 12).
- [Stu10] M. Stukow. *Generating mapping class groups of nonorientable surfaces with boundary*. *Adv. Geom.* 10.2 (2010), pp. 249–273 (↑ 69).
- [Suz05] M. Suzuki. *Geometric interpretation of the Magnus representation of the mapping class group*. *Kobe J. Math.* 22.1-2 (2005), pp. 39–47 (↑ 7, 66).
- [Sys01] I. Sysoeva. *Dimension n representations of the braid group on n strings*. *J. Algebra* 243.2 (2001), pp. 518–538 (↑ 72).
- [Sys20] I. Sysoeva. *Irreducible representations of Braid Group B_n of dimension $n + 1$* . preprint, arXiv:math2005.03105 (2020) (↑ 72).
- [TV08] B. Toën and G. Vezzosi. *Homotopical algebraic geometry. II. Geometric stacks and applications*. *Mem. Amer. Math. Soc.* 193.902 (2008), pp. x+224 (↑ 30).
- [TYM96] D.-M. Tong, S.-D. Yang and Z.-Q. Ma. *A new class of representations of braid groups*. *Comm. Theoret. Phys.* 26.4 (1996), pp. 483–486 (↑ 71).
- [Ver01] V. V. Vershinin. *On homology of virtual braids and Burau representation*. Vol. 10. 5. Knots in Hellas '98, Vol. 3 (Delphi). 2001, pp. 795–812 (↑ 6, 62, 77).
- [Wad92] M. Wada. *Group invariants of links*. *Topology* 31.2 (1992), pp. 399–406 (↑ 71).

Martin Palmer, *Institutul de Matematică Simion Stoilow al Academiei Române, 21 Calea Grivitei, 010702 București, România*. Email address: mpanghel@imar.ro

Arthur Soulié, *University of Glasgow, School of Mathematics and Statistics, 132 University Pl, Glasgow G12 8TA, United-Kingdom*. Email address: Arthur.Soulie@glasgow.ac.uk