


Aufgaben zur Topologie

Prof. Dr. C.-F. Bödigheimer

Wintersemester 2016/17

Week 8 — Homological algebra and homotopy invariance of homology

Due: 21. December 2016



The birth of homology, from *Analysis Situs*, H. Poincaré (1895).

Exercise 8.1 (The five-lemma.)

Prove the famous *five-lemma*. Let R be a ring and suppose we have the following diagram of modules over R :

$$\begin{array}{ccccccccc}
 A & \longrightarrow & B & \longrightarrow & C & \longrightarrow & D & \longrightarrow & E \\
 \downarrow \alpha & & \downarrow \beta & & \downarrow \gamma & & \downarrow \delta & & \downarrow \varepsilon \\
 A' & \longrightarrow & B' & \longrightarrow & C' & \longrightarrow & D' & \longrightarrow & E'
 \end{array}$$

Assume that this diagram is commutative and that the two horizontal rows of homomorphisms are exact. Moreover, assume that α is surjective, β and δ are bijective and ε is injective. Prove that γ is bijective.

Application: let $\phi: B \rightarrow B'$ be a homomorphism of R -modules taking a submodule $A \subseteq B$ to a submodule $A' \subseteq B'$, so that we have restricted and induced homomorphisms $\phi|_A: A \rightarrow A'$ and $\bar{\phi}: B/A \rightarrow B'/A'$. If $\phi|_A$ and $\bar{\phi}$ are both isomorphisms then so is ϕ .

Exercise 8.2 (Mapping cones and mapping cylinders of chain complexes.)

Let A and B be chain complexes with differential ∂_A resp. ∂_B and let $f: A \rightarrow B$ be a chain map.

(i) Define a new chain complex $\text{Cone}(f)$ by $\text{Cone}(f)_n = A_{n-1} \oplus B_n$ and setting its differential $\partial: A_{n-1} \oplus B_n \rightarrow A_{n-2} \oplus B_{n-1}$ to be the sum of the four maps

$$\partial_A: A_{n-1} \rightarrow A_{n-2} \quad \partial_B: B_n \rightarrow B_{n-1} \quad 0: B_n \rightarrow A_{n-2} \quad (-1)^n \cdot f_{n-1}: A_{n-1} \rightarrow B_{n-1},$$

or as a formula

$$\partial(a, b) := (\partial_A(a), (-1)^n \cdot f_{n-1}(a) + \partial_B(b)).$$

- (1) Prove that this is indeed a chain complex.
- (2) Construct chain maps $B \rightarrow \text{Cone}(f)$ and $\text{Cone}(f) \rightarrow A[1]$, where $A[1]$ simply means the chain complex A with the modified grading $A[1]_n = A_{n-1}$, and show that you have constructed a short exact sequence

$$0 \rightarrow B \longrightarrow \text{Cone}(f) \longrightarrow A[1] \rightarrow 0.$$

- (ii) Now define a chain complex $\text{Cyl}(f)$ by $\text{Cyl}(f)_n = A_{n-1} \oplus B_n \oplus A_n$ with differential $\partial: \text{Cyl}(f)_n \rightarrow \text{Cyl}(f)_{n-1}$ given in block form by the matrix

$$\begin{pmatrix} \partial_A & 0 & 0 \\ (-1)^n \cdot f_{n-1} & \partial_B & 0 \\ (-1)^{n+1} \cdot \text{id} & 0 & \partial_A \end{pmatrix}.$$


- (3) Prove that this is a chain complex.
- (4) Construct a chain homotopy equivalence $\text{Cyl}(f) \simeq B$.

Exercise 8.3 (Cones of continuous maps and dunce caps.)

Let Z be a space with subspace $A \subseteq Z$ and let $f: A \rightarrow Y$ be a continuous map. Recall (cf. Exercise 5.6) that the space $Z \cup_f Y$ is defined to be the quotient of the disjoint union $Z \sqcup Y$ by the smallest equivalence relation \sim such that $a \sim f(a)$ for all $a \in A$.


Now let $g: X \rightarrow Y$ be a continuous map and define its *mapping cylinder* to be $\text{Cyl}(g) = Z \cup_f Y$ where $Z = X \times [0, 1]$, $A = X \times \{0\}$ and f is g composed with the obvious identification $X \times \{0\} \cong X$. Define its *mapping cone* to be $\text{Cone}(g) = \text{Cyl}(g)/\sim$, where \sim is the smallest equivalence relation such that $(x, 1) \sim (x', 1)$ for all $x, x' \in X$.

- (1) Draw a picture to show what is going on geometrically in these constructions.
- (2) Construct an embedding $Y \rightarrow \text{Cone}(g)$ and a projection $\text{Cone}(g) \rightarrow \Sigma X$, where ΣX is the *suspension* of X , defined to be $X \times [0, 1]/\sim$, where \sim is the smallest equivalence relation such that $(x, 1) \sim (x', 1)$ and $(x, 0) \sim (x', 0)$ for all $x, x' \in X$.
- (3) Show that $\text{Cyl}(g)$ is homotopy equivalent to Y .
(We will see later in the course that these constructions yield those of the previous exercise after applying the singular chain functor.)
- (4) Now let $A \subseteq X$ be a closed subspace for which there exists an open neighbourhood $U \supseteq A$ that deformation retracts onto A . Let $f, g: A \rightarrow Y$ be two continuous maps which are homotopic. Prove that $X \cup_f Y$ and $X \cup_g Y$ are homotopy equivalent. Thus the operations $\text{Cyl}(-)$ and $\text{Cone}(-)$ are homotopy invariant.
- (5) Thus show that the following two spaces are contractible:



(Hint: realise each of them as $\mathbb{D}^2 \cup_f \mathbb{S}^1$ for some map $f: \partial\mathbb{D}^2 \rightarrow \mathbb{S}^1$, and consider the degree of this map.)
The left-hand space above is often called the “dunce cap”. There are many *generalised dunce caps* like the right-hand space above – each of them is the quotient of a polygon with an odd number of sides, which are all identified with certain choices of orientations.

- (6) Using a similar trick to above, show that the following two spaces each have fundamental group isomorphic to \mathbb{Z} , and draw a generator in each case:
(Note: in the left-hand space on the next page, in addition to the depicted identifications of edges, we also identify all four (not just two) vertices to a single point.)



Exercise 8.4 (Mapping tori of chain complexes.)

Let R be a ring, C a chain complex of R -modules and $f: C \rightarrow C$ be a chain map from C to itself. We can formally adjoin an invertible indeterminate t to C to obtain a chain complex \bar{C} of $R[t^{\pm 1}]$ -modules by first setting $\bar{C}_n = C_n \otimes_R R[t^{\pm 1}]$ and then defining $\bar{\partial}$ to be ∂ extended by linearity in t (more formally: $\bar{\partial} = \partial \otimes \text{id}$, where id is the identity map $R[t^{\pm 1}] \rightarrow R[t^{\pm 1}]$). Here, $R[t^{\pm 1}]$ is the ring of *Laurent polynomials* in t with coefficients in R , or, equivalently, the *group-ring* $R[\mathbb{Z}]$ of the group \mathbb{Z} with coefficients in R . The chain map f extends by linearity in t to a chain map $\bar{f}: \bar{C} \rightarrow \bar{C}$. There is also a canonical chain map $t: \bar{C} \rightarrow \bar{C}$ where each $t_n: \bar{C}_n \rightarrow \bar{C}_n$ is just multiplication by t . Define:

$$\text{Torus}(f) = \text{Cone}(\bar{f} - t).$$

- (1) Describe this explicitly in terms of the C_n , ∂_C and f_n .
- (2) Suppose that we have a commutative square

$$\begin{array}{ccc} C & \xrightarrow{f} & C \\ \alpha \downarrow & & \downarrow \alpha \\ D & \xrightarrow{g} & D \end{array}$$

Define a chain map $(\alpha, \alpha)_{\#}: \text{Torus}(f) \rightarrow \text{Torus}(g)$ and show that your construction satisfies the two functoriality properties $(\text{id}, \text{id})_{\#} = \text{id}$ and $(\alpha, \alpha)_{\#} \circ (\alpha', \alpha')_{\#} = (\alpha \circ \alpha', \alpha \circ \alpha')_{\#}$.

- (3) Show that, for any chain map $f: C \rightarrow C$, the chain map $(f, f)_{\#}: \text{Torus}(f) \rightarrow \text{Torus}(f)$ induces isomorphisms on all homology groups.

(Hint: construct a chain homotopy from $(f, f)_{\#}$ to the “multiplication by t ” chain map from $\text{Torus}(f)$ to itself. Then show that this “multiplication by t ” chain map induces isomorphisms on all homology groups and use the homotopy-invariance property of homology to deduce that the same is true for $(f, f)_{\#}$.)

- (4) Deduce that, for chain maps $f: C \rightarrow D$ and $g: D \rightarrow C$, the chain complexes $\text{Torus}(f \circ g)$ and $\text{Torus}(g \circ f)$ have the same homology groups.

(Hint: consider the chain maps $(f \circ g, f \circ g)_{\#}$ and $(g \circ f, g \circ f)_{\#}$.)

The 6-point space Σ^2 “would be homeomorphic to a 2-sphere if it were only Hausdorff.” More precisely, consider the following conditions on a topological space X : (1) *The complement of each point in X is acyclic (in singular homology)*; (2) $H_2(X) \neq 0$. We have seen that the T_0 space Σ^2 satisfies these two conditions. However, simply by adding the extra condition (3) *X is Hausdorff*, one can conclude that X is homeomorphic to the 2-sphere. (See [5].)

From *Singular homology groups and homotopy groups of finite topological spaces* by M. C. McCord (1966). In his notation, Σ^2 is the 6-point space considered in Exercise 8.5(c) on the next page. Thus you have a strong hint as to what the homology of that space “should” be!

Exercise 8.5 (Finite topological spaces.)

(a) There are three topological spaces X having exactly two points. In each case, compute the singular chain complex $S_\bullet(X)$ and the homology $H_n(X)$ for all n .

(b) Consider the 4-point topological space $\{a, b, c, d\}$ whose topology is generated by the base

$$\{a\}; \{b\}; \{a, b, c\}; \{a, b, d\}$$

and calculate its homology.

(c)* Do the same for the 6-point space $\{a, b, c, d, e, f\}$ whose topology is generated by the base

$$\{a\}; \{b\}; \{a, b, c\}; \{a, b, d\}; \{a, b, c, d, e\}; \{a, b, c, d, f\}.$$

(d)* In general, there is a $(2n + 2)$ -point space $\{a_1, b_1, \dots, a_{n+1}, b_{n+1}\}$ whose topology is generated by the base

$$\{a_1\}; \{b_1\}; \{a_1, b_1, a_2\}; \{a_1, b_1, b_2\}; \dots \dots ; \{a_1, b_1, \dots, a_n, b_n, a_{n+1}\}; \{a_1, b_1, \dots, a_n, b_n, b_{n+1}\}.$$

Make a conjecture about its homology, and about which (more familiar!) space it is homotopy equivalent to.

(e)** Prove your conjecture.

Exercise 8.6* (A chain complex of chain maps.)

Let C and D be chain complexes of R -modules. We define a *chain map of degree d* to be a collection $f = \{f_n\}_{n \in \mathbb{Z}}$ of homomorphisms of R -modules $f_n: C_n \rightarrow D_{n+d}$ such that $\partial_{n+d}^D \circ f_n = f_{n-1} \circ \partial_n^C$ for all n . We define a *pre-chain map of degree d* to be simply a collection $f = \{f_n\}_{n \in \mathbb{Z}}$ of homomorphisms $f_n: C_n \rightarrow D_{n+d}$, with no condition.

(1) Show that the set of all pre-chain maps of a fixed degree d forms an R -module, denoted $\text{PreChain}_d(C, D)$.

(2) Given $f = \{f_n\} \in \text{PreChain}_d(C, D)$, show that the formula

$$(df)_n = \partial_{n+d}^D \circ f_n - (-1)^d f_{n-1} \circ \partial_n^C$$

defines a pre-chain map $df \in \text{PreChain}_{d-1}(C, D)$.

(3) Show that $ddf = 0$, and hence that $\text{PreChain}_\bullet(C, D)$ is a chain complex.

(4) Prove that there is a natural isomorphism between $H_0(\text{PreChain}_\bullet(C, D))$ and the set of chain-homotopy-classes of chain maps (of degree 0) from C to D . (Hint: first note that $Z_0(\text{PreChain}_\bullet(C, D))$ is naturally isomorphic to the set of chain maps of degree 0.)